AnnAGNPS Version 5.51:

Input File Specifications for CSV-formatted Input Data

December 4, 2019

Input Specification Document for AnnAGNPS

USER RESPONSIBILITY	1
ANNAGNPS OPERATION MODES	1
INPUT FILES FOR ANNAGNPS	1
Control file: "AnnAGNPS.fil" Watershed input file(s): Daily Climate input file:	
Storm Type input file: RUSLE2 CSV-Formatted Erosion Input file(s) (.CSV):	
TABLE LAYOUT INFORMATION	
Data Parameter Definition Table	
Layout Matrix Table	
Input Parameter Definition: AnnAGNPS.fil	
Input File Layout Matrix: AnnAGNPS.fil	
ANNAGNPS: CSV-FORMATTED INPUT FILE DESCRIPTIONS	5
Data Section Requirements	
AnnAGNPS ID	6
Aquaculture Pond Data	7
Aquaculture Pond Management Schedule Data	
Cell Data	
Classic Gully Data	
Contour Data	
Crop Data	
Crop Growth Data	
EI Percentage Data	
Ephemeral Gully Data	
Feedlot Data	
Feedlot Management Data	
Fertilizer Application Data	
Fertilizer Reference Data	
Field Pond Data	
Field Pond Operations Data	
Geology Data	
Global Error and Warning Limits Data	
Global IDs, Factors, and Flags Data:	
Hydraulic Geometry Data (built-in)	
Hydraulic Geometry Data (user-defined in input)	
Impoundment Data	
Irrigation Application Data	
Management Field Data	
Management Operation Data	
Management Schedule Data	
Non-Crop Data	
Output Options Data – Global	
Output Options Data – AA	
Output Options Data – CSV	
Output Options Data – DPP	
Output Options Data – EV	
Output Options Data – NPT	
Output Options Data – SIM	
Output Options Data – TBL	
Output Options Data – Min/Max (Limits)	
Output Options Data – Cell	

Output Options Data – Feedlot	
Output Options Data – Field Pond	
Output Options Data – Gully, Classic	
Output Options Data – Gully, Ephemeral	
Output Options Data – Impoundment	
Output Options Data – Point Source	
Output Options Data – Reach	
Output Options Data – Wetland	
Pesticide Application Data	
Pesticide Initial Conditions Data	
Pesticide Reference Data	
PL Calibration Data	
Point Source Data	
RCN Calibration Data:	
Reach Data	
Reach Nutrient Half-life	
Riparian Buffer Data	
Runoff Curve Number Data (built-in)	
Runoff Curve Number Data	
RUSLE2 Data	
Simulation Period Data	
Soil Data	
Soil Initial Conditions Data	
Soil Layer Data	
Strip Crop Data	
Tile Drain Data	
Watershed Data	
Wetland Data	
INPUT FILE LAYOUT MATRIX	
AnnAGNPS ID	
Aquaculture Pond Data	
Aquaculture Pond Management Schedule Data	
Cell Data	
Classic Gully Data	
Contour Data	
Crop Data	
Crop Growth Data	
EI Percentage Data	
Ephemeral Gully Data	
Feedlot Data	
Feedlot Management Data	
Fertilizer Application Data	
Fertilizer Reference Data	
Field Pond Data	
Field Pond Operation Data	
Geology Data	
Global Error and Warning Limits Data	
Global IDs, Factors, and Flags Data	
Hydraulic Geometry Data (User Defined)	
Impoundment Data	
Irrigation Application Data	
Management Field Data	
Management Operation Data	
Management Schedule Data	
Non-Crop Data	
Output Options Data – Global	
Output Options Data – AA	
Output Options Data – CSV	100

Output Options Data – DPP	
Output Options Data – EV	
Output Options Data – NPT	
Output Options Data – SIM	
Output Options Data – TBL	
Output Options Data – Min/Max (Limits)	
Output Options Data - Cell	
Output Options Data – Feedlot	
Output Options Data – Field Pond	
Output Options Data – Gully, Classic	
Output Options Data – Gully, Ephemeral	
Output Options Data – Impoundment	
Output Options Data – Point Source	
Output Options Data – Reach	
Output Options Data – Wetland	
Pesticide Application Data	
Pesticide Initial Conditions Data	
Pesticide Reference Data	
PL Calibration Data:	
Point Source Data	
RCN Calibration Data	
Reach Data	
Reach Nutrient Half-life	
Riparian Buffer Data	
Runoff Curve Number Data	
RUSLE2 Data	
Simulation Period Data	
Soil Data	
Soil Initial Conditions Data	
Soil Laver Data	
Strip Crop Data	
Tile Drain Data	120
Watershed Data	120
Wetland Data	
	100
DAILY CLIMATE DATA	
Climate Station Information	
Daily Climate Information	
Climate Input File Lavout Matrix – Station	
Climate Input File Layout Matrix – Daily	
STORM TYPE DATA	
Rainfall Distribution	129
Unit Peak Discharge Regression Coefficients	129
Storm Type Input File Layout Matrix – Rainfall Distribution	130
Storm Type Input File Layout Matrix – Unit Peak Discharge Regression Coefficients	130
Storm Type Input The Edybut Mutrix Only Teak Discharge Regression Coefficients	150
APPENDIX A: OUTPUT FILES	
VERSIONS 1 & 2 OUTPUT-RELATED OPTIONS	
OUTPUT FILE STRUCTURE	131
Database. Text. & Verification Files	131
Reach Rules	131
Cell Rules	
Subarea Rules	
Minimum/Maximum Event Date Rules	
Minimum Fyont Runoff Rules	
Database (* csv) Files	

Text (*.txt) Files	
Verification (*.dpp, *.npt, & *.sim) Files	
OUTPUT FILE NAMES & INDICES	

User Responsibility

Results from the model can be extremely sensitive to the input data. It is the user's responsibility to ensure that all input data (whether it is actually entered or implied through default values for blank data fields) is appropriate for the watershed and watershed conditions being analyzed. Operations data is of particularly importance as it defines the changes that occur in the watershed over the simulation period.

AnnAGNPS Operation Modes

AnnAGNPS can now be operated in only one mode. The standard mode (AnnAGNPS) allows for continuous simulation of a watershed using a daily time step. The second mode, previously available, called the AGNPS mode is no longer supported because it can be duplicated by the AnnAGNPS mode which can be used for a simulation period of one day.

Input Files for AnnAGNPS

These specifications apply to AnnAGNPS version 5.51 csv-formatted input files. Previous versions of input specifications will apply to associated input files of those previous versions. It is recommended that the input files be brought up to the current version of AnnAGNPS.

There are four possible user-input file types to AnnAGNPS. Of the four, two are required and two are optional. AnnAGNPS requires watershed input data and daily climate input data. The user may optionally supply a control file or storm type input data. The input data types are described in more detail below.

- 1.) Control file optional
- 2.) Watershed input data required
- 3.) Daily climate input data required
- 4.) Storm type input data optional

There are currently two input formats that AnnAGNPS recognizes for watershed data, climate data, and storm type data. One format is Fixed-Formatted Data (FFD) and the other is CSV-formatted data (CSV). The following is a brief description of the two formats:

- 1.) Fixed-formatted Data (FFD) FFD is a text-based input format that has all data contained within fields of records made up of a fixed starting and ending columns. This is a very rigid format and all data must fall correctly within the starting and ending column as specified in the input specifications document "*Input_Specifications.PDF*".
- 2.) CSV-formatted Data (CSV) CSV is a format where commas are used to separate the data values. Watershed data in this format can be created and modified in common applications such as Excel and then used as input directly into AnnAGNPS. Data fields (or columns in Excel) must still be in a certain order as specified in this input specifications document.

Note: This document is only concerned with the CSV-formatted data. For information on the FFD-formatted data, please view the *"Input_Specifications.PDF"* document.

Note: For a detailed explanation of input files and formats, please view the "AnnAGNPS_User's_Guide_for_Input_Files_&_Formats.pdf" document.

CONTROL FILE: "ANNAGNPS.FIL"

AnnAGNPS has an optional FFD control file that can be used to specify the path and filename of watershed data, climate data, and storm type data. The name of the control file is required to be "AnnAGNPS.fil".

If "AnnAGNPS.fil" is not used, then two input files are required and must be in FFD format. The watershed input filename must be "AnnAGNPS.inp" and the climate filename must be "DayClim.inp". If the optional storm type input file is to be used, then the storm type filename must be "Storm_Type.inp". The tables in this document give detailed information as to the layout of the data for these three inputs.

If "AnnAGNPS.fil" is used, then AnnAGNPS reads this file first to determine the path and filename of the watershed data, climate data, and optionally, the storm type data.

"AnnAGNPS.fil" consists of three records:

- The first record is the path and filename of the CSV master list file which will be briefly described later in this document. A more detailed explanation can be found in the "AnnAGNPS_User's_Guide_for_Input_Files_&_Formats.pdf" document. The extension of the CSV master list file must be ".csv". If this record is blank then AnnAGNPS assumes the FFD default watershed input file name of "AnnAGNPS.inp" and expects all input data to be in FFD format.
- 2.) The second record is the path and filename of the FFD climate file. If the watershed input file is in CSV format, then the climate data may also be in CSV format and its path and filenames specified in the csv master list as described below. If the watershed input file is in FFD format, then the climate data must also be in FFD format. If the climate data is in FFD format and this record is blank then AnnAGNPS assumes the default climate input file name of "DayClim.inp".
- 3.) The third record is the path and filename of the optional storm type file. If the watershed input file is in CSV format, then the storm type data may also be in CSV format and its path and filenames specified in the csv master list as described below. If the watershed input file is in FFD format, then the storm type data must also be in FFD format. If the storm type data is in FFD format and this record is blank then AnnAGNPS assumes the default storm type input file name of "Storm_Type.inp".

The file path is optional and if omitted, then AnnAGNPS expects the input files to be in the same folder as the AnnAGNPS executable file.

WATERSHED INPUT FILE(S):

Since the inception of AnnAGNPS, the watershed input file has been required. The watershed input file contains all of the data that describes the watershed. It also contains control information used to specify various parameters such as the period of simulation along with any initialization period and initial conditions. If the control file "AnnAGNPS.fil" is not used then the watershed input filename must be "AnnAGNPS.inp" and uses the FFD format.

AnnAGNPS accepts the watershed input data in one of two formats; fixed-formatted (FFD) and csv-formatted (CSV). This document is only for the csv-formatted input data.

CSV formatted data – this format requires multiple CSV input files. Typically, there is one CSV input file per data section as described in this document. There are a few sections where a second CSV input file may be required. For example, soils require two csv input files; one describes the soil and the other contains the soil layer information. The first record of each CSV input file is a header record that describes the parameter expected for that column. The proceeding record(s) in the input file will contain the appropriate data.

A CSV formatted master list file is needed so that AnnAGNPS can determine the path and filename of the CSV input files to be used. There is no required data section order in the master list because AnnAGNPS will read the master list and sort it into the required order before attempting to read the data. Also, there is no requirement on the name of the master list except that it must have the ".csv" extension and be a CSV formatted file. The user would use this master list filename on the first record of the "AnnAGNPS.fil" control file to indicate to AnnAGNPS that the watershed inputs are CSV formatted input files.

The "AnnAGNPS_User's_Guide_for_Input_Files_&_Formats.pdf" document contains a more detailed explanation of the csv master list file. A table giving a brief example is included in this document below.

DAILY CLIMATE INPUT FILE:

Daily climate information must be supplied covering the entire period of simulation. A primary climate dataset is required for AnnAGNPS. If the watershed input file is FFD then the climate file must also be FFD. If the watershed inputs are from CSV files as described above then the climate data may be FFD or CSV files that contain daily climate information for the watershed. The contents, layout, and descriptions are contained in the "*Input_Specifications.PDF*" documentation for FFD and this document for CSV. If the control file "AnnAGNPS.fil" is not used then the climate data must be FFD and the filename must be "DayClim.inp".

If the climate data is in CSV format, then two entries are required in the master list:

- 1.) "Climate Data Daily" the primary station's daily climate records (required)
- 2.) "Climate Data Station" the primary station's information (required)

STORM TYPE INPUT FILE:

An optional storm type input file may be used. This file allows for user-customized storm type input data. If the watershed input file is FFD then the storm type file must also be FFD. If the watershed inputs are from CSV files as described above then the storm type data may be FFD or CSV files that contain storm type information for the watershed. The contents, layout, and descriptions are contained in the "*Input_Specifications.PDF*" documentation for FFD and this document for CSV. If the control file is not used then the storm type data must be FFD and the filename must be "Storm Type.inp".

If the storm type data is in CSV format, then two entries are possible in the master list:

- 1.) "Storm Type Data RFD" the rainfall distribution file (required)
- 2.) "Storm Type Data UPDRC" the unit peak discharge regression coefficients (optional)

RUSLE2 CSV-FORMATTED EROSION INPUT FILE(S) (.CSV):

Starting with AnnAGNPS v5.50, RUSLE2 erosion values and particle size distributions may be inputted directly via a csvformatted input file(s). The input filenames will be specified in the RUSLE2 Data section as described further in this document. The user will specify whether RUSLE2 erosion is to be used or if AnnAGNPS will calculate erosion using RUSLE as historically done.

The user may specify input parameters within the "Global Ids, Factors, and Flags Data", "Cell Data", and "RUSLE2 Data" sections to instruct AnnAGNPS to use RUSLE2 erosion values retrieved from the file(s) the user provides. There are also two output options data sections that have parameters for reporting RUSLE2 information: "OUTPUT OPTIONS – DPP" AND "OUTPUT OPTIONS – NPT".

The input file must adhere to the following format where the first record is the required header line and records 2 - n are the data records:

Date	Erosion [tns/ac]	Clay [%]	Silt [%]	Sand [%]	Small Aggregate [%]	Large Aggregate [%]
01/01/1982	0.0	5.2003	23.9975	6.5531	36.0018	28.2473
01/02/1982	1.9					
12/31/1982	0.0					

The date and erosion value are required as columns 1 and 2. However, the particle size distributions are optional. If present, only the first data record needs to be populated as shown above and total 100%. These values are used for each day of the simulation. If the distribution values are not present as shown in the table below, then AnnAGNPS will determine the particle size distributions.

Date	Erosion [tns/ac]
01/01/1982	0.0
01/02/1982	1.9
12/31/1982	0.0

Table Layout Information

There are two informational table layouts used in this document; 1.) data parameter definitions; and 2.) layout matrix.

DATA PARAMETER DEFINITION TABLE

The first portion of this document contains tables with the following information:

Description	Field	Units	Domain	Format	Record	Field
	Header for	{English}	{English}		No.	No.
	Record #1	[SI]	[SI]			

Description – describes the data parameter.

Field Header for Record #1 – every csv-formatted input file for AnnAGNPS requires a header row. Rows are commonly called "records" and is the term used throughout this document. The header record contains the column headings, commonly called "fields"

as used throughout this document, for each data parameter. This header record is always located as the first record in the csv input file. This column in the table shows the required header information expected by AnnAGNPS.

Units {English} [SI] – This column shows the units that are expected for each of the data parameters.

Domain {English} [SI] - This column shows the range of values that are acceptable to AnnAGNPS for each data parameter.

Format – This column shows the acceptable format of data that is acceptable to AnnAGNPS for each data parameter.

Record No. – There are only two records defined in the tables. Record one is always the header record. All data begins on record two and repeats as needed for remaining data. For example, a watershed data set may have 100 cells. The csv input file will have the column headings on record one. Records 2 - 101 will contain all cell information. Each record contains information for one cell. **Field No.** – This column contains the numerical column index of each data parameter.

LAYOUT MATRIX TABLE

The second portion of the document contains tables with the following information:

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field m
1	Cell_ID	Soil_ID	Mgmt_Field_ID	Reach_ID	Reach_Location_C ode	Cell_Area	
2							

Record – there are two records shown in the layout matrix. Record 1 contains the field (column) headings for each of the data parameters in the data section as required by AnnAGNPS. This header record is required for each data section. Record two is left blank in this document but illustrates that data values are stored in records 2 - n, where n is the number of data records that may exist in a data section. For example, if there are 100 cells represented in the cell data section then record 1 contains the header information and records 2 - n contain the data; one record per cell.

Field – the number of fields (columns) depends on the number of data parameters in the data section. There is one field per data parameter. For example, the cell data section currently contains 34 data parameters; therefore, there will be 34 fields per record.

INPUT PARAMETER	DEFINITION:	ANNAGNPS.FIL
-----------------	--------------------	--------------

	Description	Domain	Format	Record No.	Field No.
AnnAGNPS Input File N Input file If blank the def	ame—Path and file name to the AnnAGNPS	Computer platform	A80	1	1
filename and all input data	is expected in fixed format (FFD).	filename			
If a filename is entered in t that it is a csv file that cont AnnAGNPS. The AnnAG into separate csv files and default filename of "annag of two columns of data; 1.) defined for each data section path and filename (column following table is an examp	his field with a .csv extension, then it is assumed ains the master list of csv input files for NPS Input Editor is able to export watershed data will create the related master list csv file with a nps_master.csv. The csv master list is comprised Data Section ID (column 1) and must be exact as on described in this documentation; 2.) The file 2). The first row is a required header row. The ole:				
Data Section ID	File Path				
AnnAGNPS ID	c:\~agedit\simulation\annaid.csv				
Cell Data					
Crop Data					
Simulation Period Data	c:\~agedit\simulation\sim_period.csv				
Watershed Data	c:\~agedit\watershed\watershed_data.csv				
Wetland Data	c:\~agedit\watershed\wetland.csv				

Description	Domain	Format	Record No.	Field No.
Climate Input File Name—For climate data in CSV format, this field should be left blank and entries made in the csv master list file for the climate data. There are two required csv-formatted climate files needed; one for climate station information and the other for daily climate data. These are described later in this document. Secondary climate files may be used in AnnAGNPS. The secondary climate filenames do not go into the "AnnAGNPS.fil" file. The secondary climate files are specified in the CELL DATA section in the field labeled "Climate file ID:" and must have the same 'root' as the primary climate file. For example, if the primary climate filename is "climate_daily.csv" as specified in the csv master list file, then the related secondary files must have the same 'root' name of "climate_daily" followed with the underscore character "_" and then a user-specified appendage such as "01". For example: Primary climate: "climate_daily.csv" (Specified in the csv master list file.) Secondary filenames: "climate_daily_01.inp" "climate_daily_02.inp" "climate_daily_03.inp" "climate_daily_05.inp" "climate_daily_06.inp" Then the individual cells in the Cell Data section for which a secondary climate file is to be used would specify the secondary climate ID appendage	Computer platform acceptable path and filename	A80	2	1
of "01", "02", "03", "04", "05", or "06".				
Storm Type Data Input File Name — For storm type data in CSV format, this field should be left blank and entries made in the csv master list file for the storm type data. There are two possible csv-formatted storm type files; one for the rainfall distribution and the other for unit peak discharge regression coefficients data. These are described later in this document.	Computer platform acceptable path and filename	A80	3	1
Blank Line			Last	

INPUT FILE LAYOUT MATRIX: ANNAGNPS.FIL

Data Field 1	Data Field 2	Data Field 3	Data Field 4	Data Field 5	Data Field 6	Data Field 7	Data Field 8	
AnnAGNPS csv master list input file name								
	Daily Climate Data input file name (leave blank if csv inputs are used for climate data)							
Storm Type Data input filename (leave blank if csv inputs are used for storm type data)								

AnnAGNPS: CSV-formatted Input File Descriptions

DATA SECTION REQUIREMENTS

	Data Section Heading	Required	Optio	nal	Referenced by Data Section(s)	
			Independent	Dependent		
1.	ANNAGNPS ID*	Х				
2.	WATERSHED DATA	Х				
3.	SIMULATION PERIOD DATA	Х				
4.	GLOBAL ERROR AND WARNING LIMITS DATA		Х			
5.	GLOBAL IDS, FACTORS, AND FLAGS DATA		Х			
6.	SOIL INITIAL CONDITIONS DATA			Х	"SOIL DATA"	
7.	PESTICIDE INITIAL DATA		Х			
8.	PL CALIBRATION DATA		Х			

	Data Soction Heading	Poquirod	Optio	nal	Pafarancad by Data Soction(s)
	Data Section Heading	Keyun cu	Independent	Dependent	Referenced by Data Section(s)
9.	RCN CALIBRATION DATA		-	X	"CELL DATA"
10.	AOUACULTURE POND DATA		x		
	AOUACULTURE POND MANAGEMENT				
11.	SCHEDULE DATA			Х	"AQUACULTURE POND DATA"
12.	CELL DATA	х			"AQUACULTURE POND DATA", "CLASSIC GULLY DATA", "EPHEMERAL GULLY DATA", "FEEDLOT DATA", "FIELD POND DATA", "LANDSLIDE DATA", "POINT SOURCE DATA", "RIPARIAN BUFFER DATA"
13.	CLASSIC GULLY DATA		Х		
14.	CONTOUR DATA			X	"MANAGEMENT SCHEDULE DATA"
15.	CROP DATA			X	"MANAGEMENT SCHEDULE DATA"
16.	EPHEMERAL GULLY DATA		Х		
17.	FEEDLOT DATA		Х		
18.	FEEDLOT MANAGEMENT DATA			Х	"FEEDLOT DATA"
19.	FERTILIZER APPLICATION DATA			Х	"MANAGEMENT SCHEDULE DATA"
20.	FERTILIZER REFERENCE DATA			Х	"FERTILIZER APPLICATION DATA"
21.	FIELD POND DATA		Х		
22.	GEOLOGY DATA			Х	"CELL DATA", "GLOBAL IDS, FACTORS, AND FLAGS DATA"
23.	HYDRAULIC GEOMETRY DATA			х	"CELL DATA", "EPHEMERAL GULLY DATA", "GLOBAL IDS, FACTORS, AND FLAGS DATA", "REACH DATA"
24.	IMPOUNDMENT DATA		Х		
25.	IRRIGATION APPLICATION DATA			Х	"MANAGEMENT SCHEDULE DATA"
26.	LANDSLIDE DATA		Х		
27.	MANAGEMENT FIELD DATA	Х			"CLASSIC GULLY DATA", "EPHEMERAL GULLY DATA", "LANDSLIDE DATA"
28.	MANAGEMENT OPERATION DATA			Х	"MANAGEMENT SCHEDULE DATA"
29.	MANAGEMENT SCHEDULE DATA	Х			"MANAGEMENT FIELD DATA"
30.	NON-CROP DATA			Х	"MANAGEMENT SCHEDULE DATA"
31.	PESTICIDE APPLICATION DATA			X	"MANAGEMENT SCHEDULE DATA", "SIMULATION PERIOD DATA"
32.	PESTICIDE REFERENCE DATA			Х	"PESTICIDE APPLICATION DATA"
33.	POINT SOURCE DATA		Х		
34.	REACH DATA	х			"CELL DATA", "CLASSIC GULLY DATA", "EPHEMERAL GULLY DATA", "IMPOUNDMENT DATA"
35.	REACH NUTRIENT HALF-LIFE		Х		
36.	RIPARIAN BUFFER DATA		Х		
37.	RUNOFF CURVE NUMBER DATA	Х			"MANAGEMENT SCHEDULE DATA"
38.	RUSLE2 DATA			Х	"CELL DATA", "GLOBAL IDS, FACTORS, AND FLAGS DATA"
39.	SOIL DATA	Х			"CELL DATA", "CLASSIC GULLY DATA", "EPHEMERAL GULLY DATA", "LANDSLIDE DATA"
40.	STRIP CROP DATA			Х	"MANAGEMENT SCHEDULE DATA"
41.	TILE DRAIN DATA			Х	"MANAGEMENT FIELD DATA"
42.	WETLAND DATA		Х		
43.	OUTPUT OPTIONS DATA		Х		
44.	END DATA	Х			

Go to Table of Contents

ANNAGNPS ID

Required

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field as listed below.				Α	1	1-5

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Watershed File Input Version ID—Unique alphanumeric string identifying the watershed file's input.	"Version"		5.51	A4	2	1
Input Units code —Code identifying whether input is in English or metric units. Acceptable values are: 0 = English, 1 = SI (Blank indicates 0)	"Input_Units"		Blank, 0 or 1	I1	2	2
Output Units code —Code identifying whether output is in English or metric units. Acceptable values are: 0 = English, 1 = SI (Blank indicates 0)	"Output_Units"		Blank, 0 or 1	I1	2	3
CCHE1D Output Units code —Code used to identify output units for output file to be used with CCHE1D file. Acceptable values are: 0 = English, 1 = SI (Blank indicates no CCHE1D output desired)	"CCHE1D_Output _Units"		Blank, 0 or 1	I1	2	4
Screen Output code—Code indicating whether screen output is desired. To be used when AnnAGNPS is embedded within an preprocess/post- process code. Leave blank when directly running AnnAGNPS. Acceptable codes are: 0 = Screen output $1 = $ No screen output. (Blank indicates 1)	"Screen_Output_U nits"		Blank, 0 or 1	I1	2	5
Go to Layout Matrix		Go to	Table of Cont	tents		

AQUACULTURE POND DATA Optional

Description	Field Header for	Units {English}	Domain {English} [SI]	For mat	Record	Field No
	Record #1	[SI]		mat	110.	110.
Field Header— Required unique field header for each field listed				А	1	1-13
below.				_		
The following record repeats for t	the number of	Aquacultur	e ponds.			
Pond identifier —unique alphanumeric string identifying the aquaculture pond. Multiple aquaculture ponds within the same cell may be aggregated and entered as a single pond for simulation convenience.	"Pond_ID"			A10	2	1
Pond-Cell identifier —alphanumeric string identifying cell that contains the aquaculture pond(s). Must be the same as a cell identifier in the CELL DATA section already included within the watershed.	"Cell_ID"			A10	2	2
Pond area —area of aquaculture pond(s). Multiple aquaculture ponds in the same cell may be aggregated together as a single aquaculture pond for convenience. Defaults to cell area.	"Pond_Area"	{acres} [hectares]	Blank, {0.0—10000.0} [0.0—4000.0]	F10	2	3
Pond depth —Maximum depth of water in the aquaculture pond	"Pond_Depth"	{in} [mm]	{0.0 to 393.72} [0.0 to 10000.0]	F10	2	4
Seepage Rate Daily water loss due to seepage. If left blank, the seepage loss will be 0.	"Seepage_Rate"	{in/day} [mm/day]	Blank, {0.0 to 393.72} [0.0 to 10000.0]	F10	2	5
Sediment Delivery Ratio—Fraction of pond discharge delivered to the receiving reach. If left blank, the value is assumed to be 1.0.	"Sediment_Deli very_Ratio"		Blank, 0.0 – 1.0	F10	2	6
Relative Rotation Year —Relative year in the aquaculture pond management rotation for starting simulation for this aquaculture pond. (Example: For a 3 year rotation, starting year can be 1, 2 or 3). Blank defaults to 1.	"Relative_Rotati on_Year"		Blank, or 1 to 100	I10	2	7
Pond Management Schedule Identifier —alphanumeric string identifying the Aquaculture Pond Management Schedule for this aquaculture pond.	"Mgmt_Schd_I D"			A10	2	8
Organic Carbon Calibration Factor—used to calibrate aquaculture pond organic carbon for this pond only. Defaults to the watershed-scale organic carbon from pond sources calibration factor in the PL CALIBRATION DATA section.	"OC_Calib_Fctr "	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	9

AnnAGNPS Input File Specifications

AnnAON 5 mput Fit Specifications									
Revision: December 4, 2019 File	e Name: Input_Specifica	tions_v5.51_2	019.xx.xx.docx						
	T. 11	TT •/	D •	E D	D				

Description	Field Header for	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.	
	Record #1	[SI]					
Nitrogen Calibration Factor —used to calibrate aquaculture pond nitrogen for this pond only. Defaults to the watershed-scale nitrogen from pond sources calibration factor in the PL CALIBRATION DATA section.	"N_Calib_Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	10	
Phosphorus Calibration Factor —used to calibrate aquaculture pond phosphorus for this pond only. Defaults to the watershed-scale phosphorus from pond sources calibration factor in the PL CALIBRATION DATA section.	"P_Calib_Fctr" "	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	11	
Erosion Calibration Factor —used to calibrate aquaculture pond erosion for this pond only. Defaults to the watershed-scale sediment from pond sources calibration factor in the PL CALIBRATION DATA section.	"Erosion_Calib_ Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	12	
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	I1	2	13	
Go to Layout Matrix	Go to Table of Contents						

AQUACULTURE POND MANAGEMENT SCHEDULE DATA

Ontional	unless	referenced	in A augo	ulture Do	nd Data
Obuona	unicos	ICICICICU	i III Auuau		nu Data

	T . 11	TT •/		T	D 1	T. 11		
Description	Field	Units	Domain	For	Record	Field		
	Header for	{English}	{English} [SI]	mat	No.	No.		
	Record #1	[SI]						
Field Header— Required unique field header for each field listed				Α	1	1-45		
below.								
The following record repeats for the number of Aquaculture pond events.								
Aquaculture Pond Management Schedule identifier—unique	"Pond_Mgmt_S			A10	2	1		
alphanumeric string identifying the Aquaculture pond management	chd_ID"							
schedule. Multiple aquaculture ponds within the same cell may be								
aggregated and entered as a single pond for simulation convenience.								
(Note - Required for first event of each schedule; leave blank for								
subsequent events within a schedule.)								
Event Date—Month, day, and relative year (within a set of	"Month"	mm	1-12	I2,	2	2-4		
aquaculture pond operations) water additions and releases. This day	"Day"	dd	1-31	I2,				
will be converted internally from a relative rotation date (mm/dd/ry)	"Year"	vy	Blank, 1-99	I2				
to a 2-dimensional variable which will be the rotation year & Julian			·					
day for the event. Blank year defaults to 1.								
Water Operation code—Code that specifies the type of fill or	"Water_Operati		Blank, 0 or 1	I1	2	5		
release.	on_Code"							
0 = Fill $1 = Initial Drain$ $2 = Midseason Drain$ $3 = Harvest Drain$								
4 = Other release. Options 1,2,3 for rice only.								
Aquaculture Identifier—unique alphanumeric string identifying	"Aquaculture_I		Blank,	A10	2	6		
the aquaculture type. May be entered for the first event in the	D"		"Crawfish"					
schedule or when a change occurs. Currently, only "Crawfish" is								
allowed.								
Crop Identifier—unique alphanumeric string identifying the	"Crop_ID"		Blank, "Rice",	A10	2	7		
crop, if any. Currently only "Rice", "Sorghum", "Native" are			"Sorghum",					
allowed. To be entered on the day the crop is planted.			"Native"					
Planting Type Code – Code that specifies the planting type.	"Planting_Type		Blank, 1-4	I1	2	8		
Allowable	_Code"							
Entries are: 1 – No-till, 2 – Water-seeded with retention, 3 – Clear								
Water, 4 – Traditional water seeding Currently used for rice only.								
Gate Open/Close—Indicator for whether the filed pond gate is	"Gate_Status"		Blank, "Open",	A5	2	9		
closed (pond exists) or open (pond does not exist).			"Close"					

	T * 11	TT • /		T	D 1	T ¹ 1 1
Description	Field	Units	Domain	For	Record	Field
	Header for	{English}	{English} [SI]	mat	No.	No.
	Record #1	[SI]				
Maximum Pool Denth—Upper limit for denth of water in pond If	"Max Pool Dep	{in}	Blank {0.0 to	F10	2	10
the water level exceeds this depth due to filling or precipitation the	th"	[mm]	393 72	110	2	10
excess amount will go into the receiving reach of the cell. This		LIIIII	[0.0 to 10000.0]			
maximum will be maintained until reset or a gate status change			[0.0 10 10000.0]			
maximum will be maintained until reset, of a gate status change						
		(;)	D1 1 (0.0.)	F10	2	11
Minimum Pool Depth—Lower limit for depth of water in pond.	"Min Pool Den	{in}	Blank, $\{0.0 \text{ to}$	F10	2	11
This minimum will be maintained until reset, or a gate status change	th"	[mm]	393.72}			
occurs.			[0.0 to 10000.0]			
Fill/Release Volume—amount of water added to or released from	"Full balance Ma	$\{in\}$	Blank,	F10	2	12
the aquaculture pond on this date; depth in linear units. May be left	"Fill/Kelease_vo	[mm]	{>0. – 99999.}			
blank if release rate is entered.			[>0-4000.]			
Fill/Drain Time—time for the current fill/release. Will be used if	"Fill/Drain_Tim	hr	Blank,	F10	2	13
the fill/release rate field is blank. A blank in both the fill/drain time	e"		> 0.0-48.0			
& fill/ release rate fields will default to a 24-hr fill/drain time.						
Fill/Release Rate—rate of aquaculture pond fill/release as depth in	"Fill/Release Ra	{in/hr}	Blank.	F10	2	14
linear units per hour May be left blank in which case the rate will	te" –	[mm/hr]	$\{>0 - 99999\}$	110	-	
be calculated from the values for the volume of fill/release water		[[>040001]			
(and nond area) and fill/drain time. A blank in both the fill/drain			[² 0. 4000.]			
(and poind area) and inf/drain time. A blank in both the inf/drain time time & fill/release rate fields will default to a 24 hr fill/drain time						
Fill/Drain All and indicating that the need is to be filled the	"Fill/Drain All		Dlank 1	T1	2	15
Fin/Drain An—code indicating that the point is to be fined the	Code"		Dialik, I	11	Z	15
maximum depin or drained to the minimum depin Enter 1 for true,						
or leave blank for false.	"T-t-L S-L C		D1 1	F10		16
Total Sediment Concentration—Concentration of suspended	"Total_Sed_Con	ppm	Blank,	F10	2	16
sediment in the fill/release water			0. – 999999.			
Clay Content—Percentage of suspended sediment that is clay in the	"Clay_Content"	%	Blank,	F10	2	17
fill/release water. Default is 0. for fills. For releases, the value is			0 100.			
internally						
calculated based on pond management, if possible; is 0. otherwise.						
Silt Content—Percentage of suspended sediment that is silt in the	"Silt_Content"	%	Blank,	F10	2	18
fill/release water. Default is 0. for fills. For releases, the value is			0 100.			
internally						
calculated based on pond management, if possible; is 0. otherwise.						
Total Nitrogen—Total concentration of nitrogen in water added to	"Total_N"	ppm	Blank,	F10	2	19
or released from the aquaculture pond. Default is 0. for fills. For		11	0. – 999999.			
releases, the value is internally calculated based on pond						
management, if possible: is 0, otherwise.						
Dissolved Nitrogen —total concentration of dissolved nitrogen in	"Dissolved N"	nnm	Blank.	F10	2	20
water added to or released from the Aquaculture pond. Default is 0.	_	ppm	0 999999	110	-	
for fills. For releases the value is internally calculated based on			0. ,,,,,,,,,			
nond management if possible: is 0 otherwise						
Total Phosphorus—total concentration of Phosphorus in water	"Total P"	nnm	Blank	F10	2	21
added to or released from the Aquaculture pond. Default is 0 for		ppm	0 000000	110	2	21
fills. For releases the value is internally calculated based on pond			0)))))).			
management if possible: is 0, otherwise						
Dissolved Phosphorus concentration of dissolved Dhosphorus in	"Dissolved P"		Dlank	E10	2	22
Dissolved Filosphol us—concentration of dissolved Filosphol is in	Dissorveu_1	ppm		F10	2	22
water added to or released from the Aquaculture point. Default is 0.			0. – 999999.			
for finis. For releases, the value is internally calculated based on						
pond management, il possible; is 0. otnerwise.	When Deed A		D1 1	T10	2	22
Pesticide Applications—Number of pesticide applications	"Num_Pest_Ap		Blank	110	2	23
associated with the event.	1					
Currently not used – Reserved.						
Seasonally Adjust Concentrations—Start/Stop use of seasonal	"Season_Adjust Conc"		Blank, 'Y' or 'N'	A10	2	24
average concentrations.						
Must supply values below, if "Y" is entered. $Y = Yes(Start) N =$						
No(Stop) Blank is no change.	<u> </u>					
Sediment Concentration—Winter	"Sed_Conc_Win	ppm	Blank,	F10	2	25
	ter		0. – 999999.			
Total Nitrogen—Winter	"Total_N_Winte	ppm	Blank,	F10	2	26
	r"		0. – 999999.			

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
	"Dissolved_N_	ppm	Blank.	F10	2	27
	Winter"	11	0. – 999999.	-		
Total Phosphorus—Winter	"Total_P_Winte r"	ppm	Blank, 0. – 9999999.	F10	2	28
Dissolved Phosphorus—Winter	"Dissolved_P_W inter"	ppm	Blank, 0. – 9999999.	F10	2	29
Sediment Concentration—Spring	"Sed_Conc_Spri ng"	ppm	Blank, 0. – 999999.	F10	2	30
Total Nitrogen—Spring	"Total_N_Sprin g"	ppm	Blank, 0. – 999999.	F10	2	31
Dissolved Nitrogen—Spring	"Dissolved_N_S pring"	ppm	Blank, 0. – 999999.	F10	2	32
Total Phosphorus—Spring	"Total_P_Sprin g"	ppm	Blank, 0. – 999999.	F10	2	33
Dissolved Phosphorus—Spring	"Dissolved_P_S pring"	ppm	Blank, 0. – 999999.	F10	2	34
Sediment Concentration—Summer	"Sed_Conc_Su mmer"	ppm	Blank, 0. – 999999.	F10	2	35
Total Nitrogen—Summer	"Total_N_Sum mer"	ppm	Blank, 0. – 999999.	F10	2	36
Dissolved Nitrogen—Summer	"Dissolved_N_S ummer"	ppm	Blank, 0. – 999999.	F10	2	37
Total Phosphorus—Summer	"Total_P_Summ er"	ppm	Blank, 0. – 999999.	F10	2	38
Dissolved Phosphorus—Summer	"Dissolved_P_S ummer"	ppm	Blank, 0. – 999999.	F10	2	39
Sediment Concentration—Autumn	"Sed_Conc_Aut umn"	ppm	Blank, 0. – 999999.	F10	2	40
Total Nitrogen—Autumn	"Total_N_Autu mn"	ppm	Blank, 0. – 999999.	F10	2	41
Dissolved Nitrogen—Autumn	"Dissolved_N_A utumn"	ppm	Blank, 0. – 9999999.	F10	2	42
Total Phosphorus—Autumn	"Total_P_Autu mn"	ppm	Blank, 0. – 999999.	F10	2	43
Dissolved Phosphorus—Autumn	"Dissolved_P_A utumn"	ppm	Blank, 0. – 9999999.	F10	2	44
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	I1	2	45
<u>Go to Layout Matrix</u>		<u>Go t</u>	o Table of Conten	<u>ts</u>		

CELL DATA Required

	unea					
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed				Α	1	1-34
below.						
The following record repe	ats for the nur	nber of cells				
For cells with a Cell-Field ID of WATE	R, only fields	I-8 are used	in processing.			
Cell ID—Alphanumeric string identifying the cell.	"Cell_ID"			A10	2	1
Soil ID —Alphanumeric string identifying the soil type for the cell.	"Soil_ID"			A10	2	2
Must be the same as a soil ID in the SOIL DATA section. Leave						
blank if Management Field ID is "WATER".						

				r		
Description	Field Header for	Units (English)	Domain	For	Record	Field
Description	Record #1	{English; [SI]	{English} [SI]	mat	No.	No.
Management Field ID—Alphanumeric string identifying the field	"Mgmt Field I			A10	2	3
for the cell. Must be the same as a management field ID in the	D"			1110	2	5
MANAGEMENT FIELD DATA section. For a cell which is						
flooded with water throughout the year (such as the pool area behind						
a dam) enter "WATER",						
Reach ID—Alphanumeric string identifying the receiving reach.	"Reach_ID"			A10	2	4
Must be the same as a reach ID in the REACH DATA section .					L	
Reach Location code—Code identifying where runoff is added to	"Reach_Locatio		Blank,	I10	2	5
the specified reach. (Blank indicates 0)	II_Coue		0, or 1			
0—Runoff added at upstream end of reach.						
I—Runoff added at downstream end of reach.	"Coll Area"	(22720)	(0.000004 to	E10	2	6
Cell Area—Area within the cell.	"Cell_Area	{acres}	{0.00000410	FIU	2	0
		[licetares]	10000.03			
			4000.01			
Cell time of concentration—Time required to concentrate water at	"Time_of_Conc	hr	Blank or	F10	2	7
outlet from cell. If not a "WATER" cell and blank, AnnAGNPS will	" – –		0.01 to 100.0	1	-	
compute time of concentration from Sheet flow and concentrated						
flow variables for the cell.					l	
Cell average elevation—Representative elevation for the cell.	"Avg_Elevation	ft	{-3280.8 to	F10	2	8
	"	[m]	32808.3}			
			[-1000.0 to			
			10000.0j		<u> </u>	
RCN Calibration ID—Alphanumeric string identifying the RCN	"RCN_Calib_ID "			A10	2	9
calibration ID to be used for this cell. If non-blank, must be the same					İ	
as a RCN Calibration ID in the RCN CALIBRA HON DATA						
section. If KUN calibration has been requested, a blank defaults to the watershed scale PCN Calibration ID. If RCN calibration has not						
been requested, then this field is ignored						
Cell average land slone—Representative land slone for the cell.	"Avg Land Slo	len-vert/	0.00001 to 3.0	F10	2	10
Cell average land slope - Representative land slope for the cell	pe"	len-horz	0.00001 10 5.0	110	2	10
		(nd)				
Cell aspect—Representative land slope orientation for cell	"Aspect"	decimal °	Blank or	F10	2	11
measured from north in a clockwise direction.			0.0 to 360.0			
RUSLE/USLE 'ls' factor-RUSLE/USLE erosion equation length-	"RUSLE_ls_Fct		0.00001 to 100.0	F10	2	12
slope factor for normal erosion conditions.	r"					
RCN Retention Calibration Factor—used to calibrate the RCN	"RCN_Rtn_Cali	(nd)	Blank, or	F10	2	13
retention variable only for this cell. If RCN calibration has been	D_FCIF		0 to ∞			
requested, a blank defaults to the cell's RCN calibration ID's						
retention factor. If RCN calibration has not been requested, then a $11 - 1 - 1 - 1 - 1 - 1 - 1 - 0$						
blank defaults to 1.0.	"Secondary Cli			A 10	2	14
secondary Climate File ID—secondary Climate file ID for uns	mate_File_ID"			AIU	2	14
Shoot flow Manning's "n"_Roughness coefficient for Sheet flow	- "Sheet Flow M		Blank or	F10	2	15
within the cell Otherwise blank defaults to 0.150. Should be left	annings_n"		0.005 to 1.000	1.10	-	15
blank if cell time of concentration (t_c) has been entered. Regardless.			0.000 10 1.000			
the entered value for cell t_c is used in lieu of any value in this field.						
Geology ID—Alphanumeric string identifying the Geology data set	"Geology_ID"			A10	2	16
used for baseflow calculations for this cell. Must be the same as a						
Geology ID in the GEOLOGY DATA section. Leave Blank if						
watershed-scale defaults are to be used for baseflow calculations.						
Concentrated flow slope—Slope of concentrated flow path within	"Conc_Flow_Slo	len-vert /	Blank, or	F10	2	17
the cell. Should be left blank if cell time of concentration (t _c) has	pe	len-horz	0.00001 to 3.0			
been entered. Regardless, the entered value for cell t_c is used in lieu		(nd)				
of any value in this field.						
Diank defaults to the Cell's average land slope.					1	

	Diald	TI	-			
Description	Header for Record #1	{English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Concentrated flow length —Length of concentrated flow path within the cell after the first 100 meters (328 feet). Blank indicates value is computed from receiving reach length coefficient and exponent. Should be left blank if cell time of concentration (t_c) has been entered. Regardless, the entered value for cell t_c is used in lieu of any value in this field.	"Conc_Flow_Le ngth"	{ft} [m]	Blank, or {0.0 to 328080.0} [0.0 to 99999.0]	F10	2	18
Hydraulic Geometry ID —Alphanumeric string identifying the Hydraulic Geometry ID to be used for this cell. Must be the same as an available bulit-in Hydraulic Geometry ID or a valid Hydraulic Geometry ID created by the user in the HYDRAULIC GEOMETRY DATA section. Blank defaults to Hydraulic Geometry ID for the cell's receiving reach.	"Hydraulic_Geo m_ID"		Blank or valid Hydraulic Geometry ID	A10	2	19
Concentrated flow hydraulic depth —Representative rectangular channel hydraulic depth for concentrated flow path within the cell. Depth is for a 2 year 24-hr storm event runoff as determined by the flow area divided by the top width. Blank indicates value is computed from receiving reach width coefficient and exponent. Should be left blank if cell time of concentration (t_c) has been entered. Regardless, the entered value for cell t_c is used in lieu of any value in this field.	"Conc_Flow_Hy draulic_Depth"	{ft} [m]	Blank, or {0.0 to 131.0} [0.0 to 40.0]	F10	2	20
Concentrated flow Manning's "n" —Roughness coefficient for concentrated flow within the cell. Blank defaults to 0.040. Should be left blank if cell time of concentration (t _c) has been entered. Regardless, the entered value for cell t _c is used in lieu of any value in this field.	"Conc_Flow_M annings_n"		Blank, or 0.005 to 1.000	F10	2	21
Sheet flow slope —Slope of Sheet (sheet) flow path within the cell. Should be left blank if cell time of concentration (t _c) has been entered. Regardless, the entered value for cell t _c is used in lieu of any value in this field. Blank defaults to the Cell's average land slope.	"Sheet_Flow_Sl ope"	len-vert / len-horz (nd)	Blank, or 0.00001 to 3.0	F10	2	22
Sheet flow length —Length of Sheet (sheet) flow path within the cell. Default value is 50 meters (164 feet). Should be left blank if cell time of concentration (t_c) has been entered. Regardless, the entered value for cell t_c is used in lieu of any value in this field.	"Sheet_Flow_Le ngth"	{ft} [m]	Blank, or {0.0 to 164.} [0.0 to 50.0]	F10	2	23
Shallow Concentrated flow slope —Slope of shallow concentrated flow path within the cell. Should be left blank if cell time of concentration (t_c) has been entered. Regardless, the entered value for cell t_c is used in lieu of any value in this field. Blank defaults to .00001.	"Shallow_Conc_ Flow_Slope"	len-vert / len-horz (nd)	Blank, or 0.00001 to 3.0	F10	2	24
Shallow Concentrated flow length —Length of shallow concentrated flow path within the cell. Default value is 50 meters (164 feet). Should be left blank if cell time of concentration (t _c) has been entered. Regardless, the entered value for cell t _c is used in lieu of any value in this field.	"Shallow_Conc_ Flow_Length"	{ft} [m]	Blank, or {0.0 to 164.} [0.0 to 50.0]	F10	2	25
Delivery Ratio —Delivery ratio from all landscape sources of erosion (sheet & rill and gullies) to sediment yield at the cell's receiving stream reach. Recommended procedure is the HUSLE equation which is the default procedure when this field is left blank.	"Delivery_Ratio "	(nd)	Blank, or 0. to 1.	F10	2	26
Constant USLE C-factor —constant USLE C-factor which will supercede any internal calculated RUSLE C-factor values. Blank defaults to internal RUSLE calculated C-factor values.	"Constant_USL E_C_Fctr"		Blank, 0. to 1.	F10	2	27
Constant USLE P-factor —constant USLE P-factor which will supercede any internal calculated RUSLE P-factor values. Blank defaults to internal RUSLE calculated P-factor values.	"Constant_USL E_P_Fctr"		Blank, 0. to 1.	F10	2	28
All Organic Carbon Calibration Factor—used to calibrate all organic carbon for this cell only. Defaults to the watershed-scale organic carbon from all sources calibration factor in the PL CALIBRATION DATA section.	"All_OC_Calib_ Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	29
All Nitrogen Calibration Factor—used to calibrate all nitrogen for this cell only. Defaults to the watershed-scale nitrogen from all sources calibration factor in the PL CALIBRATION DATA section.	"All_N_Calib_F ctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	30

Description	Field Header for Record #1	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.	
All Phosphorus Calibration Factor—used to calibrate all phosphorus for this cell only. Defaults to the watershed-scale phosphorus from all sources calibration factor in the PL CALIBRATION DATA section.	"All_P_Calib_F ctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	31	
Sheet & Rill Erosion Calibration Factor —used to calibrate sheet & rill erosion for this cell only. Defaults to the watershed-scale sediment from sheet & rill sources calibration factor in the PL CALIBRATION DATA section.	"Sheet_and_Rill _Erosion_Calib_ Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	32	
Gullies Erosion Calibration Factor—used to calibrate gully erosion for this cell only. Defaults to the watershed-scale sediment from gully sources calibration factor in the PL CALIBRATION DATA section.	"Gullies_Erosio n_Calib_Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	33	
 RUSLE2 ID—Alphanumeric string identifying the RUSLE2 ID to be used for this cell. If non-blank, must be the same as a RUSLE2 ID in the RUSLE2 DATA section. If this parameter is blank and 1.) The RUSLE2 flag is set to true and a default RUSLE2 ID specified in the "Global IDs, Factors, and Flags Data" section then the default RUSLE2 ID will be used. 2.) The RUSLE2 flag is set to false then RUSLE2 parameters will not be used for this cell. 	"RUSLE2_ID"			A40	2	34	
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	Il	2	35	
Go to Layout Matrix	Go to Table of Contents						

CLASSIC GULLY DATA¹

Optional

	Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
--	-------------	----------------------------------	----------------------------	--------------------------	------------	---------------	--------------

Note: A gully is either cell-located or reach-located. The gully is cell-located if the gully is wholly contained within a single AnnAGNPS cell; the gully is reach-located if the gully is located within the thalweg of a reach. A cell-located gully is affected by only one AnnAGNPS cell; a reach-located gully is affected by more than just one AnnAGNPS cell—all cells upstream of the upstream end of the reach plus any contribution from the reach's local cells. A gully's delivery ratio is defined to be the gully yield to its receiving stream divided by the gully's sediment yield to the gully mouth. The delivery ratio for the gully erosion to its mouth is internally determined.

If the gully is cell-located, the gully's cell is a single AnnAGNPS cell which is the "Gully's Cell ID". The gully's cell's drainage area that affects the gully erosion may be only a portion or can be all of this cell's total drainage area. The drainage area affecting the cell-located gully erosion is the "Gully's Cell's Drainage Area". The delivery ratio to the gully's cell's receiving stream is also a function of the drainage area of this cell's subarea to its receiving stream. The drainage area of the gully's cell's subarea is described as the "Cell's Drainage Subarea". A gully's cell may be composed entirely of just one subarea, where the gully's subarea would be equal to the gully's cell's total drainage area, or many subareas, where the gully may be located in only one of the gully's cell's subareas.

If the gully is reach-located, the drainage area affecting the gully is all or part of each of the reach's local-contributing AnnAGNPS cells' drainage area (Gully's Reach's Local-Cells Drainage Area)—both the reach's left-bank cell and the reach's right-bank cell—and all of the drainage area above the upstream end of the reach. For a reach-located gully, the "Gully's Reach's Local-Cells Drainage Area Fraction" is ratio of the sum of all local-cells' drainage area contributing to the gully mouth to the sum of all local-cells' total drainage area. Since the reach-located gully's mouth is located within the reach, the gully's sediment yield from the mouth to the receiving reach is equal to the gully's sediment yield to the mouth and its subsequent "Delivery Ratio" must be unity. Therefore, no subarea information is required.

Field Header— Required unique field header for each field listed below.				А	1	1-21	
The following record repeats for the number of classic gullies. Multiple gullies within a cell must be consecutive.							
Classic Gully ID—Alphanumeric string identifying the classic	"Gully_ID"			A10	2	1	
gully.							

¹ A classic gully is cell-located if only one cell contributes to the flow at the mouth of the gully. A classic gully is reach-located if more than one cell contributes to the flow at the mouth of the gully.

 AnnAGNPS Input File Specifications

 2019
 File Name: Input_Specifications_v5.51_2019.xx.xx.docx
 Revision: December 4, 2019

Description	Field Header for	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.
	Record #1	[SI]	(
Cell ID—Only if gully is cell-located, alphanumeric string	"Cell_ID"			A10	2	2
identifying the cell that contains the classic gully if the gully's						
drainage area is wholly contained within a single cell (cell-located).						
Must be the same as a cell ID in the CELL DATA section.						
Reach ID—Only if gully is reach-located, alphanumeric string	"Reach_ID"			A10	2	3
identifying the reach ID whose thalweg contains the gully mouth.						
Must be the same as a reach ID in the REACH DATA section.						
Soil ID -Alphanumeric string identifying the dominant soil type for	"Soil_ID"			A10	2	4
the gully or other erosion point source. Must be the same as a soil						
ID (in the SOIL DATA section). Blank defaults to the Soil ID for						
the cell that contains the mouth of the gully, or the left-bank cell if						
the guily is in the cell's receiving reach.	"Coll Duoinago		D1 1	E10	2	
Cell's Drainage Area —Only if the gully is cell-located, the	Area"	{acres}	Blank, or $(0.000025.4)$	F10	2	2
drainage area is that portion of the cell's drainage area contributing		[hectares]	$\{0.000025\ to$			
to the mouth of the guily. A blank defaults to the entire cell			9884.}			
drainage area. The default is the entire cell's drainage area.			4000 01			
Bassh's Logal Calls Drainage Area Only if the gully is reach	"Reach Drainag	(norros)	Plank or	E10	2	6
located this drainage area is sum of that portion of the drainage	e_Area"	{acres}	$\int 0.00025 \text{ to}$	110	Z	0
areas from all of the contributing reach's local-cells' drainage areas		[neetares]	9884 3			
draining to the mouth of the gully. The default is sum of the all of			[0 00001 to			
the local-cell's drainage areas.			4000.01			
Head Cut Denth -Gully (erosion point-source) head cut denth used	"Headcut Dept	{in}}	Blank or	F10	2	7
to composite the eroded soil lavers. Blank defaults to 1 st soil laver.	h"	[mm]	$\{0.00 \text{ to } 120.\}$	110	-	,
1 5 5		LJ	[0.00 to 3000]			
Erosion Coefficient—Coefficient in classic gully erosion power	"Erosion_Coef"		[>0.0 to 50. for	F10	2	8
curve (note that the units for both the erosion (Q_s) and the			Q_s in mm] ²			
rainfall/runoff (Q _w) may be unit area or totals but must be						
consistent within a power curve):						
O _s =coef*O _w exp						
where $Q_w = rainfall/runoff$ volume [unit area-(in or mm ³); total						
units-(AF or Mg)]						
Q_s = sediment discharge [unit area-(T/ac or Mg/ha); total						
units-(T or Mg)].						
A blank is not allowed						
Erosion Exponent—Exponent in classic gully erosion power curve	"Erosion_exp"		Blank or	F10	2	9
(note that the units for both the erosion (Q_s) and the rainfall/runoff			[0.0 to 3.0 for $Q_{\rm w}$			
(Q _w)) may be unit area or totals but must be consistent within each			in mm]			
power curve):						
$Q_s = coef^*Q_w^{exp}$						
where $Q_w = rainfall/runoff$ volume [unit area-(in or mm ³); total						
units-(AF or Mg)]						
Q_s = sediment discharge [unit area–(T/ac or Mg/ha); total						
units-(T or Mg)].						
A blank defaults to zero.						
Delivery Ratio —Delivery ratio of gully erosion to gully yield.	"Delivery_Ratio	(nd)	Blank, 0 to 1	F10	2	10
Blank defaults to HUSLE delivery ratio algorithm for cell-located						
gully; one if reach-located.						
Management Field ID—Alphanumeric string identifying the field	"Mgmt_Field_I D"			A10	2	11
for the classic gully. Must be the same as a management field ID in	,					
the Management Field Data section. Blank defaults to no field						
management effects	1	1	1	1		

² Unit conversion from English to SI is non-linear. Appropriate English ranges would restrict the erosion to less than 136 T/ac.

	Theld	T. A.				
Description	Header for	{English}	Domain	For	Record	Field
	Record #1	[SI]	{English} [SI]	mat	No.	No.
Cell's Drainage Subarea —If the gully is cell-located (i.e.: the	"Cell Drainage	{acres}	Blank, or	F10	2	12
gully's drainage area is located wholly within one AnnAGNPS cell).	Subarea"	[hectares]	{0.000025 to	110	-	
the cell's total drainage area may actually be comprised of many		[]	9884.}			
subareas that drain separately into and along the receiving reach			[0.00001 to			
such as likely to be the case for either the left- or right-bank cells,			4000.0]			
even though the cell's entire drainage is designated to only enter			-			
either the upstream or downstream end of the reach. Then the gully						
may be located in one of these cell's subareas, requiring a						
designation of the drainage area of the subarea containing the gully						
to the receiving reach as the Cell's Drainage Subarea. A blank						
means that cell's drainage subarea is composed of the entirely cell's						
drainage area (such as a source cell with the gully mouth located at						
the downstream end of the cell), resulting in the subarea containing						
the gully to be equal to the cell's total drainage area. A blank is						
required if the gully is reach-located.						
Load Calibration Factor—calibration factor used to calibrate the	"Load_Calib_Fc		Blank, 0 to ∞	F10	2	13
classic gully sediment yield to its loading at a known point (usually	tr					
at a USGS gaging station). Blank defaults to 1.						
Currently not used – Reserved.						
Rainfall/Runoff Indicator—Code to indicate whether the power	"Rainfall/Runof f. Indicator"		Blank, 0 or 1	I1	2	14
curve's volume (Q) is a function of rainfall or runoff. Blank						
defaults to 1; 0-rainfall, 1-runoff						
Units Indicator—Code to indicate whether the regression	"Units_Indicato		Blank, 0 or 1	11	2	15
coefficient and exponents are for unit area or total units. Blank						
defaults to 1; 0-unit area, 1-total mass.	"Cully Logation		D1 1 4779	т 1	2	16
Gully Location Code—If gully is cell-located, code ,must be a "1"	Code"		Blank, "I" or	LI	2	16
or blank. If guily is reach-located, code must be an F. Blank			Г			
defaults to true 1. (Note to Editor's editor: this field is not to be						
determined within the Editor depending upon whether the user						
designates the gully as cell- or reach-located)						
Owner Carbon Calibration Factor wood to calibrate alagain	"OC Calib Fatr	(nd)	Dlamle 0.0 an	E10	2	17
organic Carbon Canoration Factor—used to canorate classic	"	(na)	00000001 to	F10	Z	1 /
the watershed scale organic carbon from classic gully sources						
calibration factor in the PL CALIBRATION DATA section A			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			
"(0.0" (at least one zero & a decimal) means that none of this						
nollutant originates from this source. A complete field for the real-						
number input consisting of nine 9's & a decimal ("9999999999").						
which is AnnAGNPS' real-number input for infinity, means that all						
physically-possible pollutant originates instantaneously from this						
source for each runoff event. Any real-number smaller than this						
"infinity" is taken to be exactly the value of this real-number.						
Nitrogen Calibration Factor—used to calibrate classic gully	"N_Calib_Fctr"	(nd)	Blank, 0.0, or	F10	2	18
nitrogen for this classic gully only. Blank defaults to the watershed-			.000000001 to			
scale nitrogen from classic gully sources calibration factor in the PL			9999999999.			
CALIBRATION DATA section. A "0.0" (at least one zero & a						
decimal) means that none of this pollutant originates from this						
source. A complete field for the real-number input consisting of						
nine 9's & a decimal ("999999999."), which is AnnAGNPS' real-						
number input for infinity, means that all physically-possible						
pollutant originates instantaneously from this source for each runoff						
event. Any real-number smaller than this "infinity" is taken to be						
exactly the value of this real-number.				1		

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Phosphorus Calibration Factor —used to calibrate classic gully phosphorus for this classic gully only. Blank defaults to the watershed-scale phosphorus from classic gully sources calibration factor. in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Calib_Fctr"	(nd)	Blank, 0.0, or .000000001 to 999999999.	F10	2	19
Erosion Calibration Factor —used to calibrate classic gully erosion for this classic gully only. Blank defaults to the watershed- scale sediment from classic gully sources calibration factor in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Erosion_Calib_ Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	20
Input Units Code — Code identifying whether input is in English or SI units. $0 =$ English $, 1 =$ SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	I1	2	21
Go to Layout Matrix		<u>Go to</u>	Table of Conten	ts		

CONTOUR DATA

Optional unless referenced in Management Schedule Data	a
--	---

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-6
The following record repeats	for the number	• of contour	sets.		-	-
Contour ID —Alphanumeric string identifying the Contour Data.	"Contour_ID"			A10	2	1
Ridge Height code—Code indicating the height of the contour ridges. Allowable codes are: 0 = No ridge (non-cropland only) 1 = very low (0.5"-2") ridges 2 = low (2"-3") ridges 3 = moderate (3"-4") ridges 4 = high (4"-6") ridges 5 = very high (>6") ridges 6 = variable ridge heights. Zero value is entered for non-crop landuse where only mechanical disturbance calculation is desired. If blank, 3-moderate ridges will be used.	"Ridge_Height_ Code"		Blank, or 0 to 6	I1	2	2
Furrow Slope —Slope of the furrow. Can be left blank if the landuse is non-crop and only mechanical disturbance calculation is desired.	"Furrow_Slope"	len-vert / len-horz (nd)	Blank, or 0.00001 to 1.0	F10	2	3

 AnnAGNPS Input File Specifications

 2019
 File Name: Input_Specifications_v5.51_2019.xx.xx.docx
 Revision: December 4, 2019

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.	
Disturbed cover code - Code indicating condition of cover related to soil disturbance. Acceptable codes are: <u>Cropland with EI distribution number < 400</u> 1 = C1) established = sod-forming grass 2 = C2) 1st year grass or cut for hay 3 = C3) heavy cover = And/or very rough 4 = C4) moderate cover = and/or rough 5 = C5) light cover = and/or mod = rough 6 = C6) no cover and/or min = rough. 7 = C7) clean tilled, smooth, fallow <u>Cropland with EI distribution number ≥ 400</u> 8 = VR) very rough with stubble <u>Non-cropland</u> 1 = R1) very rough; plant + rock cover > 50% 2 = R2) very rough; plant + rock cover < 50% 3 = R3) rough; plant + rock cover < 50% 4 = R5) moderately rough; plant + rock cover < 25% A blank defaults to 5.	"Disturbed_Cov er"		Blank, or 1 to 8	II	2	4	
Consolidated cover code- Code indicating condition of cover related to soil consolidation. Required only for non-crop landuse. Acceptable codes are: 1 = R3) rough; plant + rock cover > 50% 2 = R4) rough; plant + rock cover < 50% 3 = R6) moderately rough; est = veg.; cover < 40% 4 = R8) slightly rough; est = veg.; cover < 35% 5 = R9) smooth; est. veg.; plant + rock cover < 25% A blank defaults to 3.	"Consolated_Co ver"		Blank, or 1 to 5	I1	2	5	
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	I1	2	6	
Go to Layout Matrix	Go to Table of Contents						

CROP DATA

Optional unless referenced in Management Schedule Data

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-41
The following record rep	eats for numb	er of crops.				
Crop ID—Alphanumeric string identifying the crop.	"Crop_ID"			A20	2	1
Yield Units Harvested per Area—Number of units of yield per unit of area at harvest. (e.g. For a yield of 50 bushels per acre: enter 50.)	"Yield_Units_H arvested"	{yield units /ac} [yield units/ha]	[0.0 to 100000.0]	F10	2	2
Residue Mass Ratio—Mass ratio of residue to yield.	"Residue_Mass_ Ratio"	[nd]	0.0 to 1000.0	F10	2	3
Surface decomposition—Surface residue decomposition coefficient. (Blank defaults to 0.016)	"Surface_Deco mp"		Blank or 0.0 to 1.0	F10	2	4
Sub-surface decomposition —Sub-surface residue decomposition coefficient. (Blank defaults to 0.016)	"Sub- surface_Decomp "		Blank or 0.0 to 1.0	F10	2	5

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
USLE C-Factor —optional, constant USLE C-factor to be used in lieu of computed RUSLE time-varying value. This value will be used unless overridden by the Constant USLE C-factor located in the CELL DATA section. Blank field defaults to either the computed RUSLE time-varying value or the Constant USLE C-factor in the CELL DATA section.	"USLE_C_Fctr"		Blank or 0.0 to 1.0	F10	2	6
Moisture Depletion Rate of moisture depletion, used only for the Pacific Northwest. Only needed for following EI distribution numbers: 6-10, 14-15, 31-40, 58-60, 63 (EI distribution number entered with Simulation Period Data) otherwise leave blank. Blank defaults to 0.0. Example values from RUSLE: w. wheat & other deep rooted crops 1.0 spring wheat & barley 0.75 spring peas & lentils 0.67 shallow rooted crops 0.50 summer fallow 0.0	"Moisture_Depl etion"	{??} [??]	Blank, or 0.0 to 1.0	F10	2	7
Residue Adjust Amount —Minimum residue amount to adjust runoff curve number. (Blank defaults to 0) Currently not used – Reserved.	"Residue_Adjus t_Amt"	{lb / acre} [kg / hectare]	Blank, or {0.0 to 99924.} [0.0 to 112000.0]	F10	2	8
Crop Residue —Surface residue from crop. (Three values in order for 30, 60, and 90 percent cover.). At least one of the three values must be >0. (Blank defaults to 0.0)	"Crop_Residue_ 30%", Crop_Residue_6 0%, Crop_Residue_9 0%	{lb / acre} [kg / hectare]	Blank, or {0.0 to 99924.} [0.0 to 112000.0]	3F10	2	9-11
Annual crop code.—Acceptable values are: 0 = cool season 1 = Annual 2 = perennial Blank defaults to 0. Currently not used – Reserved.	"Annual_Crop_ Code"		Blank , 0, 1 or 2	I1	2	12
Legume code. - Acceptable values are : Y = Yes and N = No (Blank indicates No) Currently not used.	"Legume_Code"	[nd]	Blank, Y or N	A1	2	13
Senescence code—Code indicating whether crop senescence increases with crop residue. Acceptable values are: Y = Yes N = No (Blank indicates ves.)	"Senescence_Co de"	[nd]	Blank, Y or N	A1	2	14
Yield Unit Name (optional)—Yield unit name for crop. (Ontional—For user reference only. Not read by AnnAGNPS.)	"Yield_Unit_Na me"			A10	2	15
Yield Unit Mass—Mass of the unit of yield for the crop. (See Yield Unit Name for unit description.).	"Yield_Unit_Ma ss"	{lb per unit of yield} [kg per unit of yield]	{0.0 to 88105.} [0.0 to 40000.0]	F10	2	16
Harvest C-N Ratio—Ratio of Carbon to Nitrogen for crop at harvest.	"Harvest_C- N_Ratio"		1.0 to 200.0	F10	2	17
Pre-harvest C-N Ratio —Ratio of Carbon to Nitrogen for crop before harvest. Currently not used – Reserved.	"Pre- Harvest_C- N_Ratio"		Blank or 1.0 to 200.0	F10	2	18
Harvest Water—Water content of harvested portion of crop. Currently not used – Reserved.	"Harvest_Water	wt / wt (nd)	Blank or 0.0 to 1.0	F10	2	19
N Uptake—Nitrogen uptake per yield unit	"N_Uptake"	wt-N / wt- harvest unit (nd)	0.0 to 1.0	F10	2	20
P Uptake—Phosphorus uptake per yield unit	"P_Uptake"	wt-P / wt- harvest unit (nd)	0.0 to 1.0	F10	2	21
Harvest C-P Ratio—Ratio of Carbon to Phosphorus for crop at harvest.	"Harvest_C- P_Ratio"		50.0 to 400.0	F10	2	22
Pre-harvest C-P Ratio —Ratio of Carbon to Phosphorus for crop before harvest. Currently not used – Reserved.	"Pre- Harvest_C- P Ratio"		Blank or 50.0 to 400.0	F10	2	23

Revision: December 4, 2019 File Name: Ir	nput_Specifications_v5.51_2019.xx.xx.docx
--	---

Description	Field Header for	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.
	Record #1	[SI]	(1.00	1100
Growth Time —Accumulated fraction of time from planting to harvest for ending each of 4 growth stages: initial; development; mature; and senescence. Fourth entry must be 1.0.	"Growth_Time_ Ini", "Growth_Time_ Dev", "Growth_Time_ Mat", "Growth_Time_ Son"		0.0 to 1.0	4F10	2	24-27
Growth N Uptake —Fraction of Nitrogen uptake from planting to harvest for each of 4 growth stages: initial; development; mature; and senescence. Four fractions entered must sum to 1.0.	"Growth_N_Upt ake_Ini", "Growth_N_Upt ake_Dev", "Growth_N_Upt ake_Mat", "Growth_N_Upt ake Sen"		0.0 to 1.0	4F10	2	28-31
Growth P Uptake —Fraction of Phosphorus uptake from planting to harvest for each of 4 growth stages: initial; development; mature; and senescence. Four fractions entered must sum to 1.0.	"Growth_P_Upt ake_Ini", "Growth_P_Upt ake_Dev", "Growth_P_Upt ake_Mat", "Growth_P_Upt ake Sen"		0.0 to 1.0	4F10	2	32-35
Basal Crop Coefficient ("Kcb-ini") —Coefficient for the initial growth stage; used in adjusting the potential evapotranspiration (ET). The coefficient for all days in growth stage 1 will be equal to this value. The coefficient is internally set to 0.0 for all days outside of the growing season. Blank defaults to 0.15.	"Basal_Crop_C oef_Ini"	[nd]	0.0 to 1.2	F10	2	36
Basal Crop Coefficient ("Kcb-dev") —Coefficient for the development growth stage; used in adjusting the potential evapotranspiration (ET). (Currently reserved for future use.) Currently, the days in growth stage 2 will be linearly interpreted between the last day of stage 1 to the first day of stage 3. The coefficient is internally set to 0.0 for all days outside of the growing season. Blank defaults to 1.0	"Basal_Crop_C oef_Dev"	[nd]	0.0 to 1.2	F10	2	37
Basal Crop Coefficient ("Kcb-mid") —Coefficient for the mature growth stage; used in adjusting the potential evapotranspiration (ET). The coefficient for all days in growth stage 3 will be equal to this value unless internally adjusted based on climate conditions. The coefficient is internally set to 0.0 for all days outside of the growing season. Blank defaults to 1.15.	"Basal_Crop_C oef_Mid"	[nd]	0.0 to 1.2	F10	2	38
Basal Crop Coefficient ("Kcb-end") —Coefficient for the last day of the senescence growth stage; used in adjusting the potential evapotranspiration (ET). The days in growth stage 4 will be linearly interpreted between the last day of stage 3 to the last day of stage 4. Futher adjustment may be made based on climate conditions. The coefficient is internally set to 0.0 for all days outside of the growing season. Blank defaults to 0.15.	"Basal_Crop_C oef_End"	[nd]	0.0 to 1.2	F10	2	39
Basal Crop Coefficient Climatic Adjustment code —Code indicating whether the basal crop coefficients "Kcb-mid" and "Kcb- end" will be adjusted based on climate conditions. Acceptable values are: $Y = Adjustment$; $N = No$ adjustment. Blank will default to 'Y' unless the global basal crop climate adjustment code is set to false in the Global IDs, Factors, and Flags Data section.	"Basal_Crop_C oef_Climate_Ad just"	[nd]	Blank, Y or N	A1	2	40
Input Units Code — Code identifying whether input is in English or SI units. 0 = English ,1 = SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	8	41
Go to Layout Matrix		Go to	Table of Conten	ts		

CROP GROWTH DATA

Optional unless CROP DATA is referenced in Management Schedule Data

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-5
The following record repeats 1 to 24 times for each crop in the CROP DATA section. Each crop-growth record represents a 15+ day period in a year for the crop. At least one crop-growth record is required for each crop. Time is measured from the planting of crop.						
Crop Growth ID —Alphanumeric string identifying the crop for which this crop-growth record applies. Must match a Crop ID in the CROP DATA section. Crop-growth records must be contiguous and sequential.	"Crop_Growth_ ID"			A20	2	1
Root Mass —Array representing the live root mass in the top 100 mm (4 inch) of soil. The values start at 0 days of plant growth and increment each 15 days.	"Root_Mass"	{lb / acre} [kg / hectare]	{0.0 to 99924.} [0.0 to 112000.0]	F10	2	2
Canopy Cover —Array representing the ratio of ground covered by the crop canopy to total ground area. The values start at 0 days of plant growth and increment each 15 days.	"Canopy_Cover "		0.0 to 1.0	F10	2	3
Rain Fall Height —Array representing the average intercepted rain drop fall height from the plant canopy to the ground. The values start at 0 days of plant growth and increment each 15 days.	"Rain_Fall_Hei ght"	{ft} [m]	{0.0 to 262.} [0.0 to 80.0]	F10	2	4
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}, 1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	5
Go to Layout Matrix	Go to Table of Contents					

EI PERCENTAGE DATA

Optional unless the EI number specified in the SIMULATION PERIOD DATA section above is > 149 or replacement percentages for a built-in EI number are desired

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-24
The following record is entered only if the "EI number" specified in the SIMULATION PERIOD DATA section above is > 149 or if replacement percentages for a built-in EI number are desired (Values for EI numbers up to 149 are built-in to the AnnAGNPS model). This is a non-repeating record.						
EI Percentages (Periods 1 to 24) —Cummulative EI percentages for the 24 15+ day periods in a year for the "EI number" entered in the SIMULATION PERIOD DATA section.	"EI_Pct_xx" where "xx" = "01" to "24"		0.0 to 100.0	24F10	2	1-24
Go to Layout Matrix	Go to Table of Contents					

EPHEMERAL GULLY DATA³ Optional

³ An ephemeral gully is cell-located if only one cell contributes to the flow at the mouth of the gully. A gully is reach-located if more than one cell contributes to the flow at the mouth of the gully.

Description	Field Header for	Units {English}	Domain (English) [SI]	For mat	Record	Field No	
	Record #1	[SI]	{English} [51]	mai	110.	140.	
Record #1 SI Carget , P P And Constant , P Constant , Constant , Co							
the reach-located gully's mouth is located within the reach, the gully	s sediment yiel	d from the n	nouth to the receivi	ng rea	ch is equa	l to the	
gully's sediment yield to the mouth and its subsequent "Delivery Rat	io" must be uni	ty. Therefor	e, no subarea infor	mation	is require	ed.	
Field Header— Required unique field header for each field listed below.				A	1	1-30	
The following record repeats for the number of ephemeral	gullies. Multi	iple gullies v	within a cell must	be con	secutive.		
Ephemeral Gully ID—Alphanumeric string identifying the ephemeral gully.	"Gully_ID"			A10	2	1	
Cell ID —Only if gully is cell-located, alphanumeric string	"Cell_ID"			A10	2	2	
drainage area is wholly contained within a single cell (cell-located). Must be the same as a cell ID in the CELL DATA section.							
Reach ID —Only if gully is reach-located, alphanumeric string identifying the reach ID whose thalweg contains the gully mouth. Must be the same as a reach ID in the REACH DATA section.	"Reach_ID"			A10	2	3	
Soil ID —Alphanumeric string identifying the dominant soil type for the ephemeral gully. Must be the same as a soil ID in the SOIL DATA section. Blank defaults to the Soil ID for the cell (1 st) that contains the mouth of the gully, or the 1 st cell identified as contributing to the reach if the gully is reach-located.	"Soil_ID"			A10	2	4	
Drainage Area to Mouth —The total drainage area is the entire drainage area contributing to the mouth of the gully. A blank default for a cell-located gully is the entire cell's drainage area.	"Drainage_Area _to_Mouth"	{acres} [hectares]	Blank, or {0.000025 to 9884.} [0.00001 to 4000.0]	F10	2	5	
Local Drainage Area —Only if the gully is reach-located, this drainage area is sum of that portion of the drainage areas from all of the contributing reach's local-cells' drainage areas draining to the mouth of the gully. The default is sum of the all of the local-cell's drainage areas.	"Local_Drainag e_Area"	{acres} [hectares]	Blank, or {0.000025 to 9884.} [0.00001 to 4000.0]	F10	2	6	
Gully Slope —Land slope immediately upstream from the mouth of the gully. Blank defaults to 0.00001.	"Gully_Slope"	l-vert / l- horz (nd)	Blank, or 0.00001 to 3.0	F10	2	7	
Critical Shear Stress —Critical shear stress at which gully erosion begins. Blank defaults to internal calculation based upon the gully's clay, silt, & sand content as determined using the gully's soil ID.	"Critical_Sheer _Stress"	{lbs/ft ² } [N/m ²]	Blank, or {>0 to 2.05} [>0 to 100.0]	F10	2	8	
Gully Location Code —If gully is cell-located, code ,must be a "T" or blank. If gully is reach-located, code must be an "F". Blank defaults to true "T" (Note to Editor's editor: this field is not to be under the user's direct control, but is to be determined within the Editor depending upon whether the user completes the fields as cell-or reach-located.)	"Gully_Location _Code"		Blank, "T" or "F"	L1	2	9	

	Field	Units	Domain	For	Record	Field
Description	Header for Record #1	{English} [SI]	{English} [SI]	mat	No.	No.
Management Field ID —Alphanumeric string identifying the field for the ephemeral gully. Must be the same as a management field in the MANAGEMENT FIELD DATA section.	"Mgmt_Field_I D"		Blank or valid Management Field ID	A10	2	10
A management field ID of 'BUFFER' is internally defined as the gully headcut location in a non-crop riparian buffer, resulting in no gully erosion.						
A management field ID of 'WETLAND' is internally defined as the gully headcut location in a wetland, resulting in no gully erosion.						
Blank defaults to Management Field ID for the cell that contains the mouth of the gully, or the primary cell identified as contributing to the reach if the gully is reach-located.						
Erosion Depth —ephemeral gully depth of erosion. If blank, the erosion depth is determined by the maximum 'Operation Tillage Depth' found for all associated operations where the surface 'Area Disturbed' is 50% or greater. See the 'Management Operation Data' section for more details on the 'Operation Tillage Depth' and 'Area Disturbed' parameters.	"Erosion_Depth "	{in} [mm]	Blank, or 0 to bottom of soil column.	F10	2	11
Cell's Drainage Subcell —If the gully is cell-located (i.e.; the gully's drainage area is located wholly within one AnnAGNPS cell), the cell's total drainage area for local subareas that are created as left- & right-bank cells may actually be comprised of many subcells that drain separately into and along the receiving reach or riparian buffer such as likely to be the case for either the left- or right-bank cells, even though the cell's entire drainage is designated to only enter either the upstream or downstream end of the reach. Then the gully may be located in one of these cell's subcells, requiring a designation of the drainage area of the subcell containing the gully to the receiving reach or to the location where the flow path enters the riparian buffer, if present. A blank means that cell's drainage subcell is composed of the entirely cell's drainage area (such as a source cell with the gully mouth located at the downstream end of the cell's total drainage area. A blank is required if the gully is reach-located.	"Cells_Drainage _Subcell"	{acres} [hectares]	Blank, or {0.000025 to 9884.} [0.00001 to 4000.0]	F10	2	12
Hydraulic Geometry ID —Alphanumeric string identifying the hydraulic geometry to be used for the ephemeral gully. Must be the same as an available bulit-in Hydraulic Geometry ID or a valid Hydraulic Geometry ID created by the user in the HYDRAULIC GEOMETRY DATA section. Blank defaults to Hydraulic Geometry ID for the cell that contains the mouth of the gully, or for the reach that contains the mouth of the gully if it is reach-located. The width algorithm for the gully-location's hydraulic geometry equation (c) must also be true to be effective.	"Hydraulic_Geo metry_ID"		Blank or valid ID from the Hydraulic Geometry Data section.	A10	2	13
Width Algorithm—Flag to indicate if Nachtergaele et al's (2002) equation may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for Nachtergaele et al's (2002) equation (b).	"Width_Nachter gaele"		Blank, or T/F	L1	2	14
Width Algorithm—Flag to indicate if the gully-location's hydraulic geometry equation may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for the gully-location's hydraulic geometry equation (c).	"Width_Hydrau lic_Geometry"		Blank, or T/F	L1	2	15
Width Algorithm —Flag to indicate if non-submerging tailwater at headcut crest equation may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for the non-submerging tailwater at headcut crest equation (d).	"Width_Non- submerging_Tai lwater"		Blank, or T/F	L1	2	16

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Width Algorithm —Flag to indicate if Woodward's (1999) equilibrium gully width equation may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag Woodward's (1999) equilibrium gully width equation (e).	"Width_Woodw ards_Equilibriu m"		Blank, or T/F	L1	2	17
Width Algorithm —Flag to indicate if Woodward's (1999) ultimate gully width equation may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for Woodward's (1999) ultimate gully width equation (f).	"Width_Woodw ards_Ultimate"		Blank, or T/F	L1	2	18
Width Algorithm —Flag to indicate if Modified Wells' (2013) gully width equation #9 may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for Modified Wells' (2013) gully width equation (g).	"Width_Wells_ Eq.9"		Blank, or T/F	L1	2	19
Note: See additional available widt	h algorithms b	eginning in	field #31			
Delivery Ratio —Delivery ratio of ephemeral gully's yield at its mouth to its yield at its receiving stream. Blank defaults to the HUSLE equation.	"Delivery_Ratio	(nd)	Blank, 0. to 1.	F10	2	20
Manning's "n" —Roughness coefficient for concentrated flow within the gully. Blank defaults to 0.040.	"Mannings_n"		Blank, or 0.005 to 1.000	F10	2	21
Re-Plant Period —numbers of days between planting and sufficient crop development that, if an ephemeral gully developed, the eroded area would be replanted. Any tillage operation that distributes the surface layer resets any ephemeral gully voids to zero. Blank defaults to 30 days.	"Replant_Perio d"	{days} [days]	Blank, or 0 to 365	I10	2	22
Organic Carbon —used to calibrate ephemeral gully organic carbon for this ephemeral gully only. Blank defaults to the watershed-scale organic carbon from ephemeral gully sources calibration factor in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"OC_Calib_Fctr	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	23
Nitrogen —used to calibrate ephemeral gully nitrogen for this ephemeral gully only. Blank defaults to the watershed-scale nitrogen from ephemeral gully sources calibration factor in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Calib_Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	24
Phosphorus —used to calibrate ephemeral gully phosphorus for this ephemeral gully only. Blank defaults to the watershed-scale phosphorus from ephemeral gully sources calibration factor in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Calib_Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	25

Description	Field Header for Becord #1	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.
Erosion —used to calibrate ephemeral gully erosion for this ephemeral gully only. Blank defaults to the watershed-scale sediment from ephemeral gully sources calibration factor in the PL CALIBRATION DATA section. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be	Kecord #1 "Erosion_Calib_ Fctr"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	26
exactly the value of this real-number. Headcut Migration Barrier —the sum of the drainage areas of all immediate upstream barriers to headcut migration which, if any limit exists, may be due to the sum of: (a) upstream gullies; and (b) any natural geologic man-made barriers such as a rock outcrop, rock chute, or a drop-structure. Immediate upstream limits are the downstream most upstream gully(s) or blockages that are in parallel with and upstream from this gully. Blank defaults to internal calculation for approximate initial plot-size sheet & rill area (initial 72.6 ft. flow paths).	"Headcut_Migr ation_Barrier"	{acres} [hectares]	Blank, 0. to drainage area at mounth	F10	2	27
Headcut detachment leading coefficient "a"—a local leading coefficient used to determine the headcut detachment(K _{ch}) which is used to calculate headcut migration. The headcut detachment equation is: $K_{ch} = a \square e^{(b \square r_c)}$ The detachment & erodibility options share this field. If the detachment option has been activated, a blank field defaults to the global headcut detachment's leading coefficient. If non-blank and the detachment option has been activated, "a" must be equal to or greater than 0.	"Headcut_Dtach /Erod_Coef_a"	$\begin{array}{c} K_{ch} \text{ in} \\ \{lb_mass/s \\ / \ lb_force\} \\ [g/s-N]; \ \tau_c \\ in \\ \{psi\} \\ [Pa] \end{array}$	Blank, a≥0	F10	2	28
Headcut erodibility leading coefficient "a"—If the erodibility's global fields have been activated, then this is the local leading coefficient used to determine the headcut erodibility(K _d) which is used to calculate headcut migration. The headcut erodibility equation is: $K_d = a \Box \tau_c^{\ b}$ The detachment & erodibility options share this field. If the erodibility option has been activated, a blank field defaults to the global headcut erodibility's leading coefficient. If non-blank and the erodibility option has been activated, "a" must be equal to or greater than 0	"Headcut_Dtach /Erod_Coef_a"	$\begin{array}{c} K_{d} \ in \\ \{in^{3}\!/\!s/ \\ lb_force\} \\ [cm^{3}\!/\!s\text{-}N]; \\ \tau_{c} \ in \\ \{psi\} \\ [Pa] \end{array}$	Blank, a≥0	F10	2	28
Headcut detachment exponent coefficient "b"—a local exponent coefficient used to determine the headcut detachment(K _{ch}) which is used to calculate headcut migration. The headcut detachment equation is: $K_{ch} = a \square e^{(b \square r_c)}$ The detachment & erodibility options share this field. If the detachment option has been activated, a blank field defaults to the global headcut detachment's exponent coefficient. If non-blank and the detachment option has been activated, "b" must be equal to or less than 0.	"Headcut_Dtach /Erod_Exp_Coef _b"	$\begin{array}{c} K_{ch} \text{ in} \\ \{lb_mass/s \\ / \ lb_force\} \\ [g/s-N]; \tau_c \\ in \\ \{psi\} \\ [Pa] \end{array}$	Blank, b≤0	F10	2	29

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Headcut erodibility exponent coefficient "b" — If the erodibility's global fields have been activated, then this is the local exponent coefficient used to determine the headcut erodibility(K _d) which is used to calculate headcut migration. The headcut erodibility equation is: $K_d = a \Box \tau_c^{\ b}$ The detachment & erodibility options share this field. If the erodibility option has been activated, a blank field defaults to the global headcut erodibility's exponent coefficient. If non-blank and the detachment option has been activated, "b" must be equal to or less than 0.	"Headcut_Dtach /Erod_Exp_Coef _b"	$\begin{array}{c} K_{d} \mbox{ in } \\ \{\mbox{in}^{3}\mbox{/s}\ \\ \mbox{lb_force} \} \\ [\mbox{cm}^{3}\mbox{/s}\ N]; \\ \tau_{c} \mbox{ in } \\ \{\mbox{psi} \} \\ [\mbox{Pa}] \end{array}$	Blank, b ≤ 0	F10	2	29
Maximum Buffer Trapping Efficiency "TE-m"—Maximum buffer trapping efficiency for the gully. (Blank defaults to: if a buffer is present in the cell or reach that contains the gully then the maximum buffer trapping efficiency determined for the entire buffer would be applied to the gully; or if no buffer is present then the maximum buffer trapping efficiency is 0.)	"Max_Trapping _Efficiency"	[nd]	Blank, 0. to 1.	F10	2	30
Width Algorithm —Flag to indicate if WELLS' (2013) gully width equation #8 may be used as a candidate for this ephemeral gully width. Default for this width algorithm is the watershed flag for WELLS' (2013) gully width equation (h).	"Width_Wells_ Eq.8"		Blank, or T/F	L1	2	31
Width Algorithm—reserved	"Width_reserve d_i"		Blank, or T/F	L1	2	32
Width Algorithm— reserved	"Width_ reserved_j"		Blank, or T/F	L1	2	33
Width Algorithm— reserved	"Width_ reserved_k"		Blank, or T/F	L1	2	34
Input Units Code— Code identifying whether input is in English or SI units. 0 = English ,1 = SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"	Cat	Table of Conten	II	2	35

FEEDLOT DATA Optional

Opti	onai						
Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.				А	1	1-25	
The following record repeats for the number of feedlots. Multiple feedlots for a given cell outlet should be entered as consecutive sets.							
Feedlot ID—Alphanumeric string identifying the feedlot.	"Feedlot_ID"			A10	2	1	
Feedlot Manage ID —Alphanumeric string identifying the feedlot management schedule for the feedlot. Must be the same as a feedlot management ID (in Feedlot Management Data). Leave blank to use initial feedlot conditions as steady state values.	"Feedlot_Mgmt _ID"			A10	2	2	
Open Area—Total open (uncovered) area of the feedlot.	"Open_Area"	{acres} [hectares]	{0.0 to 9884.} [0.0 to 4000.0]	F10	2	3	
Paved Ratio—Paved open feedlot area to total open feedlot area ratio.	"Paved_Ratio"		0.00 to 1.00	F10	2	4	
Roof Area —Total roofed area (in all cells) for feedlot.	"Roof_Area"	{acres} [hectares]	{0.0 to 9884.} [0.0 to 4000.0]	F10	2	5	
Upslope Area —Total area (in all cells) upslope of feedlot whose runoff drains across the feedlot.	"Upslope_Area"	{acres} [hectares]	{0.0 to 9884.} [0.0 to 4000.0]	F10	2	6	

	Field			For	Record	Field
Description	Header for Record #1	Units	Domain	mat	No.	No.
Feedlot Initial N—Initial daily Nitrogen production for the feedlot	"Initial_N"	{lb / day /	{0.0 to 999.}	F10	2	7
expressed on a per unit area basis.		acre}	[0.0 to 1120.0]			
		hectare]				
Feedlot Initial P—Initial daily Phosphorus production for the	"Initial_P"	{lb / day /	{0.0 to 999.}	F10	2	8
feedlot expressed on a per unit area basis.		acre}	[0.0 to 1120.0]			
		[kg / day / hectare]				
Feedlot Initial OrgC—Initial daily organic Carbon production for	"Initial_OC"	{lb / day /	{0.0 to 9992.}	F10	2	9
the feedlot expressed on a per unit area basis.		acre}	[0.0 to 11200.0]			
		[kg / day /				
Delta N —Daily increase in Nitrogen production for the feedlot	"Delta N"	Alb / day	{0.0 to 999 }	F10	2	10
expressed on a per unit area basis.	_	/ acre}	[0.0 to 1120.0]	110	2	10
		[∆kg / day				
	"D-H- D"	/ hectare]		F10	2	11
Delta P —Daily increase in Phosphorus production for the feedlot expressed on a per unit area basis	"Delta_P"	$\{\Delta lb / day$	$\{0.0 \text{ to } 999.\}$	F10	2	11
expressed on a per unit area basis.		$\int \operatorname{acre}_{\mathcal{F}}$	[0.0 10 1120.0]			
		/ hectare]				
Delta OrgC —Daily increase in organic Carbon production for the	"Delta_OC"	$\Delta b / day$	{0.0 to 999.}	F10	2	12
feedlot expressed on a per unit area basis.		/ acre}	[0.0 to 1120.0]			
		[$\Delta kg / day$ / hectare]				
Feedlot Max N—Maximum concentration of Nitrogen in feedlot	"Max_N"	ppm	0.0 to 1000.0	F10	2	13
runoff.		11			_	
Feedlot Max P—Maximum concentration of Phosphorus in feedlot	"Max_P"	ppm	0.0 to 1000.0	F10	2	14
runoff. Feedlet Max OrgC Maximum concentration of organic Carbon in	"Max OC"	nnm	0.0 to 10000.0	E10	2	15
feedlot runoff.	Max_00	ppm	0.0 10 10000.0	110	2	15
Feedlot Pack N—Initial amount of Nitrogen in the feedlot manure	"Pack_N"	{lb/acre}	{0.0 to 9992.}	F10	2	16
pack expressed on a per unit area basis.		[kg /	[0.0 to 11200.0]			
Feedlot Pack P—Initial amount of Phosphorus in the feedlot		{lb / acre}	{0.0 to 9992 }	F10	2	17
manure pack expressed on a per unit area basis.	"Pack_P"	[kg /	[0.0 to 11200.0]	110	2	17
· · ·		hectare]				
Feedlot Pack OrgC—Initial amount of organic Carbon in the	"Pack_OC"	{lb / acre}	{0.0 to 9992.}	F10	2	18
reediot manure pack expressed on a per unit area basis.		[Kg / hectare]	[0.0 to 11200.0]			
Organic Carbon Calibration Factor—used to calibrate feedlot	"OC_Calib_Fctr	(nd)	Blank, 0.0 or	F10	2	19
organic carbon for this feedlot only. Defaults to the watershed-scale	"	()	.000000001 to		_	
organic carbon from feedlot sources calibration factor in the PL			9999999999.			
CALIBRATION DATA section. Nitrogen Calibration Factor—used to calibrate feedlot nitrogen	"N Calib Fctr"	(nd)	Blank 0.0 or	F10	2	20
for this feedlot only. Defaults to the watershed-scale nitrogen from		(IIC)	.000000001 to	110	2	20
feedlot sources calibration factor in the PL CALIBRATION DATA			9999999999.			
section.	"D Calib Eat"	(1)	D1 1 0 0	F10	2	21
Phosphorus Calibration Factor —used to calibrate feedlot phosphorus for this feedlot only Defaults to the watershed-scale	"F_Cand_Fet	(nd)	Blank, 0.0 or 000000001 to	F10	2	21
phosphorus for this reduct only. Dentation to the waterbled source phosphorus from feedlot sources calibration factor in the PL			9999999999.			
CALIBRATION DATA section.	(P		D1			
Erosion Calibration Factor —used to calibrate feedlot erosion for this feedlot only. Defaults to the watershed coals addiment from	"Erosion_Calib_ Fctr"	(nd)	Blank, 0.0 or	F10	2	22
feedlot sources calibration factor in the PL CALIBRATION DATA			.00000001 to 9999999999.			
section.						
Feedlot Cell ID—Alphanumeric string identifying which cell	"Cell_ID"			A10	2	23
contains the feedlot.	"Call Buffor I o	(0)	(0.0.4, 0.9.4.)	E10	2	24
defaults to 0, i.e., no buffer length thus no buffer area is considered)	ngth"	{n} [m]	$\{0.0 \text{ to } 984.\}$	F10	2	24

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}, 1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	25
Go to Layout Matrix	Go to Table of Contents				-	

FEEDLOT MANAGEMENT DATA

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.	-			А	1	1-12
The following record is repeated for each operation in the fee management schedule must	edlot managen be in sequent	ient schedul ial time ordo	e. Multiple opera er.	tions	for a feed	lot
Feedlot Management ID—Alphanumeric string identifying the feedlot management schedule.	"Mgmt_ID"			A10	2	1
Feedlot Operation Date —Month, day and year the feedlot operation occurs. Year is relative to feedlot management schedule. Blank year defaults to 1.	"Month" "Day" "Year"	mm dd yyyy	mm—1 to 12 dd—1 to 31 yyyy—1 to 1000	I2, I2, I4	2	2-4
Pack Remove Ratio —Ratio of feedlot manure pack removed by scraping operation. (Leave blank for animal operations)	"Pack_Remove_ Ratio"		Blank, or 0.0 to 1.0	F10	2	5
Pack Start N —Starting daily rate for Nitrogen produced by all animals changed on the feedlot with this operation. (Leave blank for scraping operation)	"Pack_Start_N"	{lb / day} [kg / day]	Blank, or {-88105. to 88105.} [-40000.0 to 40000.0]	F10	2	6
Pack Start P —Starting daily rate for Phosphorus produced by all animals changed on the feedlot with this operation (Leave blank for scraping operation)	"Pack_Start_P"	{lb / day} [kg / day]	Blank, or {-88105. to 88105.} [-40000.0 to 40000.0]	F10	2	7
Pack Start OrgC —Starting daily rate for organic Carbon produced by all animals changed on the feedlot with this operation. (Leave blank for scraping operation)	"Pack_Start_O C"	{lb / day} [kg / day]	Blank, or {-88105. to 88105.} [-40000.0 to 40000.0]	F10	2	8
Pack Change N —Daily rate change for Nitrogen produced by all animals changed on the feedlot with this operation. (Leave blank for scraping operation)	"Pack_Change_ N"	$\begin{array}{c} \{\Delta \ \text{lb} \ / \\ \text{day} \} \\ [\Delta kg \ / \ \text{day}] \end{array}$	Blank, or {-881. to 881.} [-400.0 to 400.0]	F10	2	9
Pack Change P —Daily rate change for Phosphorus produced by all animals changed on the feedlot with this operation. (Leave blank for scraping operation)	"Pack_Change_ P"	$\begin{array}{c} \{\Delta \ \text{lb} \ / \\ \text{day} \} \\ [\Delta kg \ / \ \text{day}] \end{array}$	Blank, or {-881. to 881.} [-400.0 to 400.0]	F10	2	10
Pack Change OrgC —Daily rate change for organic Carbon produced by all animals changed on the feedlot with this operation. (Leave blank for scraping operation)	"Pack_Change_ OC"	$ \begin{array}{c} {\Delta \overline{lb} /} \\ {day} \\ [\Delta kg / day] \end{array} $	Blank, or {-881. to 881.} [-400.0 to 400.0]	F10	2	11
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	12
<u>Go to Layout Matrix</u>		<u>Go t</u>	o Table of Conten	<u>ts</u>		

FERTILIZER APPLICATION DATA

Optional unless referenced in Management Schedule Data

		-	-	_	-	-
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-6
The following record repeats repeats for	or the number	of fertilizer	applications.			
Fertilizer Application ID—Alphanumeric string identifying the fertilizer application.	"Application_ID "			A10	2	1
Fertilizer Name ID —Alphanumeric string that is the name of the fertilizer or type manure. Must be the same as a Fertilizer reference ID (in Fertilizer Reference Data).	"Name_ID"			A20	2	2
Fertilizer Rate—Fertilizer application rate	"Application_R ate"	{lb / acre} [kg /hectare]	{0.0 to 49974.} [0.0 to 56000.0]	F10	2	3
Fertilizer Depth—Fertilizer application depth in the soil. Zero indicates fertilizer is applied to the soil surface. Currently not used – Reserved.	"Depth"	{in} [mm]	Blank or {0.0 to 59.} [0.0 to 1500.0]	F10	2	4
Fertilizer mixing code—Code indicating whether fertilizer is mixed uniformly between the soil surface and the depth of incorporation. Acceptable values are: N—no, Y—yes (Blank indicates yes)	"Mixing_Code"		Blank, Y or N	A10	2	5
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}, 1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	6
Go to Layout Matrix		<u>Go to</u>	Table of Conten	<u>ts</u>		

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

FERTILIZER REFERENCE DATA

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-13
The following record repeats for t	he number of	fertilizer re	ferences.			
Fertilizer Reference ID—Alphanumeric string that is the name of the fertilizer. Could also be a manure type.	"Reference_ID"			A20	2	1
Fertilizer Nitrite —Fertilizer fraction which is nitrite (NO ₂) to total amount. Currently not used – Reserved.	"Nitrite"	wt /wt (nd)	Blank	F10	2	2
Fertilizer Nitrate —Fertilizer fraction which is nitrate (NO ₃) to total amount. Currently not used – Reserved.	"Nitrate"	wt /wt (nd)	Blank	F10	2	3
Fertilizer Inorganic N —Fertilizer fraction which is mineralizable (inorganic) Nitrogen. Blank defaults to 0.	"Inorganic_N"	wt /wt (nd)	Blank or 0.0 to 1.0	F10	2	4
Fertilizer Organic N—Fertilizer fraction which is organic Nitrogen. Blank defaults to 0.	"Organic_N"	wt /wt (nd)	Blank or 0.0 to 1.0	F10	2	5
Fertilizer Ammonia —Fertilizer fraction which is ammonia (NH ₄). Currently not used – Reserved.	"Ammonia"	wt /wt (nd)	Blank or 0.0 to 1.0	F10	2	6
Fertilizer Mineral Ammonia—Fertilizer fraction which is mineralizable ammonia. Currently not used – Reserved.	"Mineral_Amm onia"	wt /wt (nd)	Blank	F10	2	7
Fertilizer Elemental P —Fertilizer fraction which is phosphorus (P) of any elemental P form. Currently not used – Reserved.	"Elemental_P"	wt /wt (nd)	Blank	F10	2	8
Fertilizer Soluble P —Fertilizer fraction which is soluble Phosphorus to total amount. Currently not used – Reserved.	"Soluble_P"	wt /wt (nd)	Blank	F10	2	9
Fertilizer Inorganic P —Fertilize fraction which is mineralizable (inorganic) Phosphorus. Blank defaults to 0.	"Inorganic_P"	wt /wt (nd)	Blank or 0.0 to 1.0	F10	2	10

Optional unless referenced in Fertilizer Application Data

AnnAGNPS Input File Specifications

		•	
Revision: December 4, 2019	File Name: In	nput Specifications	v5.51 2019.xx.xx.docx

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Fertilizer Organic P—Fertilizer fraction which is organic	"Organic_P"	wt /wt	Blank or	F10	2	11
Phosphorus. Blank defaults to 0.		(nd)	0.0 to 1.0			
Fertilizer Organic Matter—Fertilizer fraction which is organic	"Organic_Matte	wt /wt	Blank or	F10	2	12
matter. Blank defaults to 0.	r"	(nd)	0.0 to 1.0			
Fertilizer Consistency code—Fertilizer (or manure) consistency	"Consistency_C		Blank or	I1	2	13
code. Acceptable values are: 1= Liquid, 2 = Slurry, 3 =	ode"		1, 2, or 3			
Solid						
Currently not used – Reserved.						
Go to Layout Matrix	Go to Table of Contents					

FIELD POND DATA

Optional

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-22
The following record repeats	for the numbe	r of field po	nds.			
Field Pond ID —unique alphanumeric string identifying the field pond. Multiple field ponds within the same cell may be aggregated and entered as a single pond for simulation convenience.	"Pond_ID"			A10	2	1
Field Pond-Cell ID —alphanumeric string identifying cell that contains the field pond(s). Must be the same as a cell ID in the CELL DATA section already included within the watershed.	"Cell_ID"			A10	2	2
Field Pond area —area of field pond(s). Multiple field ponds in the same cell may be aggregated together as a single field pond for convenience.	"Pond_Area"	{acres} [hectares]	{0.0—9884.} [0.0—4000.0]	F10	2	3
Number of rotation years—number of years in rotation for the management of this field pond.	"Number_of_Ro tation_Years"		< 10000	I10	2	4
Number gate operations —total number of field pond gate openings & closures within the rotation period for this field pond. For every opening (closing) there has to be a closing (opening).	"Number_of_Ga te_Operations"		2—10000	I10	2	5
Delivery Ratio	"Delivery_Ratio		0.0 - 1.0	10	2	6
Volume of release water —amount of water released from field pond per gate open operation; depth in linear units. May be left blank if release rate is entered.	"Volume_of_Rel ease_Water"	{in} [mm]	{>0393.} [>0-10000.]	F10	2	7
Drain time —time to drain field pond's release. Will be used if the release rate field is blank. A blank in both the drain time & release rate fields will default to a 24-hr drain time.	"Drain_Time"	hr	> 0.0 - 8784.	F10	2	8
Release rate —rate of field pond release as depth in linear units per hour. May be left blank, in which case the rate will be calculated from the values for the volume of release water (and pond area) and drain time. A blank in both the drain time & release rate fields will default to a 24-hr drain time.	"Release_Rate"	{in/hr} [mm/hr]	{>0 393.} [>0 10000.]	F10	2	9
Sediment concentration —average total sediment concentration in release water. Blank field defaults to 0 ppm.	"Sediment_Con c"	ppm	0. – 1000000.	F10	2	10
Clay content —percent of clay content in sediment yield from the field pond. Clay plus silt contents must not add up to more than 100 %. Blank fields for both clay & silt contents defaults to 100% clay. A blank field for clay with a silt content greater than 0%, defaults to 0% content for clay. If the sum of clay & silt content adds up to less than 100%, the difference is assumed to be sand.	"Clay_Content"	%	0. – 100.	F10	2	11
Silt content —percent of silt content in sediment yield from the field pond. Clay plus silt contents must not add up to more than 100 %. If sum adds up to less than 100%, the difference is assumed to be sand. Blank field defaults to 0% for silt.	"Silt_Content"	%	0. – 100.	F10	2	12

 AnnAGNPS Input File Specifications

 2019
 File Name: Input_Specifications_v5.51_2019.xx.xx.docx
 Revision: December 4, 2019

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Nitrogen concentration —total concentration of nitrogen (both dissolved & attached) in release water. Blank defaults to 0. Currently not used – Reserved.	"N_Conc"	ppm	Blank or 0. – 1000000.	F10	2	13
Phosphorus concentration—total concentration of Phosphorus (both dissolved & attached) in release water. Blank defaults to 0. Currently not used – Reserved.	"P_Conc"	ppm	Blank or 0. – 1000000.	F10	2	14
Organic carbon concentration—total concentration of organic carbon (both dissolved & attached) in release water. Blank defaults to 0. Currently not used – Reserved.	"OC_Conc"	ppm	Blank or 0. – 1000000.	F10	2	15
Pesticide reference id —ID of the pesticide in the release water in pesticide reference list. ID must be in the pesticide reference list. ID will be converted to the array index of the pesticide in the pesticide reference list. Only one pesticide per field in each field pond release is allowed. Currently not used – Reserved.	"Pesticide_Refer ence_ID"			A40	2	16
Pesticide concentration —total concentration of pesticide (both dissolved & attached) in release water. Only one pesticide allowed per field pond. Blank defaults to 0. Currently not used – Reserved.	"Pesticide_Conc "	ppm	Blank or 0. – 1000000.	F10	2	17
Organic Carbon Calibration Factor —used to calibrate field pond organic carbon for this field pond only. Defaults to the watershed- scale organic carbon from field pond sources calibration factor in the PL CALIBRATION DATA section.	"OC_Calib_Fctr "	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	18
Nitrogen Calibration Factor —used to calibrate field pond nitrogen for this field pond only. Defaults to the watershed-scale nitrogen from field pond sources calibration factor in the PL CALIBRATION DATA section.	"N_Calib_Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	19
Phosphorus Calibration Factor —used to calibrate field pond phosphorus for this field pond only. Defaults to the watershed-scale phosphorus from field pond sources calibration factor in the PL CALIBRATION DATA section.	"P_Calib_Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	20
Erosion Calibration Factor —used to calibrate field pond erosion for this field pond only. Defaults to the watershed-scale sediment from field pond sources calibration factor in the PL CALIBRATION DATA section.	"Erosion_Calib_ Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	21
Input Units Code— Code identifying whether input is in English or SI units. 0 = English ,1 = SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"	Cat	Table of Contra	Il	2	22

FIELD POND OPERATIONS DATA

Optional unless FIELD POND DATA is present.

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-5
The following record repeats for the	ne number of f	ield pond op	erations.			
Field Pond ID —unique alphanumeric string identifying the field pond. Multiple field ponds within the same cell may be aggregated and entered as a single pond for simulation convenience.	"Pond_ID"			A10	2	1

AnnAGNPS Input File Specifications

		·	
Revision: December 4, 2019	File Name: Input	Specifications v5	5.51 2019.xx.xx.docx
	1		

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.		
Open/close —gate action as to whether the gate is opened (water release) or closed. The time period (days) during which gate is opened (before it is closed), the field pond will be treated as a part of the homogeneous cell. The time period during which the gate is closed (before it is opened again), will be treated as no runoff; i.e., all rainfall is assumed to be captured within the pond and infiltrates into the ground.	"Open/Close_G ate_Action"		'open' or 'close'	A10	2	2		
Opening/closure rotation day —Month, day, and relative year (within a set of field pond operations) for gate openings & closures. This day will be converted internally from a relative rotation date (mm/dd/ry) to a 2-dimensional variable which will be the rotation year & Julian day for the release/gate closure. If the operation month and day are blank, and this operation is the first in a set of field pond operations grouping, then this operation is used as initial conditions for starting the simulation. Blank year defaults to 1.	"Open/Close_Ro tation_Month", "Open/Close_Ro tation_Day", "Open/Close_Ro tation_Year"	mm/ dd/ yy	Blank, or 1—12 1—31 1—99	I2, I2, I2	2	3-5		
Go to Layout Matrix	Go to Table of Contents							

GEOLOGY DATA

Optional unless referenced by Cell Data or Global IDs, Factors, and Flags Data											
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.					
Field Header— Required unique field header for each field listed below.				А	1	1-12					
The following record repeats for the number of geology sets.											
Geology ID—Alphanumeric string identifying the geology set.	"Geology_ID"			A10	2	1					
Delay Time —Drainage time of the overlying geologic formations (days). Blank defaults to 3 days.	"Delay_Time"	days	Blank, 1 to 365	I10	2	2					
Water Table —Depth of water table. The depth at which the water pressure equals atmospheric pressure.	"Water_Table"	{ft} [m]	{0.0 to 98.424} [0.0 to 30.0]	F10	2	3					
Aquifer Saturated Hydraulic Conductivity—Hydraulic conductivity of the aquifer. Blank defaults to the hydraulic conductivity of soil layer 2.	"Aquifer_Sat_H yd_Conduct"	{in/day} [mm/day]	Blank or {0.0 to 99999999.9} [0.0 to 30480000.0]	F10	2	4					
K-vadose Saturated Hydraulic Conductivity —Hydraulic conductivity of the vadose zone. Blank defaults to the hydraulic conductivity of soil layer 2.	"Vadose_Sat_H yd_Conduct"	{in/day} [mm/day]	Blank or {0.0 to 99999999.9} [0.0 to 30480000.0]	F10	2	5					
Aquifer Porosity—Porosity of the aquifer. Blank defaults to the porosity of soil layer 2.	"Porosity"		Blank or 0.00001 to 1.0	F10	2	6					
Aquifer Field Capacity—Field capacity of the aquifer. Blank defaults to the field capacity of soil layer 2.	"Field_Capacity "		Blank, or 0.00001 to 1.0	F10	2	7					
Aquifer Specific Yield—Specific yield of the aquifer. It is also called drainable pore space. It can be approximately calculated as the difference between the porosity and the field capacity.	"Specific_Yield"		Blank, or 0.00001 to 1.0	F10	2	8					
Aquifer Thickness—Thickness of the aquifer from the water table to the bottom of the aquifer. Blank defaults to 30 m.	"Thickness"	{ft} [m]	Blank or {0.0 to 99999999.9} [0.0 to 30480000.0]	F10	2	9					
Aquifer Soluble Nitrogen—Concentration of nitrate in groundwater contribution to stream flow from sub-basin. Blank defaults to 0.0.	"Soluble_N"	{ppm} [mg/L]	Blank or {0.0 to 100.0} [0.0 to 100.0]	F10	2	10					
Aquifer Soluble Phosphorus—Concentration of phosphorus in groundwater contribution to stream flow from sub-basin. Blank defaults to 0.0.	"Soluble_P"	{ppm} [mg/L]	Blank or {0.0 to 100.0} [0.0 to 100.0]	F10	2	11					
AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

Description	Field Header for Record #1	Units {English}	Domain {English} [SI]	For mat	Record No.	Field No.
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	12
Go to Layout Matrix		Gote	Table of Conten	ts		

GLOBAL ERROR AND WARNING LIMITS DATA

		Opt	ional					
			Field	Units	Domain	For	Record	Field
	Description		Header for	{English}	(English) [SI]	mot	No	No
			Record #1	[SI]	{English} [51]	mat	INO.	INO.
Field Header— Rec	uired unique field head	er for each field listed				Α	1	1-5
below.								
	The following recor	d repeats for the numb	er of global eri	or & warni	ng sets requested.		<u>.</u>	_
Keyword ID—Alph	anumeric string identify	ing the error and	"Keyword_ID"			A10	2	1
warning record ID. C	Only warning limits can	be changed at this time.						
Allowable keywords are:								
	_							
Keyword ID	Descr	iption						
"CLB EROS"	erosion calibration fac	tor						
"CLB NIT"	nitrogen calibration fa	ctor						
"CLB ORGC"	organic carbon calibra	tion factor						
"CLB PHOS"	phosphorus calibration	n factor						
"CLB RCN"	RCN calibration factor	r						
"DA"	drainage area							
"ELE"	elevation							
"MAN N"	Manning's N							
"SLP"	slope							
"TC"	time of concentration							
Warning Minimum	-Value defined to be t	he minimum allowable	"Warning_Min"		Blank or	F10	2	2
threshold below which	ch an AnnAGNPS warn	ing message will be			{error minimum			
generated. This value	e must be between the en	rror minimum and			to maximum}			
maximum limits defi	ined within AnnAGNPS	. Blank defaults to the			[error minimum			
initial warning minin	num defined within Anr	AGNPS.			to maximum]			
Keyword ID	Default	AnnAGNPS						
Rey word ID	Warning Minimum	Error Minimum						
"CLB EROS"	0.5	0.0						
"CLB NIT"	0.5	0.0						
"CLB ORGC"	0.5	0.0						
"CLB PHOS"	0.5	0.0						
"CLB RCN"	0.5	0.0						
"DA"	0.09	.000000001						
"ELE"	0.0	-300.0						
"MAN N"	0.01	0.01						
"SLP"	0.0001	.00001						
"TC"	0.1	0.0						

AnnAGNPS Input File Specifications

		-	
Revision: December 4, 2019	File Name: I	nput_Specifications	_v5.51_2019.xx.xx.docx

	Description		Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Warning Maximum—Value defined to be the maximum allowable threshold above which an AnnAGNPS warning message will be generated. This value must be between the error minimum and maximum limits defined within AnnAGNPS. Blank defaults to the initial warning maximum defined within AnnAGNPS.		"Warning_Max "		Blank or {error minimum to maximum} [error minimum to maximum]	F10	2	3	
Keyword ID "CLB EROS" "CLB NIT" "CLB ORGC" "CLB PHOS" "CLB RCN" "DA" "ELE" "MAN N" "SLP" "TC"	Default Warning Maximum 2.0 2.0 2.0 2.0 2.0 100.0 5000.0 0.5 1.0 24.0	AnnAGNPS Error Maximum 9999999999 9999999999 9999999999 999999						
Error Minimum—Value defined to be the minimum allowable threshold below which an AnnAGNPS error message will be generated. This value must be between the error minimum and maximum limits defined within AnnAGNPS. Blank defaults to the initial error minimum defined within AnnAGNPS.		"Error_Min"		Blank or {error minimum to maximum} [error minimum to maximum]	F10	2	4	
Currently not used – Reserved. Error Maximum —Value defined to be the maximum allowable threshold above which an AnnAGNPS error message will be generated. This value must be between the error minimum and maximum limits defined within AnnAGNPS. Blank defaults to the initial error maximum defined within AnnAGNPS. Currently not used – Reserved.		maximum allowable message will be ror minimum and . Blank defaults to the GNPS.	"Error_Max"		Blank or {error minimum to maximum} [error minimum to maximum]	F10	2	5
	Go to Layout Matrix	x		<u>Go to</u>	<u>o Table of Conten</u>	ts		

⁴⁵GLOBAL IDS, FACTORS, AND FLAGS DATA: Optional

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-57
The following reco	rd is non-repe	ating		_		
Headcut detachment leading coefficient (a) —a global leading coefficient used to determine the headcut detachment(K _{ch}) which is used to calculate headcut migration. The headcut detachment equation is: $K_{ch} = a \Box e^{(b \Box r_c)}$ If the headcut erodibility fields are not activated, a blank field will default to "29.1". If non-blank, the detachment option is activated, the erodibility option may not be activated, and "a" must be equal to or greater than 0. If all fields for the detachment and erodibility options are blank, the detachment option is activated with its defaults.	"Hdct_Detachm ent_Coef_a"	$\begin{array}{c} K_{ch} \mbox{ in } \\ \{ lb_mass/s \\ /lb_force \} \\ [g/s/N]; \\ \tau_c \mbox{ in } \\ \{ lb/ft^2 \} \\ [Pa] \end{array}$	a ≥ 0	F10	2	1

⁴ This section must be placed immediately following the GLOBAL ERROR AND WARNING LIMITS DATA if included; otherwise must follow SIMULATION PERIOD DATA section.

⁵ All global factors & flags are used as defaults for local values but are subsidiary to local values if the local values are explicit input, that is, the local values are not blank fields. This is consistent with the default rule where local values override global values for the local variable.

		.	
Revision: December 4, 2019	File Name:	Input_Specifications_	v5.51_2019.xx.xx.docx

Description	Field Header for	Units	Domain	For	Record	Field
Description	Record #1	{English} [SI]	{English} [SI]	mat	No.	No.
Headcut detachment exponent coefficient (b) —a global exponent coefficient used to determine the headcut erodibility(K _d) which is used to calculate headcut migration. The headcut erodibility equation is: $K_{ch} = a \Box e^{(b\Box r_c)}$ If the headcut erodibility fields are not activated, a blank field will default to "-0.224". If non-blank, the detachment option is activated, the erodibility option may not be activated, and "b" must be equal to or less than 0. If all fields for the detachment and erodibility options are blank, the detachment option is activated with its defaults.	"Hdct_Detachm ent_Exp_Coef_b "		b ≤ 0	F10	2	2
URBAN landuse repair date —Date when any prior urban landuse ephemeral gullying will be repaired. Complete the month, day, & relative year [mm/dd/yyyy] when the repair is to be scheduled. The relative year is interpreted as follows: if yyyy="0001", then repair will take place every year; if yyyy="0002", then repair will take place every other year; if yyyy="0003", then every third year; etc. To ensure no repairs are to be made, set date to "00/00/0000". Default for this repair date is "10/01/0001"	"Urban_Repair _Month", "Urban_Repair _Day", "Urban_Repair _Year"	mm dd yyyy	0 to 12 0 to 31 0 to 9999	I2, I2, I4	2	3-5
CROPLAND landuse repair date —Date when any prior cropland landuse ephemeral gullying will be repaired. Complete the month, day, & relative year [mm/dd/yyyy] when the repair is to be scheduled. The relative year is interpreted as follows: if yyyy="0001", then repair will take place every year; if yyyy="0002", then repair will take place every other year; if yyyy="0003", then every third year; etc. To ensure no repairs are to be made, leave date blank or set date to "00/00/0000". Blank defaults to "00/00/0000".	"Cropland_Rep air_Month", "Cropland_Rep air_Day", "Cropland_Rep air_Year"	mm dd yyyy	0 to 12 0 to 31 0 to 9999	12, 12, 14	2	6-8
FOREST landuse repair date —Date when any prior forest landuse ephemeral gullying will be repaired. Complete the month, day, & relative year [mm/dd/yyyy] when the repair is to be scheduled. The relative year is interpreted as follows: if yyyy="0001", then repair will take place every year; if yyyy="0002", then repair will take place every other year; if yyyy="0003", then every third year; etc. To ensure no repairs are to be made, set date to "00/00/0000". Default for this repair date is "11/01/0001".	"Forest_Repair_ Month", "Forest_Repair_ Day", "Forest_Repair_ Year"	mm dd yyyy	0 to 12 0 to 31 0 to 9999	I2, I2, I4	2	9-11
PASTURE landuse repair date —Date when any prior pasture landuse ephemeral gullying will be repaired. Complete the month, day, & relative year [mm/dd/yyyy] when the repair is to be scheduled. The relative year is interpreted as follows: if yyyy="0001", then repair will take place every year; if yyyy="0002", then repair will take place every other year; if yyyy="0003", then every third year; etc. To ensure no repairs are to be made, set date to "00/00/0000". Default for this repair date is "10/01/0001".	"Pasture_Repai r_Month", "Pasture_Repai r_Day", "Pasture_Repai r_Year"	mm dd yyyy	0 to 12 0 to 31 0 to 9999	I2, I2, I4	2	12-14
RANGELAND landuse repair date —Date when any prior rangeland landuse ephemeral gullying will be repaired. Complete the month, day, & relative year [mm/dd/yyyy] when the repair is to be scheduled. The relative year is interpreted as follows: if yyyy="0001", then repair will take place every year; if yyyy="0002", then repair will take place every other year; if yyyy="0003", then every third year; etc. To ensure no repairs are to be made, set date to "00/00/0000". Default for this repair date is "00/00/0000".	"Rangeland_Re pair_Month", "Rangeland_Re pair_Day", "Rangeland_Re pair_Year"	mm dd yyyy	0 to 12 0 to 31 0 to 9999	12, 12, 14	2	15-17

Description	Field Header for	Units {English}	Domain {English} [SI]	For mat	Record No	Field No
	Record #1	[SI]	{English} [51]	mat	110.	110.
Headcut erodibility leading coefficient (a)—a global leading coefficient used to determine the headcut erodibility(K _d) which is used to calculate headcut migration. The headcut erodibility equation is: $K_d = a \Box \tau_c^{\ b}$ A blank field will default to headcut detachment. If non-blank, the detachment option fields must be blank and are deactivated, and the erodibility option is activated. "a" must be equal to or greater than 0. If all fields for the detachment and erodibility options are blank, the detachment option is activated with its defaults.	"Hdct_Erodibili ty_Coef_a"	$\begin{array}{c} K_{d} \ in \\ \{in^{3}/s / \\ lb_force\} \\ [cm^{3}/s / N]; \\ \tau_{c} \ in \\ \{lb/ft^{2}\} \\ [Pa] \end{array}$	a ≥ 0	F10	2	18
Headcut erodibility exponent coefficient (b)—a global exponent coefficient used to determine the headcut erodibility(K _d) which is used to calculate headcut migration. The headcut erodibility equation is: $K_d = a \Box \tau_c^{\ b}$ If non-blank, the detachment option fields must be blank, the erodibility field for "a" must be non-blank, and "b" must be equal to or less than 0. If "a" is non-blank and this field is blank, "b" will default to "-0.5". If all fields for the detachment and erodibility options are blank, the detachment option is activated with its defaults.	"Hdct_Erodibili ty_Exp_Coef_b"	$\begin{array}{c} K_{d} \mbox{ in } \\ \{\mbox{in}^{3}\mbox{s} / \\ \mbox{lb_force} \} \\ \mbox{[cm}^{3}\mbox{s} / N]; \\ \tau_{c} \mbox{ in } \\ \{\mbox{lb}\mbox{ft}^{2} \} \\ \mbox{[Pa]} \end{array}$	b ≤ 0	F10	2	19
Width Algorithm —Flag to indicate if Nachtergaele et al's (2002) equation may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Nachter gaele"		T/F	L1	2	20
Width Algorithm—Flag to indicate if gully-location's hydraulic geometry equation may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Hydrau lic_Geometry"		T/F	L1	2	21
Width Algorithm —Flag to indicate if non-submerging tailwater at headcut crest equation may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Non- submerging_Tai lwater"		T/F	L1	2	22
Width Algorithm —Flag to indicate if Woodward's (1999) equilibrium gully width equation may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Woodw ards_Equilibriu m"		T/F	L1	2	23
Width Algorithm —Flag to indicate if Woodward's (1999) ultimate gully width equation may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Woodw ards_Ultimate"		T/F	L1	2	24
Width Algorithm —Flag to indicate if Modified Wells' (2013) gully width equation #9 may be used as a candidate for the ephemeral gully width. Default for this width algorithm is false.	"Width_Wells_ Eq.9"		T/F	L1	2	25
Note: See additional available widt	th algorithms b	beginning in	field #52			
Erosion verification file —Flag to indicate if the ephemeral gully erosion file (AnnAGNPS_SIM_Ephemeral_Gully_Erosion.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to true). Default for this verification file is set to true.	"Erosion_Vrfy"		T/F	L1	2	26
Hydrograph & section geometry verification file—Flag to indicate if the ephemeral gully hydrograph & section geometry verification file (AnnAGNPS_SIM_Ephemeral_Gully_Section_&_Hydrograph.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to true). Default for this verification file is set to true.	"Hydrograph_V rfy"		T/F	L1	2	27

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Nickpoint verification file—Flag to indicate if the ephemeral gully nickpoint verification file (AnnAGNPS_SIM_Ephemeral_Gully_Nickpoint.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to	"Nickpoint_Vrf y"		T/F	L1	2	28
Repair dates verification file —Flag to indicate if the ephemeral gully repair dates verification file (AnnAGNPS_SIM_Ephemeral_Gully_Repair_Date.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to true). Default for this verification file is set to true.	"Repair_Dates_ Vrfy"		T/F	L1	2	29
Sediment yield to the gully's mouth verification file—Flag to indicate if the ephemeral gully's sediment yield to its mouth verification file (AnnAGNPS_SIM_Ephemeral_Gully_Yield_to_Mouth.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to true). Default for this verification file is set to true.	"Sed_Vield_to_ Gully_Mouth_V rfy"		T/F	L1	2	30
Sediment yield to its receiving stream verification file—Flag to indicate if the ephemeral gully's sediment yield to its receiving stream verification file (AnnAGNPS_SIM_Ephemeral_Gully_yield.sim) will be included in the output requests. The "Process_Gully" field in the Simulation Verification Files submenu within the OUTPUT OPTIONS DATA menu must also be requested by the user (set to true). Default for this verification file is set to true.	"Sed_Vield_to_ Rcvg_Reach_Vr fy"		T/F	L1	2	31
Minimum Interception Evaporation —maximum amount of expected intercepted precipitation subject to evaporation prior to infiltration (surface retention) at 100% relative humidity. Blank defaults to 1.000 [mm].	"Min_Intercepti on_Evaporation "	{in} [mm]	Blank, or {0.0 to 0.250} [0.0 to 6.350]	F10	2	32
Maximum Interception Evaporation—maximum amount of expected intercepted precipitation subject to evaoporation prior to infiltration (surface retention) at 0% relative humidity. Blank defaults to 2.500 [mm].	"Max_Intercepti on_Evaporation "	{in} [mm]	Blank, or {0.0 to 0.250} [0.0 to 6.350]	F10	2	33
Detention Coefficient "a" —rainfall/runoff detention's constant coefficient to account for depressions (puddles, etc.) Blank defaults to 0.	"Detention_Coef _a"	{in} [mm]	Blank, or {0.0 to 39.37} [0.0 to 1000]	F10	2	34
Detention Coefficient "b" — rainfall/runoff detention's saturated hydraulic conductivity coefficient multiplier to account for the ability of the water to infiltrate into the soil column. Blank defaults to 1.	"Detention_Coef _b"	{in} [mm]	Blank, or {0.0 to 3.937} [0.0 to 100]	F10	2	35
RCN Convergence Tolerance —RCN calibration iterations will continue until the calculated water load at the subwatershed outlet is within +- of the tolerance from the target water load or until the number of iterations reaches the maximum. Blank defaults to 6.35 [mm] (1/4 [in]).	"RCN_Converg ence_Tolerance"	{in} [mm]	Blank or {4.0·10 ⁻⁹ to 1.0} [0.0000001 to 25.4]	F10	2	36
RCN Maximum Number of Iterations —maximum number of iterations before calibration creases regardless of the specified tolerance. Blank defaults to 10.	"RCN_Max_Ite rations"		1 to 100	I10	2	37
Available Soil Moisture Ratio for AMC II—soil column's available soil moisture for antecedent moisture condition II (AMC II). It is the ratio of the available soil moisture, which is the amount of water in the soil column less the amount of pore space at the wilting point, to the difference between the amount of pore space at the field capacity less the amount of pore space at the wilting point. Value must be between 0 & 1; blank defaults to 0.5.	"Avbl_Soil_Moi st_Ratio_AMC_ II"	[nd]	Blank, or 0.0 to 1.0	F10	2	38

				r.		
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Maximum Available Sediment Concentration for Sheet Flow— maximum allowable sediment concentration of sediment yield from sheet flow erosion at the cell's receiving reach.	"Max_Avbl_Sed _Conc_for_Sht_ Flw"	[%]	Blank, or 0.0 to 100.0	F10	2	39
Maximum Available Sediment Concentration for Concentrated Flow—maximum allowable sediment concentration of sediment yield from concentrated flow erosion at the cell's receiving reach.	"Max_Avbl_Sed _Conc_for_Con c_Flw"	[%]	Blank, or 0.0 to 100.0	F10	2	40
Average Annual Unit-Area Baseflow—Average annual baseflow which will be used when reach routing. Currently not used – Reserved.	"AA_Unit_Area _Baseflow"	{in/yr} [mm/yr]	Blank or [0 to 10.} [0 to 254]	F10	2	41
RCN Calibration Only — flag to indicate if execution will only calibrate the runoff curve numbers and not continue with any other loading analyses; true if calibration only, false otherwise. Blank defaults to false.	"RCN_Calib_O nly"		Blank, or T/F	L1	2	42
Calculate Baseflow — Baseflow calculations will be performed if True. No baseflow is calculated if False. Default = False	"Calculate_Base flow"		Blank, or T/F	L1	2	43
FAO ET Enhancement —If true then the FAO (Allen et al., 1998) enhancement for calculating the evapotranspiration will be used. If false then no enhancement will be performed and all enhancement related crop and non-crop parameters will be ignored. Default = True	"FAO_ET_Enh ancement"		Blank, or T/F	L1	2	44
Basal Crop Coefficient Climatic Adjustment code —If true then the basal crop coefficient for "Kcb-mid" and "Kcb-end" will be adjusted, if necessary, based on climate conditions for crop data and non-crop data unless overridden by the local flag in the Crop Data section and the Non-Crop Data section. Default = True	"Basal_Crop_C oef_Climate_Ad just"		Blank, or T/F	L1	2	45
Watershed Storm Type ID—ID indicating one of the preset synthetic or user-requested storm types to use with pre-calculated solutions using extended TR-55: Acceptable preset IDs are: blank which defaults to Std SCS Type II, (1) Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type II; (4) Std. SCS Type III; (5) Std. Uniform; (6) Std. SCS NM60; (7) Std. SCS NM65; (8) Std. SCS NM70; & (9) Std. SCS NM75; other IDs, limited to eleven (11) additional, require user-defined storm type input for both the rainfall distribution & optional unit peak discharge regression coefficients.	"Wshd_Storm_ Type_ID"			A20	2	46
Default Geology ID —This geology set will be used for all cells that do not have a specific Geology ID assigned within the CELL DATA section. Must be the same as a Geology ID in the GEOLOGY DATA section. If blank, a pre-defined Geology set called "geol_set_1" will be used for baseflow calculations for all cells that do not have a Geology ID specified in the CELL DATA section with corresponding Geology IDs in the GEOLOGY DATA section.	"Dflt_Geology_1 D"			A10	2	47

Revision: December 4, 2019	File Name:	Input_Specif	fications_v5.51	2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
 Default Hydraulic Geometry ID—Alphanumeric string identifying the hydraulic geometry data which will be used for the CELL DATA & REACH DATA sections as the default for their Hydraulic Geometry fields. Blank in this field defaults to Curve B. The bulitin data are: Curve A—Mediterranean climate of winter rainfall such as San Francisco region at 30 inches annual precipitation (Leopo&ld et al); Curve B: high-rainfall areas such as Pennsylvania, with annual accumulation precipitation of 45 inches (Leopold et al); Curve C: mountain areas in the Upper Green River, Wyoming (Leopold et al); Curve E—Pacific Maritime Mountainsnorth Cascades, Puget lowland, coast range, & Willamette valley (Castro); Curve F—Western CordilleraKlamath mountains, Cascades, eastern Cascades, Blue mountains, northern Rockies, middle Rockies, and Wasatch & Uinta mountains (Castro); Curve G—Western Interior Basin & RangesColumbia Basin, Snake River Basin/high Desert, northern Basin & range, ands Wyoming Basin (Castro); Curve I—Mission Creek, Oregoneastern slopes of the Umatilla Mountains (Theurer & Pedone); Curve I—Seco Creek, Texasnorth of San Antonio in the Edwards Aquifer (Theurer et al, 1995); Curve K—Wrights Brook & Kiff BrookNew York City water supply watersheds in the West Branch of the Delaware River Basin, New York (Theurer et al, 1995); Curve L—Cole Gully & Wikoff Bayousubwatersheds within the Bayou Plaquemine Brule in the Mermentau River Basin, southwestern Louisiana (Kolian & Theurer, 2001); Curve M—Carneros Creek in Monterey & San Benito Counties, central coast, California (Merkel, 1999). Any other Hydraulic Geometry IDs must be created using the HYDRAULIC GEOMETRY DATA section. 	"Dfl_Hydraulic _Geom_ID"		Blank, Curve A, Curve B, Curve C, Curve D, Curve F, Curve F, Curve G, Curve H, Curve J, Curve J, Curve K, Curve M, or a valid Hydraulic Geometry ID from the HYDRAULIC GEOMETRY DATA section	A10	2	48
Default Initial Soil Conditions ID —ID of the initial soil conditions record set that will be used for all soils that do not have a specific Initial Soil Conditions ID assigned within the SOIL DATA section. If present, must be the same as an Initial Soil Conditions ID in the SOIL INITIAL CONDITIONS DATA section. If blank, pre-defined values will be used.	"Dflt_Init_Soil_ Conditions_ID"			A10	2	49
Default Crop RCN ID —Curve Number ID of the RCN set that will be used as a default for the first event of each "crop" schedule in the MANAGEMENT SCHEDULE DATA. If present, must be the same as a Curve Number ID in the RUNOFF CURVE NUMBER DATA section. If blank, an internally defined curve number set with the ID of "Default Crop CN" will be used. The curve numbers supplied for A, B, C, and D are 72, 81, 88, and 91 respectively.	"Dflt_Crop_RC N_ID"			A40	2	50
Default Non-Crop RCN ID—Curve Number ID of the RCN set that will be used as a default for the first event of each "non-crop" schedule in the MANAGEMENT SCHEDULE DATA. If present, must be the same as a Curve Number ID in the RUNOFF CURVE NUMBER DATA section. If blank, an internally defined curve number set with the ID of "Default Non-Crop CN" will be used. The curve numbers supplied for A, B, C, and D are 68, 79, 86, and 89 respectively.	"Dflt_Non- Crop_RCN_ID"			A40	2	51

	-	-	-	r	r	-
Description	Field	Units	Domain	For	Record	Field
Description	Header for Record #1	{English}	{English} [SI]	mat	No.	No.
	With Will		TE /E	T 1	2	
Width Algorithm—Flag to indicate if WELLS' (2013) gully width	"width_wells_ Eq.8"		T/F	LI	2	52
width Default for this width algorithm is false						
Width Algorithm—reserved	"Width Reserve		T/F	L1	2	53
	d_i"				-	55
Width Algorithm—reserved	"Width_ Reserved_i"		T/F	L1	2	54
Width Algorithm—reserved	"Width_ Reserved k"		T/F	L1	2	55
Critical Shear Stress—Critical shear stress at which gully erosion	"Critical_Shear	$\{lbs/ft^2\}$	Blank, or	F10	2	56
begins. If a value is supplied here, it is used as a global default for	_Stress"	$[N/m^2]$	{>0 to 2.05}			
all gullies that do not have a local critical shear stress value supplied			[>0 to 100.0]			
in the Ephemeral Gully Data section. Blank means that for all gullies						
that do not have a local critical shear stress value supplied, a critical						
shear stress value will be internally calculated based upon the gully's clay silt & sand content as determined using the gully's soil						
ID.						
RUSLE2 —This serves as a toggle (on/off) switch for RUSLE2.	"RUSLE2_		T/F	L1	2	57
If this parameter is true (on) and	Flag"					
1.) A default RUSLE2 ID is specified in this section then the						
RUSLE2 parameters selected in that RUSLE2 ID's record						
will be used for all cells. If the cell's RUSLE2 ID is						
present then it will be used; otherwise, the default						
RUSLE2 ID will be used.						
2.) A default RUSLE2 ID is -not- specified in this section then only cells with a specified RUSLE2 ID will use						
RUSLE2 data: otherwise RUSLE2 will not be used						
If this parameter is false (off) then RUSLE2 will not be used for any						
cells even if RUSLE2 IDs are specified.						
Default = false.						
Default RUSLE2 ID —ID of the RUSLE2 set that will be used as a	"Dflt_RUSLE2_			A40	2	58
global default. If present, must be the same as a RUSLE2 ID in the	ID"					
RUSLE2 DATA section.						
If the RUSLE2 flag is true and:						
1.) This parameter is specified then it will be used for all cells						
 with no locally specified KUSLE2 ID. This parameter is -not- specified then only cells with a 						
2.) This parameter is -not- specified then only cens with a locally specified RUSI F2 ID will use RUSI F2						
parameters as selected in the RUSLE2 ID's record.						
Input Units Code— Code identifying whether input is in English or	"Input_Units_C			I1	2	59
SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code	ode"					
in the AnnAGNPS ID data section)						
Go to Layout Matrix		Cot	Table of Conten	ts		

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

HYDRAULIC GEOMETRY DATA (BUILT-IN)

There are several built-in reach geometry coefficient sets. The built in sets are:

Hydroulia			Cha	nnel		Vallav	Width	
Geometry ID	Representing	Widt	h	Depth	ı	vancy wiuth		
Geometry ID		Coeff. (m)	Exp.	Coeff. (m)	Exp.	Coeff. (m)	Exp.	
Curve A	Mediterranean climate of winter rainfall such as San Francisco region at 30 inches annual precipitation	0.5889	.38	0.5889	.38	0.5889	.38	
Curve B	High-rainfall areas such as Pennsylvania, with annual accumulation precipitation of 45 inches	0.4901	.39	0.4901	.39	0.4901	.39	

AnnAGNPS Input File Specifications

Hadavalla			Cha	annel		Vallay Width		
Hydraulic Coornectory ID	Representing	Widt	h	Dept	h	Valley	Width	
Geometry ID		Coeff. (m)	Exp.	Coeff. (m)	Exp.	Coeff. (m)	Exp.	
Curve C	Mountain areas in the Upper Green River, Wyoming	0.1878	.45	0.1878	.45	0.1878	.45	
Curve D	Mountain areas in the Upper Salmon River, Idaho	0.2546	.39	0.2546	.39	0.2546	.39	
Curve E	Pacific Maritime Mountains (N. Cascades, Puget Lowland, coast range, & Willamette valley)	0.3462	.43	0.3462	.43	0.3462	.43	
Curve F	Western CordilleraKlamath mountains, Cascades, eastern Cascades, Blue mountains, northern Rockies, middle Rockies, and Wasatch & Uinta mountains	0.2777	.42	0.2777	.42	0.2777	.42	
Curve G	Western Interior Basin & RangesColumbia Basin, Snake River Basin/high Desert, northern Basin range, ands Wyoming Basin	0.0586	.51	0.0586	.51	0.0586	.51	
Curve H	Mission Creek, Oregon—eastern slopes of the Umatilla Mountains	0.3008	.378	0.3008	.378	3.1002	.294	
Curve I	Seco Creek, Texasnorth of San Antonio in the Edwards Aquifer	1.4926	.3151	1.4926	.3151	1.4926	.3151	
Curve J	Davis Hollow Basin & Hole Basin, West Virginiatributaries to the Greenbriar River in karst geomorphology	0.4016	.4193	0.4016	.4193	0.4016	.4193	
Curve K	Wrights Brook & Kiff BrookNew York City water supply watersheds in the West Branch of the Delaware River Basin, New York	0.0132	.8033	0.0132	.8033	0.0132	.8033	
Curve L	Cole Gully & Wikoff Bayousubwatersheds within the Bayou Plaquemine Brule in the Mermentau River Basin, southwestern Louisiana	5.9843	.1448	5.9843	.1448	34.7450	.1448	
Curve M	Carneros Creek in Monterey & San Benito Counties, central coast, California	1.1985	.7338	1.1985	.7338	2.5476	.5141	
1	No layout matrix available	_	Go to	Table of C	Conten	ts		

Revision: December 4, 2019 File Name: Input_Specifications_v5.51_2019.xx.xx.docx

All built-in sets use the same Geometry Length Coefficient (79.19) and Exponent (0.60).

HYDRAULIC GEOMETRY DATA (USER-DEFINED IN INPUT)

If sets other than the built-in ones are desired then enter using format below.

Optional unless referenced in Cell Data, Global IDs, Factors, and Flags Data, or Reach Data.

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed				Α	1	1-9
below.						
The following record repeats for the nu	mber of reach	geometry c	oefficient sets.			
Hydraulic Geometry ID—Alphanumeric string identifying a reach	"Hydraulic_Geo			A10	2	1
geometric coefficient and exponent set. Blank defaults to what is set	m_ID"					
in						
Simulation Period section.						

		TT • (-	
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Channel Length Coefficient—Geomorphic length coefficient in equation: length = coef * Da ^{exp} . Where length = distance from hydraulically most distant point in watershed (ft or m) Da = total drainage area (acres or hectares) at the reach outlet Reach length is determined by the difference in two solutions of this relationship. One at the downstream end and the other at the unstream end of the reach	"Channel_Lengt h_Coef"		Blank or {0.00000001 to 99999999.9} [0.00000001 to 99999999.9]	F10	2	2
Channel Length Exponent—Geomorphic length exponent in equation: length = coef * Da ^{exp} . Where length = distance from hydraulically most distant point in watershed (ft or m) Da = total drainage area (acres or hectares) at the reach outlet. Reach length is determined by the difference in two solutions of this relationship. One at the downstream end and the other at the upstream end of the reach	"Channel_Lengt h_Exp"		Blank or {0.00000001 to 99999999.9} [0.00000001 to 99999999.9]	F10	2	3
Channel Width Coefficient—Geomorphic width coefficient in equation: width = coef * Da ^{exp} . Where width = channel bank full width (ft or m) Da = total drainage area (acres or hectares) at the reach outlet.	"Channel_Widt h_Coef"		{0.00000001 to 99999999.9} [0.00000001 to 99999999.9]	F10	2	4
Channel Width Exponent—Geomorphic width exponent in equation: width = coef * Da ^{exp} . Where width = channel bank full width (ft or m) Da = total drainage area (acres or hectares) at the reach	"Channel_Widt h_Exp"		{0.0 to 99999999.9} [0.0 to 99999999.9]	F10	2	5
Channel Depth Coefficient—Geomorphic depth coefficient in equation: depth = coef * Da ^{exp} . Where depth = channel bank full depth (ft or m) Da = total drainage area (acres or hectares) at the reach outlet	"Channel_Dept h_Coef"		{0.00000001 to 99999999.9} [0.00000001 to 99999999.9]	F10	2	6
Channel Depth Exponent—Geomorphic depth exponent in equation: depth = coef * Da ^{exp} . Where depth = channel bank full depth (ft or m) Da = total drainage area (acres or hectares) at the reach outlet.	"Channel_Dept h_Exp"		{0.0 to 999999999.9} [0.0 to 999999999.9]	F10	2	7
Valley Width Coefficient—Geomorphic valley width coefficient in equation: width = coef * Da ^{exp} . Where valley width = valley width (ft or m) Da = total drainage area (acres or hectares) at the reach outlet. Blank defaults to Channel Width Coefficient; note—Valley Width Exponent field must also be blank.	"Valley_Width_ Coef"		Blank, or {0.00000001 to 999999999.9} [0.00000001 to 999999999.9]	F10	2	8

Revision: December 4, 2019 Fi	File Name: Input_Specifications_v5.51_2019.xx	k.xx.docx
-------------------------------	---	-----------

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Valley Width Exponent—Geomorphic valley width exponent in equation: width = coef * Da ^{exp} . Where valley width = valley width (ft) Da = total drainage area (acres or hectares) at the reach outlet. Blank defaults to Channel Width Exponent; note—Valley Width Coefficient field must also be blank.	"Valley_Width_ Exp"		Blank, or {0.0 to 999999999.9} [0.0 to 999999999.9]	F10	2	9
Go to Layout Matrix		Go to	Table of Conten	ts		

IMPOUNDMENT DATA

Optional

Opt	Ionai					
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-12
The following record repeats fo	r the number	of impound	nents.	-	_	_
Impoundment ID—Alphanumeric string identifying the impoundment.	"Impoundment_ ID"			A10	2	1
Impoundment Infiltration —Infiltration rate for bottom the impoundment. Blank will default to 0.	"Infiltration"	{in / hr} [mm / hr]	Blank, or {0.0 to 19.} [0.0 to 500.0]	F10	2	2
Impoundment Seepage —Constant value seepage rate through the embankment. Blank will default to 0.	"Seepage"	${cfs} {m^{3}/sec}$	Blank, or {0.0 to 105.} [0.0 to 3.0]	F10	2	3
Permanent Pool Depth —Depth used as the base for impoundment temporary storage and spillway discharge. If zero or blank, there is no permanent pool.	"Permanent_Po ol_Depth"	{ft} [m]	Blank, or {0.0 to 328.0} [0.0 to 100.0]	F10	2	4
Impound Volume Coefficient—Coefficient in power curve describing the volume- depth relationship: Vol = coef (Depth) ^{exp} where: Vol = Storage volume above channel bottom elevation. (acre- feet or hectare-meter). Depth = Vertical distance above channel bottom (feet or meters)	"Volume_Coef"		[>0.0 to 260.] ⁶	F10	2	5
Impound Volume Exponent—Exponent in power curve describing the volume-depth relationship: Vol = coef (Depth) ^{exp} where: Vol = Storage volume above channel bottom elevation. (acre- feet or hectare-meter). Depth = Vertical distance above channel bottom (feet or meters) Blank defaults to 1.	"Volume_Exp"		Blank, or 0. to 10.	F10	2	6
Impound Discharge Coefficient—Coefficient in power curvedescribing the discharge- depth relationship: $Q = coef (Depth)^{exp}$ where: $Q = Principal spillway discharge (cfs or m³/sec).$ Depth = Vertical distance above channel bottom (for pressure flow) or above permanent pool (for weir flow) (feet or meters).	"Discharge_Coe f"		[>0.0 to 2000.0] ⁷	F10	2	7

⁶ Unit conversion from English to SI is non-linear. Appropriate English ranges are: {>0.0 to 642.} for minimum Impound Volume Exponent; and {>0.0 to 59.7} for maximum Impound Volume Exponent.

⁷ Unit conversion from English to SI is non-linear. Appropriate English ranges are: {>0.0 to 39000.} for minimum Impound Discharge Exponent; and {>0.0 to 3620.} for maximum Impound Discharge Exponent.

	-	-	
Revision: December 4, 2019	File Name:	Input_Specifications_	_v5.51_2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Impound Discharge Exponent—Exponent in power curve describing the discharge- depth relationship:	"Discharge_Exp "		Blank, 0.5 , 1.5, or 2.5	F10	2	8
$Q = coef (Depth)^{exp}$						
where: Q = Principal spillway discharge (cfs or m ³ /sec). Depth = Vertical distance above channel bottom (for pressure flow) or above permanent pool (for weir flow) (feet or meters).						
Use 0.5 for pressure flow and 1.5 for horizontal weir flow and 2.5 for v-notch weir flow. Blank defaults to 0.5						
Sediment Clean Out Depth—- Depth of sediment accumulation	"Sed_Clean_Ou	(ft.)	Blank, or	F10	2	9
before clean out.	t_Deptn	m	0 to permenant			
A blank defaults to the permanent pool depth.			pool depth			
Sediment Clean Out Year—- Number of years of sediment accum-	"Sed_Clean_Ou		Blank, or	I10	2	10
ulation before clean out. A blank in, "Permanent Pool Depth" and	t_real		>=0			
"Sediment Clean Out Depth" fields and this field will default to 5						
years. Conversely, a nonblank in either "Permanent Pool Depth"						
or "Sediment Clean Out Depth" fields and a blank in this field will						
default to no clean out.	(75) 1 154					
Reach ID —Alphanumeric string identifying the reach in which the	"Reach_ID"			A10	2	11
impoundment is located. Must be the same as a reach ID in the						
REACH DATA section.						
Input Units Code— Code identifying whether input is in English or	"Input_Units_C			I1	2	12
SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code	oue					
in the AnnAGNPS ID data section)						
Go to Layout Matrix		Go to	o Table of Conten	ts		

IRRIGATION APPLICATION DATA

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-18
The following record repeats for the	e number of Ir	rigation Ap	plications.			
Irrigation Application ID —Alphanumeric string identifying the irrigation schedule. Each application method may be automatic, single-manual, or interval-manual irrigation. The automatic method must have a: (1) begin irrigation season which is the "Management Schedule Event Date" field within the MANAGEMENT SCHEDULE DATA when also referring there to this "Irrigation Application ID"; (2) end irrigation season date which is the "Irrigation Depletion Limit" field within this Irrigation Application ID; and (3) the "Irrigation Depletion Limit" field within this Irrigation Application ID. The single-manual irrigation method must leave theis "Irrigation End Date" field totally blank while completing .the other necessary fields. The interval-manual irrigation method must have an end of irrigation season which is the "Irrigation End Date" field within the first and the method must have an end of irrigation season which is the "Irrigation End Date" field while also completing the appropriate other fields.	"Application_ID "			A10	2	1
Season End Date—Month, day and relative operation (rotation) year that the irrigation season ends. Entry requires that one (and only one) of either Irrigation Depletion Limit, Interval Number, or Interval Days be entered. Leave blank for single-manual application. Blank year defaults to 1. Note–each automatic irrigation season (rotation year) must have its own end of season date or automatic irrigation will continue through the winter months until the next entered rotaion year's end date.	"Season_End_M onth", "Season_End_D ay", "Season_End_Y ear"	mm dd yyyy	1 to 12 1 to 31 1 to 1000	12, 12, 14	2	2-4

Optional unless referenced in Management Schedule Data

	Theld	TI				
Description	Fleid Hoodon fon	Units (English)	Domain	For	Record	Field
Description	Decord #1	{English}	{English} [SI]	mat	No.	No.
	Record #1	[51]				
Method Code—Acceptable values are::	"Method_Code"		1-13	I10	2	5
1 = Furrow—open ends, alternating						
2 = Furrow—open ends, adjacent						
3 = Furrow—blocked, alternating						
4 = Furrow—blocked, adjacent						
5 = Surge						
6 = Border (level or graded)						
7 = Level Basin						
8 = Center Pivot (inc LEPA)						
9 = Linear Move						
10 = Sprinkler (set Move or Solid Set)						
11 = Big Gun (Moving or Solid Set)						
12 = Trickle (Drip, Micro Spray etc)						
13 = Sub-surface						
Cycle Duration—Duration of irrigation application cycle. This	"Cycle_Duratio	hr	Blank or	F10	2	6
field is ignored for automatic applications. Blank defaults to 24	n"		0.1 to 24.0			
hours for manual applications.						
Amount Lost—Percentage of applied irrigation water that runs off.	"Amount_Lost"	%	Blank or	F10	2	7
Only applies to method codes 1, 2, and 5. Blank defaults to 0.0			0.0 to 100.0		_	,
Application Rate—Irrigation application rate expressed as inches	"Application R	$\{in/24-hr\}$	Blank or	F10	2	8
[English] or mm [SI] in a 24-hour period Blank defaults to 1 inch	ate"	[mm/24]	$\begin{cases} 0.0 \text{ to } 36.0 \end{cases}$	110	2	0
[English] of 1111 [51] in a 24-hour period. Drank defaults to 1 men [Fnglish] or 25.4 mm [SI] ner 24-hr		hr]	[0.0 to 914 4]			
Tailwatar Racovary Effectiveness of tailwater recovery: blank	"Tailwater Rec		0.0 to 100.0	F10	2	0
defaults to 100. Only applies to method codes 1. 2. and 5	overy"	70	0.0 10 100.0	110	2	,
Depletion Lower Limit Soil moisture depletion lower limit for	"Depletion Low	(nd)	Blank or	F10	2	10
automatic irrigation scheduling to begin. The depletion lower limit	er_Limit"	(iiu)	0.0 to 1.0	110	2	10
is defined as the fraction of total available soil moisture in the soil			0.0 10 1.0			
column where irrigation will begin. This is equivalent to the soil						
column's sail maisture where irrigation will begin divided by the						
columni s son moisture where inigation will begin divided by the						
moisture or water consolity in the soil column is defined to be the						
amount of water expansion between field conacity and the wilting						
noint. Only used with automatic irrigation and requires irrigation						
season's Irrigation End Date entry. Blank defaults to a manual						
irrigation application (can be either single or interval)						
inigation application (can be entier single of interval).		(;)	D1 1	F10	-	11
Application Amount—Amount of water to be applied during an irrigation	"Application A	{in}	Blank	F10	2	11
cycle. Blank defaults to calculation between rate & duration.	mount"	[mm]	$\{0. \text{ to } 4.0\}$			
	<i>"</i>		[0. to 101.6]			
Area Fraction-Fraction of cell (field) area which has irrigation	"Area_Fraction		Blank or	F10	2	12
water applied. Blank defaults to 1.0			0.0 to 1.0			
Interval Number—Number of fixed interval irrigations during the	"Interval_Numb		Blank or	I10	2	13
irrigation period. Only used with fixed interval irrigation where	er		1 to 100			
Interval is not specified. Requires Irrigation End Date entry. Leave						
blank for automatic irrigation or single-manual irrigation						
application.						
Interval Days—Fixed number of days between irrigations. Only	"Interval_Days"	Days	Blank or	I10	2	14
used with manually-scheduled irrigation applications at fixed			1 to 100			
intervals where the Interval Number is not specified. Requires an						
Irrigation End Date entry. Leave blank for automatic irrigation or a						
single manually-scheduled irrigation application.						
Chemical Multiple—Multiple of manual irrigation applications	"Chemical_Mult		Blank or	I10	2	15
between irrigation applications with chemicals added to the	ipie		1 to 100			
irrigation water. Requires entry of Interval Number or Interval Days.						
Leave blank for automatic irrigation or single-manual application						
irrigation.						

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Sediment Rate—Sediment yield rate (including all particle sizes) at end of field. (Blank defaults to 0.0)	"Sediment_Rate "	{tons / acre / in of irrigation water yield} [metric tons / hectare / mm of irrigation water yield]	Blank or {0.0 to 6.0} [0.0 to 0.53]	F10	2	16
Depletion Upper Limit —Soil moisture depletion upper limit for automatic irrigation scheduling to stop. This depletion upper limit is defined as the fraction of total available soil moisture in the soil column at the irrigation's highest controllable level where irrigation will stop. This is equivalent to the soil column's soil moisture where irrigation will stop divided by the soil column's total available soil moisture. The total available soil moisture or water capacity in the soil column is defined to be the amount of water available between field capacity and the wilting point. Blank defaults to 1.0 (at field capacity).	"Depletion_Upp er_Limit"	(nd)	Blank or 0.0 to 1.0	F10	2	17
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	18
<u>Go to Layout Matrix</u>		<u>Go ta</u>	Table of Conten	ts		

MANAGEMENT FIELD DATA

Required

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-11
The following record repea	ts for the num	ber of fields	5.			
Field ID—Alphanumeric string identifying the field.	"Field_ID"			A10	2	1
Field Landuse Type ID—Alphanumeric string describing the landuse type. Acceptable values are: "Cropland", "Pasture", "Rangeland", "Forest", "Urban".	"Landuse_Type _ID"			A10	2	2
Management Schedule ID —Alphanumeric string identifying the management schedule for the field. Must be the same as a management schedule ID (in Management Schedule Data).	"Mgmt_Schd_I D"			A10	2	3
Gregorian Year for a 1st Year of Rotation —A Gregorian calendar year for a 1 st year of rotation in the designated Management Schedule. (Example: For a Management Schedule involving a 3 year rotation, assigning a Gregorian year of 2000 for a 1 st year of rotation would result in the first rotation year in the management schedule to occur in Gregorian calendar years 2000, 2003, 2006, etc. and in 1997, 1994, 1991, etc.) The date when a management schedule event would occur during the simulation period would be based on this association. Note that there are many such years that can be used which will all results in the same rotation schedule with respect to the Gregorian calendar. Blank defaults to Gregorian calendar year 1 C.E./A.D.	"Greg_Yr_for_1 st _Yr_of_Rotatio n"		Blank, or -100 to 9999	I10	2	4
Percent Rock Cover—Percent surface area covered by rocks. Blank defaults to 0.0.	"Percent_Rock_ Cover"		Blank, or 0.0 to 100.0	F10	2	5

AnnAGNPS Input File Specifications

		-			
Revision: December 4, 2019	File Name:	Input S	Specifications	v5.51	2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Inter-rill Erosion code —Beta code indicating the ratio of rill to inter-rill erosion. Acceptable values are: 1 = rill/inter-rill erosion equal for bare soil (ratio = 0.035) 2 = interrill erosion dominant for bare soil (ratio = 0.025) 3 = rill erosion dominant for bare soil (ratio = 0.050) 4 = coarse soil; low ppt.; cover strongly affects runoff (ratio = 0.045) Blank defaults to 3.	"Interrill_Erosi on_Code"		Blank or 1—4	110	2	6
Random Roughness —Long term random roughness. Surface random roughness resulting from rocks, roots, or any other vegetative effects on surface at the time the field is left undisturbed for greater than number of years it takes for the soil to fully consolidate. Blank defaults to 1.25 in (32 mm).	"Random_Roug hness"	{in} [mm]	Blank, or {0.000004 to 19.6} [0.00001 to 500.0]	F10	2	7
Terrace Horizontal Distance —Distance between terraces on the field. Leave blank if there are no terraces on the field.	"Terrace_Horiz ontal_Distance"	{ft} [m]	Blank or {0.000033 to 9842.} [0.00001 to 3000.0]	F10	2	8
Terrace grade .—Grade in terrace to outlet Zero or blank indicates a flat bottom detention terrace. Must be blank if Terrace Horizontal Distance is blank.	"Terrace_Grade "	len-vert / len-horz (nd)	Blank or 0.0 to 10.0	F10	2	9
Tile Drain ID .—Alphanumeric string identifying the tile drainage applied to the field. Leave blank if no tile drains present	"Tile_Drain_ID "			A10	2	10
Input Units Code — Code identifying whether input is in English or SI units. 0 = English ,1 = SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	11
<u>Go to Layout Matrix</u>		Go to	Table of Conten	ts		

MANAGEMENT OPERATION DATA

\sim			
()a	ntı	Or	۱n
C J	DU	UI.	a
-			

Description Field Header— Required unique field header for each field listed	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat A	Record No.	Field No. 1-19
The following record repeats for the	e number of o	perations re	ferences.	l		
Management Operation ID —Alphanumeric string identifying the management operation.	"Mgmt_Operati on_ID"			A20	2	1
Effect Codes—Operation effects codes describing field changes. Enter up to 5 codes. Allowable codes are: 1 = no effect 2 = soil surface disturbed 3 = current crop residue added to surface 4 = other residue added to the field 5 = current residue removed from field 6 = current crop harvested 7 = plant crop (growth begins) 8 = current crop killed 9 = call in a new crop growth set 10 = current and previous residue removed from field. Only # 1 can be repeated in the array. Blanks default to #1.	"Effect_Code_0 1", Effect_Code_02, Effect_Code_03, Effect_Code_04, Effect_Code_05		1 to 10	12, 12, 12, 12, 12, 12	2	2-6

AnnAGNPS Input File Specifications 2019 File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

	Field	Unite				
Description	Header for	(Fnglish)	Domain	For	Record	Field
Description	Record #1	(English)	{English} [SI]	mat	No.	No.
Residue Cover Remaining—Percent residue cover remaining on	"Residue Cover	%	Blank or	F10	2	7
the surface after a soil surface disturbing operation. Only used if an	_Remaining"	/0	>0. to 100.0	110	4	,
oneration effect is 2 and % of residue weight remaining is not						
available. If the effect code is 2, either "Residue Cover Remaining"						
or "Residue Weight Remaining" must be >0 while the other is 0.0 or						
blank. Blank defaults to 0.0. Must be blank or 0.0 for other than						
effect code 2.						
Residue Weight Remaining—Percent residue weight remaining on	"Residue_Weigh	%	Blank or	F10	2	8
the surface after a soil surface disturbing operation. Only used if an	t_Remaining"		>0. to 100.0			
Effect code is 2 and Residue Cover Remaining is not available. If						
the effect code is 2, either "Residue Cover Remaining" or "Residue						
Weight Remaining" must be >0 while the other is 0.0 or blank.						
Blank defaults to 0.0. Must be blank or 0.0 for other than effect code						
2.						
Area Disturbed—Percent surface area disturbed by operation.	"Area_Disturbe	%	Blank or	F10	2	9
Only used if Effect code is 2. Effect code of 2 requires a value >0.	u		>0. to 100.0			
Must be blank or 0.0 for other than effect code 2.						
Initial Random Roughness—Initial random roughness. Only used	"Initial_Rando m_Roughness"	{in}	Blank or	F10	2	10
if Effect code is 2. Effect code of 2 requires a value >0. Must be	m_Roughness	[mm]	$\{>0. \text{ to } 10.0\}$			
blank for other than effect code 2.			[>0. to 254.0]			
Final Random Roughness—Final consolidated random roughness.	"Final_Random Boughness"	{in}	Blank or	F10	2	11
Only used if Effect code is 2. Effect code of 2 requires a value >0.	_Koughness	[mm]	{>0. to 10.0}			
Must be blank for other than effect code 2.			[>0. to 254.0]			
Operation Tillage Depth —Depth of tillage operation. Only used if	"Operation_Till	{in}	Blank or	F10	2	12
Effect Code is 2. Effect code of 2 requires a value >0. If any Effect	age_Depth	[mm]	{>. to 100.}			
Code is 2, then a blank defaults to 200 [mm]; othetrwise the default			[>0.0 to 2540.}			
IS ZETO	"A JJ-J CC	0 (D1 1	F10	-	10
Added Surface Residue—Additional residue applied that remains	"Added_Surface Residue"	%	Blank or	F10	2	13
on the surface. Only used if Effect code is 4. Blank defaults to 0.0			0. to 100.0			
meaning no added residue is applied. Must be blank for other than						
effect code 4.	"Surface Deco		Dlamlr on	E10	2	14
surface Decomposition—Surface decomposition coefficient for	mp"		Dialik of 0 ± 10.0	F10	Z	14
0.016 Must be blank	-		0.1010.0			
for other than effect code A						
Sub-surface Decomposition Sub-surface decomposition	"Subsurface De		Blank or	F10	2	15
coefficient for added residue. Only used if Effect code is 4 Blank or	comp"		0 to 10.0	110	2	15
0.0 defaults to 0.016 Must be			0.10 10.0			
blank for other than effect code 4.						
Surface Residue Added surface residue amounts at three cover	"Surface Resid	/lb / acre}	(>0 to 99924)	3E10	2	16-18
percentages. The order and value of the cover percentages are: 30%	ue_30%",		$\{>0, 10, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0$	51 10	2	10-10
60% and $90%$ Effect code of 4 requires a value >0 If effect code =	Surface_Residue	hectare]	[> 0. 10 112000.0]			
4 and all values are blank then the surface residue at 30% defaults	_00%, Surface Residue	neeunej				
to 0.01. This parameter is only used for non-crop in the RUSLE1	_90%					
version within AnnAGNPS.						
Must be blank for other than effect code 4.						
Input Units Code— Code identifying whether input is in English or	"Input_Units_C ode"			I1	2	19
SI units. $0 = \text{English}, 1 = \text{SI}$ (Blank defaults to the input units code	out					
in the AnnAGNPS ID data section)						
Go to Layout Matrix		Go to	Table of Conten	ts		

MANAGEMENT SCHEDULE DATA

Required

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-18

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
The following record repeats for the num All events within a grouping	ber of events w y must be seau	vithin a sche ential in tim	duled grouping.			
Management Schedule ID—Alphanumeric string identifying the management schedule. Must be present for the start of a new schedule. Use only once per schedule; leave blank for subsequent events within a schedule	"Mgmt_Schd_I D"			A10	2	1
Event Date—Complete date (month, day, & relative year within a scheduled grouping) for the event. Blank defaults to 1 for the event year. NOTE: for management schedule purposes, leap year is not considered when determining the Julian day, Gregorian day, and Gregorian date for the scheduled events. This is because there can be many management field IDs that use the same management schedule but with different "Gregorian Year for 1st Year of Rotation".	"Event_Month", "Event_Day", "Event_Year"	mm dd yyyy	Blank, or 1 to 12 1 to 31 1 to 1000	I2, I2, I4	2	2-4
Event Contour ID —Alphanumeric string identifying contour data. Must be the same as a contour ID (in Contour Data). Only needed for the first event involving contours or when a change occurs.	"Contour_ID"			A10	2	5
Event New Crop ID —Alphanumeric string identifying new crop information for the event . Must be the same as a crop ID (in Crop Data). Required for one event in a cropland event set with additional entries if a change occurs.	"New_Crop_ID "			A20	2	6
Event Strip Crop ID —Alphanumeric string identifying strip crop data for the event . Must be the same as a strip crop ID (in Strip Crop Data). Only needed for the first event involving strip crops in a management schedule or when a change occurs.	"Strip_Crop_ID "			A10	2	7
Event New Non-crop ID —Alphanumeric string identifying new non-crop landuse data for the event. Must be the same as a non-crop ID (in Non-Crop Data). Required for one event of a non-crop management schedule .	"New_Non- Crop_ID"			A20	2	8
Curve Number ID—A runoff curve number ID that must match a corresponding ID in the RUNOFF CURVE NUMBER DATA section and is scheduled to become effective on or shortly after the scheduled date. A curve number ID may be entered for the first scheduled date for each Management Schedule ID and when a change in runoff curve	"Curve_Numbe r_ID"			A40	2	9
number is to occur. . This field may be left blank. If blank, AnnAGNPS checks for the presence of a user-entered "Default Crop RCN ID" or "Default Non- Crop RCN ID" in the GLOBAL IDS, FACTORS, AND FLAGS DATA section based on the "Event New Crop ID" or "Event New Non-crop ID" of this section. If a default RCN ID is present, AnnAGNPS will use that RCN ID. If a default RCN ID is not present (blank), AnnAGNPS will use an internally defined RCN ID as described in the GLOBAL IDS, FACTORS, AND FLAGS DATA section.						
When a scheduled Curve Number ID is specified alone without a New Crop ID also being specified, the curve number becomes effective that day. When a Scheduled Curve Number ID is specified along with a New Crop ID and a Management Operation ID that includes a planting (7) effect code also being specified, the curve number transitions from the old to the new curve number as a function of the days from planting to harvest. The curve number represents a mature crop when a planting management operation ID is specified.						
Post Event Manning's n —Manning's n value to use after operation occurs. Required for first operation in a management schedule and when a change occurs Defaults to cell's current value.	"Post_Event_M annings_n"		blank, 0.005 to 1.0	F10	2	10

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Post Event Surface Constant —Surface condition constant to use after operation occurs. Required for first operation in a management schedule and when a change occurs Defaults to cell's current value.	"Post_Event_Su rface_Constant"		blank, 0.0 to 1.0	F10	2	11
 Operation Residue Change—Residue amount added or subtracted for the operation. Amount is always a positive number. Data interpretation depends on the effect codes (in Operation Reference Data) associated with the operation. Blank defaults to 0.0. Actions by effect codes are: 3, (optional). If 0, the current crop potential residue is added otherwise this amount is added to the current crop residue. 4, (required) added as a unique residue. 5, (optional) If 0, then all of the current crop surface residue will be removed otherwise only this amount is subtracted. 10, (optional) If 0, then all surface residues are removed, otherwise this value is divided by the total of all surface residues to determine a fraction to remove from each residue. 	"Operation_Res idue_Change"	{lb/acre} [kg/hectare]	Blank or {0.0 to 99924.} [0.0 to 112000.0]	F10	2	12
Event Fertilizer Application ID —Alphanumeric string identifying the fertilizer information for the operation. Must be the same as a fertilizer application ID (in Fertilizer Application Data). Blank indicates no fertilizer applied with event.	"Fertilizer_Appl ication_ID"			A10	2	13
 Event Irrigation Application ID—Alphanumeric string identifying the irrigation application record to be used for this event. Must be the same as an irrigation application ID in the Irrigation Application Data section. Note: Irrigation will begin on the relative year event date specified in this management schedule and will end based on the relative year "Season End Date" in the corresponding "Irrigation Application Data" record. If the irrigation season end date is in a relative year prior to the irrigation begin date in this management schedule then irrigation will continue until the relative year rotation has occurred and the irrigation application is designated then the season end date should be left blank in the corresponding irrigation application record. Blank indicates no irrigation applied with event. 	"Irrigation_App lication_ID"			A10	2	14
Management Operation ID—Alphanumeric string identifying the operation information for the operation. Must be the same as an operation ID (in Management Operation Data). Blank indicates a "NO OPERATION". Note–if a "NO OPERATION" ID is not found in the MANAGEMENT OPERATION DATA section, AnnAGNPS inserts one with all effect codes equal to 1.	"Mgmt_Operati on_ID"			A20	2	15
Tile Drain Controlled Status —Alphanumeric string indicating a change in controlled tile drain status. "Open" or "close" are acceptable inputs. The initial tile drain status is "Open" when there is a tile drain. Blank defaults to the previous controlled status. This field is ignored if a tile drain has not been assigned to the cell when this specific schedule is active.	"Tile_Drain_Co ntrolled_Status"		Blank, or "Open" or "Close"	A10	2	16
Tile Drain Controlled Depth —Numeric value indicating a change in the controlled tile drain depth. Entry of a non-zero value indicates a change in the tile drain depth via controlling the drain outlet. Entering a value of '0.0' in this field is the same as entering "close" in the Tile Drain Controlled Status field. Entering a value equal to or greater than the tile drain invert depth returns the control to the invert (e.g., a value of 2000. [mm]). This field is ignored if a tile drain has not been assigned to the active cell	"Tile_Drain_Co ntrolled_Depth"	{in} [mm]	Blank, or {0. to 78.74} [0. to 2000].	F10	2	17

 AnnAGNPS Input File Specifications

 2019
 File Name: Input_Specifications_v5.51_2019.xx.xx.docx
 Revision: December 4, 2019

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	18
Event Pesticide Application ID —Alphanumeric string identifying the pesticide application information for operation. Must be the same as a pesticide application ID (in Pesticide Application Data). Leave fields blank if no pesticides are applied in the operation. The maximum allowable number of pesticide ids for any event is 5.	"Pest_App_ID_ 1", "Pest_App_ID_ 2", "Pest_App_ID_ 3", "Pest_App_ID_ 4", "Pest_App_ID_ 4", "Pest_App_ID_ 5",			A10	2	19-23
Go to Layout Matrix	Go to Table of Contents					

NON-CROP DATA

Optional unless referenced in Management Schedule Data

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				A	1	1-14
The following record repeats	beats for the number of Non-Crops.					
Non-Crop ID —Alphanumeric string identifying a non-cropland landuse.	"Non-Crop_ID"			A20	2	1
Non-Crop Description—Description of the non-crop landuse. (For user reference only. Not used within AnnAGNPS.)	"Non- Crop_Descriptio n"			A60	2	2
Annual Root Mass—Average annual live root mass in the top 4 in (100 mm) of soil. This parameter is required if the 'USLE C-Factor' below is blank. If the 'USLE C-Factor' parameter below is populated then this parameter is not used.	"Annual_Root_ Mass"	{lb / acre} [kg / hectare]	Blank or {0.0 to 99924.} [0.0 to 112000.0	F10	2	3
Annual Cover Ratio—Average annual ratio of ground covered by canopy cover to total ground area. This parameter is required if the 'USLE C-Factor' below is blank -or- if the FAO ET Enhancement feature is enabled (true - default). May be left blank if the 'USLE C- Factor' parameter below is populated -and- the FAO ET Enhancement feature is disabled (set to false) in the GLOBAL IDS, FACTORS, AND FLAGS DATA section.	"Annual_Cover _Ratio"		Blank or 0.0 to 1.0	F10	2	4
Annual Rain Fall Height—Average annual distance rainfall falls after being intercepted by the crop canopy. This parameter is required if the 'USLE C-Factor' below is blank -or- if the FAO ET Enhancement feature is enabled (true - default). May be left blank if the 'USLE C-Factor' parameter below is populated -and- the FAO ET Enhancement feature is disabled (set to false) in the GLOBAL IDS, FACTORS, AND FLAGS DATA section.	"Annual_Rain_ Fall_Height"	{ft} [m]	Blank or {0.0 to 262.0} [0.0 80.0]	F10	2	5
Surface Residue Cover—Percent surface residue cover. Plant basal area is not considered as being part of the ground cover. Blank = 0.0%	"Surface_Cover _Residue"	%	Blank or 0 to 100.0	F10	2	6
USLE C-Factor—optional, constant USLE C-factor to be used in lieu of computed RUSLE time-varying value. Blank field defaults to computed RUSLE time-varying value.	"USLE_C-Fctr"		Blank or 0.0 to 2.0	F10	2	7
Basal Crop Coefficient ("Kcb-mid") —Average annual coefficient at the mature growth stage; used in adjusting the potential evapotranspiration (ET). The coefficient for all days in the growing season will be equal to this value unless internally adjusted based on climate conditions. The coefficient is internally set to 0.0 for all days outside of the growing season. Blank defaults to 1.0.	"Basal_Crop_C oef_Mid"	[nd]	Blank or 0.0 to 1.2	F10	2	8

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Growing Season Start Month —Month when the growing season begins. Blank = 01	"Growing_Seaso n_Start_Month"	[month]	Blank or 1 to 12	I5	2	9
Growing Season Start Day —Day when the growing season begins. Blank = 01	"Growing_Seaso n_Start_Day"	[day]	Blank or 1 to 31	15	2	10
Growing Season End Month —Month when the growing season ends. Ending date cannot be chronologically earlier than the beginning date. Blank = 12	"Growing_Seaso n_End_Month"	[month]	Blank or 1 to 12	15	2	11
Growing Season End Day—Day when the growing season ends. Ending date cannot be chronologically earlier than the beginning date. Blank = 31	"Growing_Seaso n_End_Day"	[day]	Blank or 1 to 31	15	2	12
Basal Crop Coefficient Climatic Adjustment code — Code indicating whether the basal crop coefficient "Kcb-mid" will be adjusted based on climate conditions. Acceptable values are: $Y =$ Adjustment; N = No adjustment. Blank will default to 'Y' unless the global basal crop climate adjustment code is set to false in the Global IDs, Factors, and Flags Data section.	"Basal_Crop_C oef_Climate_Ad just"		Blank or Y or N	A1	2	13
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	14
Go to Layout Matrix		<u>Go to</u>	Table of Conten	ts	_	

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

OUTPUT OPTIONS DATA – GLOBAL

See Attachment A for a detailed explanation

\sim			1	
()1	nt1/	٦n	ดเ	

Description Field Header— Required unique field header for each field listed below.	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No. 1-40
The following reco	rd does not repeat.	·				·	
Global request for all version 3 comma separated variable format database files (*.csv); default is false.	"Glbl_All_V3_csv"			blank, T, or F	A1	2	1
Global request for all version 3 data preparation verification files (*.dpp); default is false.	"Glbl_All_V3_dpp"			blank, T, or F	A1	2	2
Global request for all version 3 input data verification files (*.npt); default is false.	"Glbl_All_V3_npt"			blank, T, or F	A1	2	3
Global request for all version 3 simulation verification files (*.sim); default is false.	"Glbl_All_V3_sim"			blank, T, or F	A1	2	4
Global request for all version 3 formatted individual table files (*.txt); default is false.	"Glbl_All_V3_txt"			blank, T, or F	A1	2	5
Global request for the program execution log file (AnnAGNPS.log); default is true.	"Log_to_File"			blank, T, or F	A1	2	6
Global request for the program execution log to the screen; default is true.	"Log_to_Screen"			blank, T, or F	A1	2	7
Global request for the warning file; default is true.	"Warning_File"			blank, T, or F	A1	2	8
Global request for the version 1 & 2 formated table output files; default is false.	"V1/2_Output_Files"			blank, T, or F	A1	2	9
Reserved.	"Reserved"			blank only	_	2	10
Global request for all cells to be included in the output; default is true.	"Glbl_All_Cells"			blank, T, or F	A1	2	11
Global request for all feedlots to be included in the output; default is true.	"Glbl_All_Feedlots"			blank, T, or F	A1	2	12

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

Description	Field	Code		Density	For	Record	Field
Description	Header for Becord #1	Index	Col.	Domain	mat	No.	No.
Global request for all field nonds to be included in the output: default is	"Glb1 All Fld Ponds"			blank T or			
true.	Gibi_/in_riu_ronus			F	A1	2	13
	"Glbl_All_Gullies"			blank, T, or		-	1.4
Global request for all gullies to be included in the output; default is true.				F	Al	2	14
Global request for all point sources to be included in the output; default is	"Glbl_All_Pt_Srcs"			blank, T, or	Δ1	2	15
true.				F			10
Global request for all reaches to be included in the output; default is	"GIDI_AII_Reaches"			blank, T, or	A1	2	16
Global request for all impoundments to be included in the output: default	"Glbl All Impound"			r blank T or			
is true.				F	A1	2	17
Global request for all wetlands to be included in the output; default is	"Glbl_All_Wetlands"			blank, T, or	A 1	2	10
true.				F	AI	Z	10
Global request for all nutrients to be included in the accumulation &	"Glbl_All_AA_Nutr"			blank, T, or	A1	2	19
average annual output; default is true.	"Cibl All AA Dost"			F			-
average appual output: default is true	GIDI_AII_AA_I est			blank, 1, or	A1	2	20
Reserved	"Reserved"			blanks only		2.	21
Reserved.	"Reserved"			blanks only		2	22
Global request for all sediment to be included in the accumulation &	"Glbl_All_AA_Sed"			blank, T, or	. 1		
average annual output; default is true.				F	AI	2	23
Global request for all water to be included in the accumulation & average	"Glbl_All_AA_Wtr"			blank, T, or	Δ1	2	24
annual output; default is true.				F	111	2	24
Global request for all nutrients to be included in the event output; default	"Glbl_All_EV_Nutr"			blank, T, or	A1	2	25
is true.	"Clb1 All EV Post"			F blank T on			
default is true	GIDI_AII_E v_I est			blank, 1, or	A1	2	26
Global request for all sediment to be included in the event output: default	"Glbl All EV_Sed"			blank, T. or			
is true.				F	Al	2	27
Global request for all water to be included in the event output; default is	"Glbl_All_EV_Wtr"			blank, T, or	Δ 1	r	28
true.				F	AI		20
Reserved.	"Reserved"			blanks only		2	29
Reserved.	"Reserved"			blanks only		2	30
Global request for version 2 & 3 output to be in total mass units [water &	"Glbl_All_V2/3_Mass"			blank, T, or	A 1	2	21
or kg (SI): default is false				F	AI	Z	51
Global request for version 2 & 3 output to be in ratio units [total mass	"Glbl All V2/3 Ratio"						
from contributing source divided by total mass at reference reach				blank, T, or	A1	2	32
location]; default is false.				F			
Global request for version 2 & 3 output to be in units of total mass	"Glbl_All_V2/3_UA"						
divided by contributing area (unit area) [water & sediment are in tons/ac				blank, T, or	A1	2	33
(English) or Mg/ha (SI), chemicals are in lbs/ac (English) or kg/ha (SI)];				F		_	
default is false. I $\lfloor lbs/ac \rfloor = 1.120848 \lfloor kg/na \rfloor$	"Reserved"			blank only		2	24
Local request for version 2 CONCEPTS formatted output file	"V2 Concepts"			blank T or		2	54
(AnnAGNPS.cpt): default is false.	·			F	A1	2	35
Reserved.	"Reserved"			blank only		2	36
Local request for version 2 average annual formated output file	"V2_AA"			blank, T, or	Λ1	n	27
(AnnAGNPS_AA.csv); default is true.				F	AI	Z	57
Local request for version 2 event formated output file	"V2_EV"			blank, T, or	A1	2	38
(AnnAGNPS_EV.csv); default is false.	«X/1 A A ??			F			
Local request for version 1 accumulated data output file	•• v 1_AA**			blank, T, or	A1	2	39
(AnnAGNI S.acc), uclauit is faise.	"V1 EV"			r blank T or			
default is false.				F	A1	2	40
Go to Layout Matrix		Go to	Tahle	of Contents		<u> </u>	

OUTPUT OPTIONS DATA – AA

See Attachment A for a detailed explanation Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.		
Field Header— Required unique field header for each field listed below.					Α	1	1-55		
The following record does not repeat.									

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_aa _01= _1000 +					
AnnAGNPS_AA_Feedlots_(mass).txt (Reserved – Not currently produced)	"Reserved"	1.		blank, T, or F	A1	2	1.
AnnAGNPS_AA_Feedlots_(ratio).txt (Reserved – Not currently produced)	"Reserved"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_AA_Feedlots_(unit area).txt (Reserved – Not currently produced)	"Reserved"	3.		blank, T, or F	A1	2	3.
AnnAGNPS_AA_Field_Ponds_(mass).txt (Reserved – Not currently produced)	"Reserved"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_AA_Field_Ponds_(ratio).txt (Reserved – Not currently produced)	"Reserved"	5.		blank, T, or F	A1	2	5.
AnnAGNPS_AA_Field_Ponds_(unit area).txt (Reserved – Not currently produced)	"Reserved"	6.		blank, T, or F	A1	2	6.
AnnAGNPS_AA_Gullies_(erosion).csv	"AA_Gullies_Erosion"	7.		blank, T, or F	A1	2	7.
AnnAGNPS_AA_Gullies_(sediment).txt (Reserved – Not currently produced)	"Reserved"	8.		blank, T, or F	A1	2	8.
AnnAGNPS_AA_Gullies_(nutrients).txt (Reserved – Not currently produced)	"Reserved"	9.		blank, T, or F	A1	2	9.
AnnAGNPS_AA_Nitrogen_load_(mass).csv	"AA_N_Ld_Mass"	10.		blank, T, or F	A1	2	10.
AnnAGNPS_AA_Nitrogen_load_(ratio). csv	"AA_N_Ld_Ratio"	11.		blank, T, or F	A1	2	11.
AnnAGNPS_AA_Nitrogen_load_(unit_area). csv	"AA_N_Ld_UA"	12.		blank, T, or F	A1	2	12.
AnnAGNPS_AA_Nitrogen_yield_(mass). csv	"AA_N_Yld_Mass"	13.		blank, T, or F	A1	2	13.
AnnAGNPS_AA_Nitrogen_yield_(ratio). csv	"AA_N_Yld_Ratio"	14.		blank, T, or F	A1	2	14.
AnnAGNPS_AA_Nitrogen_yield _(unit_area). csv	"AA_N_YId_UA"	15.		blank, T, or F	A1	2	15.
AnnAGNPS_AA_Organic_Carbon_load_(mass). csv	"AA_OC_Ld_Mass"	16.		blank, T, or F	A1	2	16.
AnnAGNPS_AA_Organic_Carbon_load_(ratio). csv	"AA_OC_Ld_Ratio"	17.		blank, T, or F	A1	2	17.
AnnAGNPS_AA_Organic_Carbon_load_(unit_area). csv	"AA_OC_Ld_UA"	18.		blank, T, or F	A1	2	18.
AnnAGNPS_AA_Organic_Carbon_yield_(mass). csv	"AA_OC_Yld_Mass"	19.		blank, T, or F	A1	2	19.
AnnAGNPS_AA_Organic_Carbon_yield_(ratio). csv	"AA_OC_Yld_Ratio"	20.		blank, T, or F	A1	2	20.
AnnAGNPS_AA_Organic_Carbon_yield_(unit_area). csv	"AA_OC_YId_UA"	21.		blank, T, or F	A1	2	21.
AnnAGNPS_AA_Pesticides_load_(mass).txt (Reserved – Not currently produced)	"Reserved"	22.		blank, T, or F	A1	2	22.
AnnAGNPS_AA_Pesticides_load_(ratio).txt (Reserved – Not currently produced)	"Reserved"	23.		blank, T, or F	A1	2	23.
AnnAGNPS_AA_Pesticides_load_(unit_area).txt (Reserved – Not currently produced)	"Reserved"	24.		blank, T, or F	A1	2	24.
AnnAGNPS_AA_Pesticides_yield_(mass).txt (Reserved – Not currently produced)	"Reserved"	25.		blank, T, or F	A1	2	25.
AnnAGNPS_AA_Pesticides_yield_(ratio).txt (Reserved – Not currently produced)	"Reserved"	26.		blank, T, or F	A1	2	26.
AnnAGNPS_AA_Pesticides_yield_(unit_area).txt (Reserved – Not currently produced)	"Reserved"	27.		blank, T, or F	A1	2	27.
AnnAGNPS_AA_Phosphorus_load_(mass).csv	"AA_P_Ld_Mass"	28.		blank, T, or F	A1	2	28.

Description	Field	Code		Density	For	Record	Field
Description	Record #1	Index	C01.	Domain	mat	No.	No.
AnnAGNPS_AA_Phosphorus_load_(ratio). csv	"AA_P_Ld_Ratio"	29.		blank, T, or F	A1	2	29.
AnnAGNPS_AA_Phosphorus_load _(unit_area). csv	"AA_P_Ld_UA"	30.		blank, T, or F	A1	2	30.
AnnAGNPS_AA_Phosphorus_yield_(mass). csv	"AA_P_Yld_Mass"	31.		blank, T, or F	A1	2	31.
AnnAGNPS_AA_Phosphorus_yield_(ratio). csv	"AA_P_Yld_Ratio"	32.		blank, T, or F	A1	2	32.
AnnAGNPS_AA_Phosphorus_yield_(unit_area). csv	"AA_P_YId_UA"	33.		blank, T, or F	A1	2	33.
AnnAGNPS_AA_Point_Sources_(mass).txt (Reserved – Not currently produced)	"Reserved"	34.		blank, T, or F	A1	2	34.
AnnAGNPS_AA_Point_Sources_(ratio).txt (Reserved – Not currently produced)	"Reserved"	35.		blank, T, or F	A1	2	35.
AnnAGNPS_AA_Point_Sources_(unit area).txt (Reserved – Not currently produced)	"Reserved"	36.		blank, T, or F	A1	2	36.
AnnAGNPS_AA_Sediment_Erosion_(mass). csv	"AA_Sed_Eros_Mass"	37.		blank, T, or F	A1	2	37.
AnnAGNPS_AA_Sediment_Erosion_(ratio). csv	"AA_Sed_Eros_Ratio"	38.		blank, T, or F	A1	2	38.
AnnAGNPS_AA_Sediment_Erosion_(unit_area). csv	"AA_Sed_Eros_UA"	39.		blank, T, or F	A1	2	39.
AnnAGNPS_AA_Sediment_Load_(mass). csv	"AA_Sed_Ld_Mass"	40.		blank, T, or	A1	2	40.
AnnAGNPS_AA_Sediment_Load_(ratio). csv	"AA_Sed_Ld_Ratio"	41.		blank, T, or	A1	2	41.
AnnAGNPS_AA_Sediment_Load_(unit_area). csv	"AA_Sed_Ld_UA"	42.		blank, T, or	A1	2	42.
AnnAGNPS_AA_Sediment_Yield_(mass). csv	"AA_Sed_Yld_Mass"	43.		blank, T, or	A1	2	43.
AnnAGNPS_AA_Sediment_Yield_(ratio). csv	"AA_Sed_Yld_Ratio"	44.		blank, T, or	A1	2	44.
AnnAGNPS_AA_Sediment_Yield_(unit_area). csv	"AA_Sed_Yld_UA"	45.		blank, T, or	A1	2	45.
AnnAGNPS_AA_Water_load_(mass). csv	"AA_Wtr_Ld_Mass"	46.		blank, T, or	A1	2	46.
AnnAGNPS_AA_Water_load_(ratio). csv	"AA_Wtr_Ld_Ratio"	47.		blank, T, or	A1	2	47.
AnnAGNPS_AA_Water_load_(unit_area). csv	"AA_Wtr_Ld_UA"	48.		blank, T, or	A1	2	48.
AnnAGNPS_AA_Water_yield_(mass). csv	"AA_Wtr_Yld_Mass"	49.		blank, T, or F	A1	2	49.
AnnAGNPS_AA_Water_yield_(ratio). csv	"AA_Wtr_Yld_Ratio"	50.		blank, T, or F	A1	2	50.
AnnAGNPS_AA_Water_yield_(unit_area). csv	"AA_Wtr_Yld_UA"	51.		blank, T, or F	A1	2	51.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_aa _chld= 1100 +	Par't Field				
NOTE: Separate ranked ratio output files are produced for each reach for which output has been requested and for which one or more cells have contributed to the reach. Separate files are produced for the OUTLET, the upstream end, and the downstream end of the reach.							
AnnAGNPS_AA_Nitrogen_load_UA_RR_Total_reach-ID_reach-location.csv		1.	AA 01 12				
AnnAGNPS_AA_Nitrogen_load_UA_RR_Attached_ <i>reach-ID_reach-location</i> .csv		2.	AA 01 12				
AnnAGNPS_AA_Nitrogen_load_UA_RR_Soluble_reach-ID_reach-location.csv		3.	AA 01				

 AnnAGNPS Input File Specifications

 2019
 File Name: Input_Specifications_v5.51_2019.xx.xx.docx
 Revision: December 4, 2019

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
NOTE: Separate ranked ratio output files are produced for each reach for which output has been requested and for which one or more cells have contributed to the reach. Separate files are produced for the OUTLET, the upstream end, and the downstream end of the reach.							
AnnAGNPS_AA_Organic_Carbon_load_UA_RR_Total_ <i>reach-ID_reach-location</i> .csv		4.	AA 01 18				
AnnAGNPS_AA_Organic_Carbon_load_UA_RR_Attached_ <i>reach-ID_reach-location</i> .csv		5.	AA 01 18				
AnnAGNPS_AA_Organic_Carbon_load_UA_RR_Soluble_reach-ID_reach- location.csv		6.	AA 01 18				
NOTE: Separate ranked ratio output files are produced for each reach for which output has been requested and for which one or more cells have contributed to the reach. Separate files are produced for the OUTLET, the upstream end, and the downstream end of the reach.							
AnnAGNPS_AA_Phosphorus_load_UA_RR_Total_ <i>reach-ID_reach-location</i> .csv		7.	AA 01 30				
AnnAGNPS_AA_Phosphorus_load_UA_RR_Attached_Organic_ <i>reach-ID_reach-location</i> .csv		8.	AA 01 30				
AnnAGNPS_AA_Phosphorus_load_UA_RR_Attached_Inorganic_ <i>reach-ID_reach-location</i> .csv		9.	AA 01 30				
AnnAGNPS_AA_Phosphorus_load_UA_RR_Soluble_Inorganic_ <i>reach-ID_reach-location</i> .csv		10.	AA 01 30				
AnnAGNPS_AA_Sediment_load_by_source_(mass).csv		11.	AA 01 40				
AnnAGNPS_AA_Sediment_load_by_source_(ratio). csv		12.	AA 01 41				
AnnAGNPS_AA_Sediment_load_by_source_(unit-area). csv		13.	AA 01 42				
NOTE: Separate ranked ratio output files are produced for each reach for which output has been requested and for which one or more cells have contributed to the reach. Separate files are produced for the OUTLET, the upstream end, and the downstream end of the reach.							
AnnAGNPS_AA_Sediment_load_UA_RR_Total_ <i>reach-ID_reach-location</i> .csv		14.	AA 01 42				
AnnAGNPS_AA_Sediment_load_UA_RR_Clay_ <i>reach-ID_reach-location</i> .csv		15.	AA 01 42				
AnnAGNPS_AA_Sediment_load_UA_RR_Silt_ <i>reach-ID_reach-location</i> .csv		16.	AA 01 42				
AnnAGNPS_AA_Sediment_load_UA_RR_Sand_ <i>reach-ID_reach-location</i> .csv		17.	AA 01 42				
AnnAGNPS AA Sediment load UA RR Total SnR Gly Pnd reach- ID_reach-location.csv		18.	AA 01 42				
AnnAGNPS_AA_Sediment_load_by_reach_(mass).csv		19.	AA 01 40				
NOTE: Separate ranked ratio output files are produced for each reach for which output has been requested and for which one or more cells have contributed to the reach. Separate files are produced for the OUTLET, the upstream end, and the downstream end of the reach.			01				
AnnAGNPS_AA_Water_load_UA_RR_Total_ <i>reach-ID_reach-location</i> .csv		20.	AA 01 48				
AnnAGNPS AA Water load UA RR Baseflow <i>reach-ID_reach-location.</i> csv (Reserved - Not currently produced)		21.	AA 01 48				
AnnAGNPS_AA_Water_load_UA_RR_Direct_ <i>reach-ID_reach-location</i> .csv (Reserved - Not currently produced)		22.	AA 01 48		-		
<u>Go to Layout Matrix</u>		<u>Go to</u>	Table	of Contents			

OUTPUT OPTIONS DATA – CSV

See Attachment A for a detailed explanation

Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.			
Field Header— Required unique field header for each field listed below.					Α	1	1-61			
The following record does not repeat.										

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_csv _01= 1200 +					
All event loadings from cell to downstream reaches	"All_Evt_Lds_Cell_to_D S Rchs"	1.		blank, T, or F	A1	2	1.
All annual accumulations	"All_AA"	2.		blank, T, or F	A1	2	2.
All events	"All_Events"	3.		blank, T, or F	A1	2	3.
All nitrogen	"All_N"	4.		blank, T, or F	A1	2	4.
All organic carbon	"All_OC"	5.		blank, T, or F	A1	2	5.
All pesticides	"All_Pesticides"	6.		blank, T, or F	A1	2	6.
All phosphorus	"All_P"	7.		blank, T, or F	A1	2	7.
All sediment	"All_Sediment"	8.		blank, T, or F	A1	2	8.
All water	"All_Water"	9.		blank, T, or F	A1	2	9.
Annual accumulation nitrogen load from cell to downstream reaches	"AA_N_Ld_Cel_to_DS_ Rchs"	10.		blank, T, or F	A1	2	10.
Annual accumulation nitrogen load in reaches	"AA_N_Ld_in_Rchs"	11.		blank, T, or F	A1	2	11.
Annual accumulation nitrogen yield from cell to receiving reach	"AA_N_Yld_Cel_to_Rcv Rch"	12.		blank, T, or F	A1	2	12.
Annual accumulation organic carbon load from cell to downstream reaches	- "AA_OC_Ld_Cel_to_DS _Rchs"	13.		blank, T, or F	A1	2	13.
Annual accumulation organic carbon load in reaches	"AA_OC_Ld_in_Rchs"	14.		blank, T, or F	A1	2	14.
Annual accumulation organic carbon yield from cell to receiving reach	"AA_OC_Yld_Cel_to_Rc v_Rch"	15.		blank, T, or F	A1	2	15.
Annual accumulation pesticide load from cell to downstream reaches	"AA_Pest_Ld_Cel_to_DS _Rchs"	16.		blank, T, or F	A1	2	16.
Annual accumulation pesticide load in reaches	"AA_Pest_Ld_in_Rchs"	17.		blank, T, or F	A1	2	17.
Annual accumulation pesticide yield from cell to receiving reach	"AA_Pest_Yld_Cel_to_R cv_Rch"	18.		blank, T, or F	A1	2	18.
Annual accumulation phosphorus load from cell to downstream reaches	"AA_P_Ld_Cel_to_DS_R chs"	19.		blank, T, or F	A1	2	19.
Annual accumulation phosphorus load in reaches	"AA_P_Ld_in_Rchs"	20.		blank, T, or F	A1	2	20.
Annual accumulation phosphorus yield from cell to receiving reach	"AA_P_Yld_Cel_to_Rcv_ Rch"	21.		blank, T, or F	A1	2	21.
Annual accumulation bed & bank erosion within reach	"AA_BB_Eros_in_Rch"	22.		blank, T, or F	A1	2	22.
Annual accumulation bed & bank load in downstream reaches	"AA_BB_Ld_in_DS_Rch s"	23.		blank, T, or F	A1	2	23.
Annual accumulation erosion within cells	"AA_Eros_in_Cels"	24.		blank, T, or F	A1	2	24.
Annual accumulation gully yield from cell to receiving reach	"AA_Gly_Yld_Cel_to_Rc v_Rch"	25.		blank, T, or F	A1	2	25.
Annual accumulation landscape erosion within cells	"AA_LS_Eros_in_Cels"	26.		blank, T, or F	A1	2	26.
Annual accumulation landscape load from cell to downstream reaches	"AA_LS_Ld_Cels_to_DS _Rchs"	27.		blank, T, or F	A1	2	27.
Annual accumulation landscape load in downstream reaches	"AA_LS_Ld_in_DS_Rch s"	28.		blank, T, or F	A1	2	28.

	Field	Code					
Description	Header for Record #1	Index	Col.	Domain	For mat	Record No.	Field No.
Annual accumulation landscape yield from cell to receiving reach	"AA_LS_Yld_Cel_to_Rc v_Rch"	29.		blank, T, or F	A1	2	29.
Annual accumulation & rill erosion within cells	"AA_Rill_Eros_in_Cels"	30.		blank, T, or F	A1	2	30.
Annual accumulation sheet & rill yield from cell to receiving reach	"AA_SR_Yld_Cel_to_Rc v_Rch"	31.		blank, T, or F	A1	2	31.
Annual accumulation water load from cell to downstream reaches	"AA_Wtr_Ld_Cel_to_DS _Rchs"	32.		blank, T, or F	A1	2	32.
Annual accumulation water load in downstream reaches	"AA_Wtr_Ld_in_DS_Rc hs"	33.		blank, T, or F	A1	2	33.
Annual accumulation water yield from cell to receiving reach	"AA_Wtr_Yld_Cel_to_R cv_Rch"	34.		blank, T, or F	A1	2	34.
Nitrogen event loading from cell to downstream reaches	"N_Evt_Ld_Cel_to_DS_ Rchs"	35.		blank, T, or F	A1	2	35.
Nitrogen event loading in reaches	"N_Evt_Ld_in_Rchs"	36.		blank, T, or F	A1	2	36.
Nitrogen event yield from cell to receiving reach	"N_Evt_Yld_Cel_to_Rcv _Rch"	37.		blank, T, or F	A1	2	37.
Organic Carbon event loading from cell to downstream reaches	"OC_Evt_Ld_Cel_to_DS _Rchs"	38.		blank, T, or F	A1	2	38.
Organic Carbon event loading in reaches	"OC_Evt_Ld_in_Rchs"	39.		blank, T, or F	A1	2	39.
Organic Carbon event yield from cell to receiving reach	"OC_Evt_Yld_Cel_to_Rc v_Rch"	40.		blank, T, or F	A1	2	40.
Pesticides event loading from cell to downstream reaches	"Pest_Evt_Ld_Cel_to_DS _Rchs"	41.		blank, T, or F	A1	2	41.
Pesticides event loading in reaches	"Pest_Evt_Ld_in_Rchs"	42.		blank, T, or F	A1	2	42.
Pesticides event yield from cell to receiving reach	"Pest_Evt_Yld_Cel_to_R cv_Rch"	43.		blank, T, or F	A1	2	43.
Phosphorus event loading from cell to downstream reaches	"P_Evt_Ld_Cel_to_DS_ Rchs"	44.		blank, T, or F	A1	2	44.
Phosphorus event loading in reaches	"P_Evt_Ld_in_Rchs"	45.		blank, T, or F	A1	2	45.
Phosphorus event yield from cell to receiving reach	"P_Evt_Yld_Cel_to_Rcv _Rch"	46.		blank, T, or F	A1	2	46.
Sediment event bed & bank erosion within reach	"Sed_Evt_BB_Eros_in_R ch"	47.		blank, T, or F	A1	2	47.
Sediment event bed & bank load in downstream reaches	"Sed_Evt_BB_Ld_in_DS _Rchs"	48.		blank, T, or F	A1	2	48.
Sediment event gully erosion within cells	"Sed_Evt_Gly_Eros_in_ Cels"	49.		blank, T, or F	A1	2	49.
Sediment event gully yield from cell to receiving reach	"Sed_Evt_Gly_Yld_Cel_t o_Rcv_Rch"	50.		blank, T, or F	A1	2	50.
Sediment event landscape erosion within cells	"Sed_Evt_LS_Eros_in_C els"	51.		blank, T, or F	A1	2	51.
Sediment event landscape load from cell to downstream reaches	"Sed_Evt_LS_Ld_Cel_to _DS_Rchs"	52.		blank, T, or F	A1	2	52.
Sediment event landscape load in reaches	"Sed_Evt_LS_Ld_in_Rc hs"	53.		blank, T, or F	A1	2	53.
Sediment event landscape yield from cell to receiving reach	"Sed_Evt_LS_Yld_Cel_t o_Rcv_Rch"	54.		blank, T, or F	A1	2	54.
Sediment event sheet & rill erosion within cells	"Sed_Evt_SR_Eros_in_C els"	55.		blank, T, or F	A1	2	55.
Sediment event sheet & rill yield from cell to receiving reach	"Sed_Evt_SR_Yld_Cel_t o_Rcv_Rch"	56.		blank, T, or F	A1	2	56.
Water event load from cell to downstream reaches	"Wtr_Evt_Ld_Cel_to_DS _Rchs"	57.		blank, T, or F	A1	2	57.
Water event load in downstream reaches	"Wtr_Evt_Ld_in_DS_Rc hs"	58.		blank, T, or F	A1	2	58.

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.	
Water event Peak discharges in downstream reach	"Wtr_Evt_Pk_Disch_in_ DS_Rch"	59.		blank, T, or F	A1	2	59.	
Water event yield from cell to receiving reach	"Wtr_Evt_Yld_Cel_to_R cv_Rch"	60.		blank, T, or F	A1	2	60.	
Water event baseflow	"Wtr_Evt_Baseflow"	61.		blank, T, or F	A1	2	61.	
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_csv _chld= 1300 +	Par't Field					
(Currently, there are no child files for this section)								
<u>Go to Layout Matrix</u>	Go to Table of Contents							

OUTPUT OPTIONS DATA – DPP

See Attachment A for a detailed explanation

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.					Α	1	1-47	
The following record does not repeat.								

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_d pp_01= 140 0+					
AnnAGNPS_DPP_Accumulation_Setup.dpp	"Acc_Setup"	1.		blank, T, or F	A1	2	1.
Note: This flag produces all of the cell initialization children files listed below but does not produce an output file itself as parent. The reserved filename for this flag is: AnnAGNPS DPP Cell Initialization.dpp	"Cell_Initial"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_DPP_Cell_Time_of_Concentration.dpp	"Cell_TOC"	3.		blank, T, or F	A1	2	3.
AnnAGNPS_DPP_Crop_Growth.dpp	"Crp_Grwth"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_DPP_Pointers_Data_Prep.dpp	"Data_Prep_Pointers"	5.		blank, T, or F	A1	2	5.
Note: This flag produces all of the climate station and weather related children files listed below but does not produce an output file itself as parent. If this flag is set to 'T' then it overrides the parent flag in column 56 and produces the Climate Station Daily Weather file even if that flag is 'F'. The reserved filename for this flag is: AnnAGNPS DPP Weather.dpp	"Weather"	6.		blank, T, or F	A1	2	6.
AnnAGNPS_DPP_Operation_Rotation.dpp	"Opr_Rotation"	7.		blank, T, or F	A1	2	7.
AnnAGNPS_DPP_Pesticide_Metabolite_Reordering.dpp	"Pest_Metabolite"	8.		blank, T, or F	A1	2	8.
AnnAGNPS_DPP_Process_Flag_Set.dpp	"Process_Flag"	9.		blank, T, or F	A1	2	9.
AnnAGNPS_DPP_Quadrature.dpp	"Quadrature"	10.		blank, T, or F	A1	2	10.
AnnAGNPS_DPP_Hydraulic_Geometry.dpp	"Hydraulic_Geom"	11.		blank, T, or F	A1	2	11.
AnnAGNPS_DPP_Reach_Routing_Order.dpp	"Rch_Routing"	12.		blank, T, or F	A1	2	12.
AnnAGNPS_DPP_Reach_Time_of_Concentration.dpp	"Rch_TOC"	13.		blank, T, or F	A1	2	13.
AnnAGNPS_DPP_RUSLE_C_Factors.dpp	"RUSLE_C_Fctr"	14.		blank, T, or F	A1	2	14.
AnnAGNPS_DPP_RUSLE_C_Factors_Soil_Consoldation.dpp	"RUSLE_C_Fctr_SC"	15.		blank, T, or F	A1	2	15.
AnnAGNPS_DPP_RUSLE_Canopy_Cover.dpp	"Canopy_Cover"	16.		blank, T, or F	A1	2	16.
AnnAGNPS_DPP_RUSLE_Crop_Residue.dpp	"Crp_Residue"	17.		blank, T, or F	A1	2	17.
AnnAGNPS_DPP_RUSLE_Dead_Roots.dpp	"Dead_Roots"	18.		blank, T, or F	A1	2	18.
AnnAGNPS_DPP_RUSLE_Preprocessed_C_Factors.dpp	"PreProc_C_Fctr"	19.		blank, T, or F	A1	2	19.
AnnAGNPS_DPP_RUSLE_Dominate_Contour.dpp	"Dom_Contour"	20.		blank, T, or F	A1	2	20.
AnnAGNPS_DPP_RUSLE_EI_Percentages.dpp	"EI_Pcts"	21.		blank, T, or F	A1	2	21.
AnnAGNPS_DPP_RUSLE_Growth_Days.dpp	"RUSLE_Grwth_Days"	22.		blank, T, or F	A1	2	22.
AnnAGNPS_DPP_RUSLE_Initialize_Local_Operations.dpp	"RUSLE_Init_Loc_Oprs	23.		blank, T, or F	A1	2	23.
AnnAGNPS_DPP_RUSLE_K_Factors.dpp	"RUSLE_K_Fctr"	24.		blank, T, or	A1	2	24.
AnnAGNPS_DPP_RUSLE_LS_Factors.dpp (Reserved – Not currently produced)	"Reserved"	25.		blank, T, or	A1	2	25.
AnnAGNPS_DPP_RUSLE_Non_Cropland_C_Factors.dpp	"RUSLE_Non- crp_C_Fctr"	26.		blank, T, or F	A1	2	26.

	Field	Code					
Description	Header for Record #1	Inde x	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_DPP_RUSLE_Num_Soil_Layers_Soil_Residue.dpp	"RUSLE_Num_SLyr_SR es"	27.		blank, T, or F	A1	2	27.
AnnAGNPS_DPP_RUSLE_P_Factors.dpp	"RUSLE_P_Fctr"	28.		blank, T, or F	A1	2	28.
AnnAGNPS_DPP_RUSLE_P_Factors_Contours.dpp	"RUSLE_P_Fctr_Cntrs"	29.		blank, T, or F	A1	2	29.
AnnAGNPS_DPP_RUSLE_P_Factors_Strip.dpp	"RUSLE_P_Fctr_Strp"	30.		blank, T, or	A1	2	30.
AnnAGNPS_DPP_RUSLE_P_Factors_Strip_Rotation.dpp	"RUSLE_P_Fctr_Strp_R ot"	31.		blank, T, or	A1	2	31.
AnnAGNPS_DPP_RUSLE_Prior_Landuse.dpp	"RUSLE_Prior_LU"	32.		blank, T, or	A1	2	32.
AnnAGNPS_DPP_RUSLE_Residue_Coefficients.dpp	"RUSLE_Res_Coef"	33.		blank, T, or	A1	2	33.
AnnAGNPS_DPP_RUSLE_Seg_Residue.dpp	"RUSLE_Seg_Res"	34.		г blank, T, or	A1	2	34.
AnnAGNPS_DPP_RUSLE_Setup_Prd_Seg.dpp	"RUSLE_Setup_Prd_Seg	35.		F blank, T, or	A1	2	35.
AnnAGNPS_DPP_RUSLE_Soil_Moisture.dpp	"RUSLE_Soil_Moisture"	36.		F blank, T, or	A1	2	36.
AnnAGNPS_DPP_RUSLE_Surface_Cover.dpp	"RUSLE_Surf_Cover"	37		F blank, T, or	A1	2	37.
AnnAGNPS_DPP_RUSLE_Surface_Roughness.dpp	"RUSLE_Surf_Rough"	38		F blank, T, or	A1	2	38.
AnnAGNPS_DPP_RUSLE_Unique_Residue.dpp	"RUSLE_Unique_Res"	39		F blank, T, or	A1	2	39.
AnnAGNPS_DPP_Sediment_Particle_Distribution.dpp	"Sed_Part_Distrib"	40		F blank, T, or	A1	2	40.
AnnAGNPS_DPP_Seg_EI_Prcp.dpp	"Seg_EI_Prcp"	41		F blank, T, or	A1	2	41.
AnnAGNPS_DPP_Setup_Seg.dpp	"Setup_Seg"	41.		F blank, T, or	A1	2	42
AnnAGNPS_DPP_Soil_Composite_(surface).dpp	"Soil_Comp_Surf"	42.		F blank, T, or	A 1	2	42.
AnnAGNPS_DPP_Soil_Composite_(layers).dpp	"Soil_Comp_Lyrs"	43.		F blank, T, or	A1	2	44
AnnAGNPS_DPP_Storm_Types.dpp	"Storm_Types"	44.		F blank, T, or	A1	2	44.
AnnAGNPS DPP Climate Station Daily Weather.dpp	"Climate_Daily_Wthr"	43.		F blank, T, or	AI	2	45.
AnnAGNPS DPP Ephemeral Gully Information.csv	"Eph_Gully_Info"	46.		F blank, T, or	AI	2	46.
AnnAGNPS DPP RUSEF2 Information csv	"RUSLE2 Info"	47.		F blank T or	Al	2	47.
		48.		F	A1	2	48.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_d pp_chl d= 1500 +	Par't Field				
AnnAGNPS_DPP_Cell_Initial_Crop_and_Non-Crop.dpp		1.	DPP 01 02				
AnnAGNPS_DPP_Cell_Initial_Frozen_Soil_n_Snow.dpp		2.	DPP 01 02				
AnnAGNPS_DPP_Cell_Initial_Irrigation.dpp		3.	DPP 01 02				
AnnAGNPS_DPP_Cell_Initial_Nutrient.dpp		4.	DPP 01				
AnnAGNPS_DPP_Cell_Initial_Pesticide.dpp		5.	02 DPP 01				
AnnAGNPS_DPP_Cell_Initial_Precipitation.dpp		6.	DPP 01				
AnnAGNPS_DPP_Cell_Initial_RCN_and_Retention.dpp		7.	DPP 01				
AnnAGNPS_DPP_Cell_Initial_Soil_Moisture.dpp		8.	DPP 01 02				

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_DPP_Cell_Initial_Surface_Conditions.dpp		9.	DPP 01 02				
AnnAGNPS_DPP_Cell_Initial_Temperature_Air.dpp		10.	DPP 01 02				
AnnAGNPS_DPP_Cell_Initial_Tile_Drain.dpp		11.	DPP 01 02				
AnnAGNPS_DPP_Climate_Station_Monthly_Precipitation.dpp		12.	DPP 01 06				
AnnAGNPS_DPP_Primary_Station_Initial_Weather.dpp		13.	DPP 01 06				
AnnAGNPS_DPP_Primary_Station_Typical_Weather.dpp		14.	DPP 01 06				
AnnAGNPS_DPP_Ephemeral_Gully_Section_Development.dpp		15.	orphan				
AnnAGNPS_DPP_Cell.dpp (Note: The parent flag for this file is located in the NPT 01 record)		1.	NPT 01 01				
AnnAGNPS_DPP_Reach_Routing_Strahler_Stream_Order.csv		2.	DPP 01 12				
<u>Go to Layout Matrix</u>		<u>Go t</u>	o Table	of Contents			

OUTPUT OPTIONS DATA – EV

See Attachment A for a detailed explanation Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.					Α	1	1-54	
The following record does not repeat.								

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_ev_ 01= 1600 +					
AnnAGNPS_EV_Feedlots_(mass).txt (Reserved – Not currently produced)	"Reserved"	1.		blank, T, or F	A1	2	1.
AnnAGNPS_EV_Feedlots_(ratio).txt (Reserved – Not currently produced)	"Reserved"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_EV_Feedlots_(unit area).txt (Reserved – Not currently produced)	"Reserved"	3.		blank, T, or F	A1	2	3.
AnnAGNPS_EV_Field_Ponds_(mass).txt (Reserved – Not currently produced)	"Reserved"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_EV_Field_Ponds_(ratio).txt (Reserved – Not currently produced)	"Reserved"	5.		blank, T, or F	A1	2	5.
AnnAGNPS_EV_Field_Ponds_(unit area).txt (Reserved – Not currently produced)	"Reserved"	6.		blank, T, or F	A1	2	6.
AnnAGNPS_EV_Gullies_(mass).txt (Reserved – Not currently produced)	"Reserved"	7.		blank, T, or F	A1	2	7.
AnnAGNPS_EV_Gullies_(ratio).txt (Reserved – Not currently produced)	"Reserved"	8.		blank, T, or F	A1	2	8.
AnnAGNPS_EV_Gullies_(unit area).txt (Reserved – Not currently produced)	"Reserved"	9.		blank, T, or F	A1	2	9.
AnnAGNPS_EV_Nitrogen_load_(mass). csv	"EV_N_Ld_Mass"	10		blank, T, or F	A1	2	10.
AnnAGNPS_EV_Nitrogen_load_(ratio). csv	"EV_N_Ld_Ratio"	11		blank, T, or F	A1	2	11.
AnnAGNPS_EV_Nitrogen_load_(unit_area). csv	"EV_N_Ld_UA"	12		blank, T, or F	A1	2	12.
AnnAGNPS_EV_Nitrogen_yield_(mass). csv	"EV_N_Yld_Mass"	13		blank, T, or F	A1	2	13.
AnnAGNPS_EV_Nitrogen_yield_(ratio). csv	"EV_N_Yld_Ratio"	14		blank, T, or F	A1	2	14.
AnnAGNPS_EV_Nitrogen_yield_(unit_area). csv	"EV_N_YId_UA"	15		blank, T, or F	A1	2	15.
AnnAGNPS_EV_Organic_Carbon_load_(mass). csv	"EV_OC_Ld_Mass"	16		blank, T, or F	A1	2	16.
AnnAGNPS_EV_Organic_Carbon_load_(ratio). csv	"EV_OC_Ld_Ratio"	17		blank, T, or F	A1	2	17.
AnnAGNPS_EV_Organic_Carbon_load_(unit_area). csv	"EV_OC_Ld_UA"	18		blank, T, or F	A1	2	18.
AnnAGNPS_EV_Organic_Carbon_yield_(mass). csv	"EV_OC_Yld_Mass"	19		blank, T, or F	A1	2	19.
AnnAGNPS_EV_Organic_Carbon_yield_(ratio). csv	"EV_OC_Yld_Ratio"	20		blank, T, or F	A1	2	20.
AnnAGNPS_EV_Organic_Carbon_yield_(unit_area). csv	"EV_OC_Yld_UA"	21		blank, T, or F	A1	2	21.
AnnAGNPS_EV_Pesticides_load_(mass).txt (Reserved – Not currently produced)	"Reserved"	22		blank, T, or F	A1	2	22.
AnnAGNPS_EV_Pesticides_load_(ratio).txt (Reserved – Not currently produced)	"Reserved"	23		blank, T, or F	A1	2	23.
AnnAGNPS_EV_Pesticides_load_(unit_area).txt (Reserved – Not currently produced)	"Reserved"	24		blank, T, or F	A1	2	24.
AnnAGNPS_EV_Pesticides_yield_(mass).txt (Reserved – Not currently produced)	"Reserved"	25		blank, T, or F	A1	2	25.
AnnAGNPS_EV_Pesticides_yield_(ratio).txt (Reserved – Not currently produced)	"Reserved"	26		blank, T, or F	A1	2	26.
AnnAGNPS_EV_Pesticides_yield_(unit_area).txt (Reserved – Not currently produced)	"Reserved"	27		blank, T, or F	A1	2	27.
AnnAGNPS_EV_Phosphorus_load_(mass). csv	"EV_P_Ld_Mass"	28		blank, T, or F	A1	2	28.

Description	Field Header for	Code Index	Col.	Domain	For	Record	Field
-	Record #1				mat	INO.	INO.
AnnAGNPS_EV_Phosphorus_load_(ratio). csv	"EV_P_Ld_Ratio"	29.		blank, T, or F	A1	2	29.
AnnAGNPS_EV_Phosphorus_load_(unit_area). csv	"EV_P_Ld_UA"	30.		blank, T, or F	A1	2	30.
AnnAGNPS_EV_Phosphorus_yield_(mass). csv	"EV_P_Yld_Mass"	31.		blank, T, or F	A1	2	31.
AnnAGNPS_EV_Phosphorus_yield_(ratio). csv	"EV_P_Yld_Ratio"	32.		blank, T, or F	A1	2	32.
AnnAGNPS_EV_Phosphorus_yield_(unit_area). csv	"EV_P_Yld_UA"	33.		blank, T, or F	A1	2	33.
AnnAGNPS_EV_Point_Sources_(mass).txt (Reserved – Not currently produced)	"Reserved"	34.		blank, T, or F	A1	2	34.
AnnAGNPS_EV_Point_Sources_(ratio).txt (Reserved – Not currently produced)	"Reserved"	35.		blank, T, or F	A1	2	35.
AnnAGNPS_EV_Point_Sources_(unit area).txt (Reserved – Not currently produced)	"Reserved"	36.		blank, T, or F	A1	2	36.
AnnAGNPS_EV_Sediment_Erosion_(mass). csv	"EV_Sed_Eros_Mass"	37.		blank, T, or F	A1	2	37.
AnnAGNPS_EV_Sediment_Erosion_(ratio). csv	"EV_Sed_Eros_Ratio"	38.		blank, T, or F	A1	2	38.
AnnAGNPS_EV_Sediment_Erosion_(unit_area). csv	"EV_Sed_Eros_UA"	39.		blank, T, or F	A1	2	39.
AnnAGNPS_EV_Sediment_Load_(mass). csv	"EV_Sed_Ld_Mass"	40.		blank, T, or F	A1	2	40.
AnnAGNPS_EV_Sediment_Load_(ratio). csv	"EV_Sed_Ld_Ratio"	41.		blank, T, or F	A1	2	41.
AnnAGNPS_EV_Sediment_Load_(unit_area). csv	"EV_Sed_Ld_UA"	42.		blank, T, or F	A1	2	42.
AnnAGNPS_EV_Sediment_Yield_(mass). csv	"EV_Sed_Yld_Mass"	43.		blank, T, or F	A1	2	43.
AnnAGNPS_EV_Sediment_Yield_(ratio). csv	"EV_Sed_Yld_Ratio"	44.		blank, T, or F	A1	2	44.
AnnAGNPS_EV_Sediment_Yield_(unit_area). csv	"EV_Sed_Yld_UA"	45.		blank, T, or F	A1	2	45.
AnnAGNPS_EV_Water_load_(mass). csv	"EV_Wtr_Ld_Mass"	46.		blank, T, or F	A1	2	46.
AnnAGNPS_EV_Water_load_(ratio). csv	"EV_Wtr_Ld_Ratio"	47.		blank, T, or F	A1	2	47.
AnnAGNPS_EV_Water_load_(unit_area). csv	"EV_Wtr_Ld_UA"	48.		blank, T, or F	A1	2	48.
AnnAGNPS_EV_Water_yield_(mass). csv	"EV_Wtr_Yld_Mass"	49.		blank, T, or F	A1	2	49.
AnnAGNPS_EV_Water_yield_(ratio). csv	"EV_Wtr_Yld_Ratio"	50.		blank, T, or F	A1	2	50.
AnnAGNPS_EV_Water_yield_(unit_area). csv	"EV_Wtr_Yld_UA"	51.		blank, T, or F	A1	2	51.
AnnAGNPS_EV_Landscape_Runoff_(all_sources).csv	"EV_LS_Rnof_All_Srcs"	52.		blank, T, or F	A1	2	52.
AnnAGNPS_EV_Landscape_Yield_(all_sources).csv (Reserved - Not currently produced)	"EV_LS_Yld_All_Srcs"	53.		blank, T, or F	A1	2	53.
AnnAGNPS_EV_Gullies_(erosion).csv	"EV_Gullies_Erosion"	54.		blank, T, or F	A1	2	54.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_ev_ chld= 1700 +	Par't Field				
AnnAGNPS_EV_Sediment_load_by_source_(mass). csv		1.	EV 01				
AnnAGNPS_EV_Sediment_load_by_source_(ratio). csv		2	40 EV 01				
AnnAGNPS_EV_Sediment_load_by_source_(unit-area). csv		3.	41 EV 01				

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_EV_Landscape_Runoff_(sheet_rill).csv		4.	EV 01 52				
AnnAGNPS_EV_Landscape_Runoff_(feedlot).csv		5.	EV 01 52				
AnnAGNPS_EV_Landscape_Runoff_(point_source).csv		6.	EV 01 52			-	
AnnAGNPS_EV_Landscape_Runoff_(gully).csv		7.	EV 01				
AnnAGNPS_EV_Landscape_Runoff_(pond).csv		8.	EV 01 52				
AnnAGNPS_EV_Landscape_Runoff_(irrigation).csv		9.	EV 01 52				
AnnAGNPS_EV_Landscape_Yield_(sheet_rill).csv (Not currently produced)		10.	EV 01 53				
AnnAGNPS_EV_Landscape_Yield_(feedlot).csv (Not currently produced)		11.	EV 01 53				
AnnAGNPS_EV_Landscape_Yield_(point_source).csv (Not currently produced)		12.	EV 01 53				
AnnAGNPS_EV_Landscape_Yield_(gully).csv (Not currently produced)		13.	EV 01 53				
AnnAGNPS_EV_Landscape_Yield_(pond).csv (Not currently produced)		14.	EV 01 53				
AnnAGNPS_EV_Landscape_Yield_(irrigation).csv (Not currently produced)		15.	EV 01 53				
The following children files are produced automatically if the associated parent is set to 'T'. The filenames are built internally based on the parent filename and used a different base index.		Unit_nu m_base_ 2= 20000 + base_ev_ 01 +	Par't Field				
AnnAGNPS_EV_Outlet_Sediment_load_(mass). csv		40	EV 01 40				
AnnAGNPS_EV_Outlet_Water_load_(mass). csv		46	EV 01 46				
AnnAGNPS_EV_Outlet_Water_load_(unit-area). csv		48	EV 01 48				
Go to Layout Matrix		Go to	Table	of Contents			

OUTPUT OPTIONS DATA – NPT

See Attachment A for a detailed explanation

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.					Α	1	1-36	
The following record does not repeat.								

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_n pt_01= 180 0 +					
AnnAGNPS_NPT_AnnAGNPS_Identifier.npt	"AnnAGNPS_ID"	1.		blank, T, or F	A1	2	1.
AnnAGNPS_NPT_Cell.npt (Note: this is the parent flag for "AnnAGNPS_DPP_Cell.dpp" listed in the DPP 01 record.)	"Cell"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_NPT_Climate_Station_Information.npt	"Climate_Station"	3.		blank, T, or F	A1	2	3.
AnnAGNPS_NPT_Contour.npt	"Contour"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_NPT_Crop.npt	"Crop"	5.		blank, T, or F	A1	2	5.
AnnAGNPS_NPT_Feedlot_&_Management.npt	"Feedlot"	6.		blank, T, or	A1	2	6.
AnnAGNPS_NPT_Fertilizer_Application_&_Reference.npt	"Fertilizer"	7.		blank, T, or	A1	2	7.
AnnAGNPS_NPT_Management_Sequence.npt	"Mgmt_Seq"	8.		blank, T, or	A1	2	8.
AnnAGNPS_NPT_Field_Pond.npt	"Field_Pond"	9.		blank, T, or	A1	2	9.
AnnAGNPS_NPT_Global_Output_Options.npt	"Glbl_Output_Opts"	10.		blank, T, or	A1	2	10.
AnnAGNPS_NPT_Gully.npt	"Gully"	11.		blank, T, or	A1	2	11.
AnnAGNPS_NPT_Hydraulic_Geometry.npt	"Hydraulic_Geom"	12.		blank, T, or	A1	2	12.
AnnAGNPS_NPT_Impoundment.npt	"Impoundment"	13.		blank, T, or	A1	2	13.
AnnAGNPS_NPT_Irrigation_Application. csv	"Irrigation"	14.		blank, T, or	A1	2	14.
AnnAGNPS_NPT_Landuse_Reference.npt	"Landuse_Ref"	15.		blank, T, or	A1	2	15.
AnnAGNPS NPT 016 reserved.npt	"Reserved"	16.		r blank onlv		2	16.
AnnAGNPS_NPT_Output_Options.npt	"Output_Options"	17.		blank, T, or F	A1	2	17.
AnnAGNPS_NPT_Pesticide_Application_&_Reference.npt	"Pesticide"	18.		blank, T, or F	A1	2	18.
AnnAGNPS_NPT_Point_Source.npt	"Point_Source"	19.		blank, T, or F	A1	2	19.
AnnAGNPS_NPT_Reach.npt	"Reach"	20.		blank, T, or F	A1	2	20.
AnnAGNPS_NPT_Runoff_Curve_Number.npt	"Runoff_Curve_Num"	21.		blank, T, or F	A1	2	21.
AnnAGNPS_NPT_Simulation_Period_Data.npt	"Simulation_Period"	22.		blank, T, or F	A1	2	22.
AnnAGNPS_NPT_Soil_Actual_(Surface).npt	"Soil_Actual_Surface"	23.		blank, T, or	A1	2	23.
AnnAGNPS_NPT_Strip_Crop.npt	"Strip_Crop"	24.		blank, T, or	A1	2	24.
AnnAGNPS_NPT_Tile_Drain.npt	"Tile_Drain"	25.		blank, T, or	A1	2	25.
AnnAGNPS_NPT_Management_Field.npt	"Mgmt_Field"	26.		blank, T, or	A1	2	26.
AnnAGNPS_NPT_Management_Schedule.npt	"Mgmt_Sched"	27.		blank, T, or F	A1	2	27.
AnnAGNPS_NPT_Management_Operation.npt	"Mgmt_Opr"	28.	<u> </u>	blank, T, or F	A1	2	28.

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_NPT_Soil_Actual_(Layers).npt	"Soil_Actual_Layers"	29.		blank, T, or F	A1	2	29.
AnnAGNPS_NPT_Aquaculture_Pond.npt	"Aquaculture_Pond"	30.		blank, T, or F	A1	2	30.
AnnAGNPS_NPT_Aquaculture_Pond_Management_Schedule_A.npt	"Aquaculture_Pond_Mg mt_Schd_A"	31.		blank, T, or F	A1	2	31.
AnnAGNPS_NPT_Global_Error_&_Warning.npt	"Glbl_Err/Wrn"	32.		blank, T, or F	A1	2	32.
AnnAGNPS_NPT_Soil_Initial_Conditions.npt	"Soil_Init_Cond"	33.		blank, T, or F	A1	2	33.
AnnAGNPS_NPT_Pesticide_Inital_Conditions.npt	"Pest_Init_Cond"	34.		blank, T, or F	A1	2	34.
AnnAGNPS_NPT_Wetland.npt	"Wetland"	35.		blank, T, or F	A1	2	35.
AnnAGNPS_NPT_Riparian_Buffers.npt	"Riparian_Buffers"	36.		blank, T, or F	A1	2	36.
AnnAGNPS_NPT_RUSLE2. csv	"RUSLE2"	37.		blank, T, or F	A1	2	37.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_n pt_chld = 1900 +	Par't Field				
AnnAGNPS_NPT_Feedlot.npt		1.	NPT 01 06				
AnnAGNPS_NPT_Feedlot_Management.npt		2.	NPT 01 06				
AnnAGNPS_NPT_Classic_Gully.npt		3.	NPT 01 11				
AnnAGNPS_NPT_Ephemeral_Gully.npt		4.	NPT 01 11				
AnnAGNPS_NPT_Landslide_Gully.npt		5.	NPT 01 11				
AnnAGNPS_NPT_Aquaculture_Pond_Management_Schedule _B.npt		6.	NPT 01 31				
<u>Go to Layout Matrix</u>		<u>Go t</u>	o Table	of Contents			

OUTPUT OPTIONS DATA – SIM

See Attachment A for a detailed explanation

Opti	ional						
Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1-40
The following record does not repeat.							
	Field	Code					
---	-------------------------------	-------------------------------	------	-------------------	-----	--------	--------------
Description	Header for	Inde	Col.	Domain	For	Record	Field No.
	Record #1	X			mat	190.	INU.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_si m_01= 200 0+					
AnnAGNPS_SIM_Cell_Components_Accumulation.sim	"Cell_Components"	1.		blank, T, or F	A1	2	1.
AnnAGNPS_SIM_Conversion_Units.sim	"Conversion_Units"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_SIM_Sheet_&_Rill_Erosion_&_Sediment_Yield.csv	"Sht/Rill_Eros_Sed_Yld"	3.		blank, T, or F	Al	2	3.
AnnAGNPS_SIM_Feedlots.sim	"Feedlots"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_SIM_Insitu_Nitrogen_Inorganic. csv	"Insitu_N_Inorg"	5.		blank, T, or F	A1	2	5.
AnnAGNPS_SIM_Insitu_Nitrogen_Organic. csv	"Insitu_N_Org"	6.		blank, T, or F	A1	2	6.
AnnAGNPS_SIM_Insitu_Residue. csv	"Insitu_Residue"	7.		blank, T, or F	A1	2	7.
AnnAGNPS_SIM_Insitu_Organic_Carbon. csv	"Insitu_OC"	8.		blank, T, or F	A1	2	8.
AnnAGNPS_SIM_Insitu_Phosphorus_Inorganic. csv	"Insitu_P_Inorg"	9.		blank, T, or F	A1	2	9.
AnnAGNPS_SIM_Insitu_Phosphorus_Organic. csv	"Insitu_P_Org"	10.		blank, T, or F	A1	2	10.
AnnAGNPS_SIM_Insitu_Soil_Moisture_Daily_Cell_Data. csv	"Insitu_Soil_Moist_Daily "	11.		blank, T, or F	A1	2	11.
AnnAGNPS_SIM_Irrigation_Applications.sim	"Irrigation"	12.		blank, T, or F	A1	2	12.
AnnAGNPS_SIM_Pesticide_Application.sim	"Pesticide_App"	13.		blank, T, or F	A1	2	13.
AnnAGNPS_SIM_Pesticides_Insitu.sim	"Pesticide_Insitu"	14.		blank, T, or F	A1	2	14.
AnnAGNPS_SIM_Gully.sim	"Gully"	15.		blank, T, or F	A1	2	15.
AnnAGNPS_SIM_Reach_Accumulation_(mass).sim	"Reach_Acc_Mass"	16.		blank, T, or F	A1	2	16.
AnnAGNPS_SIM_Reach_Accumulation_(ratios).sim	"Reach_Acc_Ratio"	17.		blank, T, or F	A1	2	17.
AnnAGNPS_SIM_ Landscape_Yield_(all_sources).sim (Reserved – Not currently produced)	"LS_Yld_All_Srcs"	18.		blank, T, or F	A1	2	18.
AnnAGNPS_SIM_Reach_Loadings_Nutrients. Csv	"Reach_Ld_Nutr"	19.		blank, T, or F	A1	2	19.
AnnAGNPS_SIM_Reach_Loadings_Pesticide.sim (Reserved – Not currently produced)	"Reserved"	20.		blank, T, or F	A1	2	20.
AnnAGNPS_SIM_Reach_Loadings_Sediment. csv	"Reach_Ld_Sed"	21.		blank, T, or F	A1	2	21.
AnnAGNPS_SIM_Reach_Loadings_Water. csv	"Reach_Ld_Wtr"	22.		blank, T, or F	A1	2	22.
AnnAGNPS_SIM_Impoundment_Routings_Part_A.sim	"Impound_Routing_A"	23.		blank, T, or F	A1	2	23.
AnnAGNPS_SIM_Reach_Routing_Nutrients.sim (Reserved – Not currently produced)	"Reserved"	24.		blank, T, or F	A1	2	24.
AnnAGNPS_SIM_Reach_Routing_Pesticide.sim	"Reach_Routing_Pest"	25.		blank, T, or F	A1	2	25.
AnnAGNPS_SIM_Reach_Routing. csv	"Reach_Routing"	26.		blank, T, or F	A1	2	26.
AnnAGNPS_SIM_Reach_Routing_Water.csv (Reserved – Not currently produced)	"Reach_Routing_Wtr"	27.		blank, T, or F	A1	2	27.
AnnAGNPS_SIM_Runoff_Curve_Number. csv	"Runoff_Curve_Num"	28.		blank, T, or F	A1	2	28.

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_SIM_Scheduled_Operations.sim	"Schd_Oprs"	29.		blank, T, or F	A1	2	29.
AnnAGNPS_SIM_Soil_Particle_Distribution.sim	"Soil_Part_Distrib"	30.		blank, T, or F	A1	2	30.
AnnAGNPS_SIM_Pond_Release_n_Yield.sim	"Pond_Release/Yield"	31.		blank, T, or F	A1	2	31.
AnnAGNPS_SIM_Winter_Routines_Thermal_Layers.sim	"Winter_Thermal"	32.		blank, T, or F	A1	2	32.
AnnAGNPS_SIM_Winter_Routines_Summary.sim (Reserved – Not currently produced)	"Reserved"	33.		blank, T, or F	A1	2	33.
AnnAGNPS_SIM_USLE_Parameters. Csv	"USLE_Params"	34.		blank, T, or F	A1	2	34.
AnnAGNPS_SIM_Baseflow.sim	"Baseflow"	35.		blank, T, or F	A1	2	35.
AnnAGNPS_SIM_Insitu_Soil_Moisture_Watershed_Annual_Summaries	"Insitu_Soil_Moist_Wsh d_Sum"	36.		blank, T, or F	A1	2	36.
AnnAGNPS_SIM_Wetland_Effects.csv	"Wetland_Effects"	37.		blank, T, or F	A1	2	37.
AnnAGNPS_SIM_Potential_ET_Adjustment. csv	"Pot_ET_Adjust"	38.		blank, T, or F	A1	2	38.
AnnAGNPS_SIM_Landscape_Runoff_(all_sources).sim (Reserved – Not currently produced)	"LS_Rnof_All_Srcs"	39.		blank, T, or	A1	2	39.
AnnAGNPS_SIM_Riparian_Buffers.csv	"Riparian_Buffers"	40.		blank, T, or F	A1	2	40.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_si m_chld = 2100 +	Par't Field				
AnnAGNPS_SIM_Soil_Moisture_Time_Step_Parameters.sim		1.	SIM 01 11				
AnnAGNPS_SIM_Solar_and_Atmospheric_Radiation.sim		2.	SIM 01 11				
AnnAGNPS_SIM_Insitu_Soil_Moisture_Tile_Drain_Data. csv		3.	SIM 01 11				
AnnAGNPS_SIM_Classic_Gully_erosion.sim		4.	SIM 01 15				
AnnAGNPS_SIM_Classic_Gully_yield.sim		5.	SIM 01 15				
AnnAGNPS_SIM_Ephemeral_Gully_erosion. csv		6.	SIM 01				
AnnAGNPS_SIM_Ephemeral_Gully_yield. csv		7.	SIM 01				
AnnAGNPS_SIM_Landslide_Gully_erosion.sim		8.	SIM 01				
AnnAGNPS_SIM_Landslide_Gully_yield.sim		9.	SIM 01				
AnnAGNPS_SIM_Landscape_Yield_(CCHE1D).sim (Not currently produced)		10.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(sheet_rill).sim (Not currently produced)		11.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(feedlot).sim (Not currently produced)		12.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(point_source).sim (Not currently produced)		13.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(gully).sim (Not currently produced)		14.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(pond).sim (Not currently produced)		15.	SIM 01 18				
AnnAGNPS_SIM_Landscape_Yield_(irrigation).sim (Not currently produced)		16.	SIM 01 18				
AnnAGNPS_SIM_Impoundment_Routings_Part_B.sim		17.	SIM 01 23				
AnnAGNPS_SIM_Scheduled_Events.sim		18.	SIM 01				
AnnAGNPS_SIM_Pond_Sediment_release_and_yield.sim		19.	SIM 01				

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
AnnAGNPS_SIM_Pond_Nutrient_release_and_yield.sim		20.	SIM 01 31				
AnnAGNPS_SIM_Landscape_Runoff_(CCHE1D).sim (Not currently produced)		21.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(sheet_rill).sim (Not currently produced)		22.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(feedlot).sim (Not currently produced)		23.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(point_source).sim (Not currently produced)		24.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(gully).sim (Not currently produced)		25.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(pond).sim (Not currently produced)		26.	SIM 01 39				
AnnAGNPS_SIM_Landscape_Runoff_(irrigation).sim (Not currently produced)		27.	SIM 01 39				
AnnAGNPS_SIM_Ephemeral_Gully_Sections. csv		28.	SIM 01 15				
AnnAGNPS_SIM_Ephemeral_Gully_Yield_to_Mouth. csv		29.	SIM 01 15				
AnnAGNPS_SIM_Ephemeral_Gully_Hydrograph. csv		30.	SIM 01 15				
AnnAGNPS_SIM_Ephemeral_Gully_Nickpoint. csv		31.	SIM 01 15				
AnnAGNPS_SIM_Ephemeral_Gully_Repair_Date. csv		32.	SIM 01 15				
AnnAGNPS_SIM_Insitu_Soil_Moisture_AET_AA_Totals.csv		33.	SIM 01 36				
AnnAGNPS_SIM_Insitu_Soil_Moisture_AET_AA_Monthly_Totals.csv		34.	SIM 01 36				
AnnAGNPS_SIM_Insitu_Soil_Moisture_AET_Monthly_Totals.csv		35.	SIM 01 36				
AnnAGNPS_SIM_Sheet_and_Rill_Erosion.csv		36.	SIM 01 03				
AnnAGNPS_SIM_EI_AA_Totals.csv		37.	SIM 01 03				
AnnAGNPS_SIM_EI_AA_Monthly_Totals.csv		38.	SIM 01				
AnnAGNPS_SIM_EI_Monthly_Totals.csv		39.	SIM 01				
AnnAGNPS_SIM_Insitu_Soil_Moisture_Daily_Cell_Data_ht-H2O.csv		40.	SIM 01				
AnnAGNPS_SIM_Irrigation_Cycle_Parameters_(Automatic).csv		41.	SIM 01				
AnnAGNPS_SIM_Irrigation_Cycle_Parameters_(Manual).csv		42.	SIM 01 12				
AnnAGNPS_SIM_EI_Daily_Values.csv		43.	SIM 01 03				
Go to Layout Matrix		Go t	o Table	of Contents			

OUTPUT OPTIONS DATA – TBL

See Attachment A for a detailed explanation

Optio	onal						
Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1-5
The following record does not repeat.							

Description	Field Header for Record #1	Code Inde x	Col.	Domain	For mat	Record No.	Field No.
The following lines in this section are for parent files with associated flags in the OUTPUT OPTIONS DATA section		Base_t bl_01= 220 0+					
AnnAGNPS_TBL_CCHE1D.txt	"CCHE1D"	1.		blank, T, or F	A1	2	1.
AnnAGNPS_TBL_CONCEPTS.xml	"CONCEPTS_XML"	2.		blank, T, or F	A1	2	2.
AnnAGNPS_TBL_Gaging_Station_Data_Hyd.txt Blank defaults to true; "T".	"Gaging_Station_Hyd"	3.		blank, T, or F	A1	2	3.
AnnAGNPS_TBL_REMM_Input.txt	"REMM"	4.		blank, T, or F	A1	2	4.
AnnAGNPS_TBL_Gaging_Station_Data_Evt.txt	"Gaging_Station_Evt"	5.		blank, T, or F	A1	2	5.
The remaining lines in this section are for children files with no associated flags in the OUTPUT OPTIONS DATA section		Base_t bl_chld = 2300 +	Par't Field				
AnnAGNPS_WRK_TIEGEM.inp			orphan				
Verify_Aqua_Pond_Process.csv			orphan				
Verify_Aqua_Pond_Yield.csv			orphan				
AnnAGNPS_TBL_Gaging_Station_Data_Hyd.csv			TBL 01 03				
Go to Layout Matrix		<u>Go to</u>	o Table	of Contents		-	

OUTPUT OPTIONS DATA – MIN/MAX (LIMITS) See Attachment A for a detailed explanation

0	ational
U	Juonai

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1-12
The following reco	rd does not repeat.						
Minimum Event Date—minimum event date to include runoff data into event output files [mm/dd/yyyy]; defaults to begin simulation date.	"Min_Evt_Date"			mm-1-12 dd-1-31 yyyy-1- 9999	I2/I2 /I4	2	1
Maximum Event Date—maximum event date to include runoff data into event output files [mm/dd/yyyy]; defaults to end simulation date.	"Max_Evt_Date"			mm-1-12 dd-1-31 yyyy-1- 9999	I2/I2 /I4	2	2
Maximum Number Events—maximum number of events to be included in the event output files; defaults to 3000 events in excess of the minimum runoff at the outlet.	"Max_Number_Evts"			0—100000	I10	2	3
Minimum Runoff for Event Output—minimum water runoff at watershed outlet to be included in the event output files, runoff greater than 1mm is always accumulated and included with the average annual data; defaults to 6.35 mm (1/4 in).	"Min_Rnof_Evt"			0 – 508 mm (20 in)	110	2	4
Minimum Runoff for Cell—minimum water runoff for a cell to be included as runoff for the event, accumulated, and included in the average annual data; defaults to 0.10 mm (0.04 in).	"Min_Rnof_Cell"			0 – 508 mm (20 in)	I10	2	5
Minimum Runoff at Outlet—minimum water runoff at watershed outlet to be included as runoff for the event, accumulated, and included in the average annual data; defaults to 0.01 mm (0.004 in).	"Min_Rnof_Outlet"			0 – 508 mm (20 in)	I10	2	6
Minimum Subarea ID—excludes cell & reach data from output files whose subarea ID is outside of this lower limit; defaults to 0. Works only with TopAGNPS generated cell & reach numeric IDs.	"Min_Subarea_ID"			1—100000	I10	2	7
Maximum Subarea ID—excludes cell & reach data from output files whose subarea ID is outside of this upper limit; defaults to 2^31. Works only with TopAGNPS generated cell & reach numeric IDs.	"Max_Subarea_ID"			1—100000	I10	2	8

AnnAGNPS Input File Specifications

Revision: December 4, 2019	File Name: Input_Specifications_v5.51_2019.xx.xx.docx

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Subarea Units Position—includes only those subareas whose ID's unit position is listed. Blank defaults to no exclusion; includes source, leftside, and rightside cells and related reaches. Works only with TopAGNPS generated cell & reach numeric IDs: 0–same as blank; 1– source cells only; 2–leftside cells only; 3–rightside cells only; & 4– reachs only.	"Subarea_Units_Positn"			blank, 0, 1, 2, 3, & 4	I10	2	9
Maximum Number Verification File Accesses—maximum number of write accesses to each verification file; defaults to 1000.	"Max_Vrfy_File_Access"			1—4000	I10	2	10
Maximum Number Verification File Bytes—maximum number of bytes written to each verification file; defaults to 2^24	"Max_Vrfy_File_Bytes"			16,777,216	I10	2	11
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_Code"				I1	2	12
<u>Go to Layout Matrix</u>	Go to Table of Contents						

OUTPUT OPTIONS DATA – CELL

See Attachment A for a detailed explanation

Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1
The following record repe	ats for each requested	d cell.					
Cell ID —ID for cell selected to be included in version 3 output; Must match a valid cell id in the CELL DATA section.	"Cell_ID"				A10	2	1
<u>Go to Layout Matrix</u>	Go to Table of Contents						

OUTPUT OPTIONS DATA – FEEDLOT

See Attachment A for a detailed explanation

Opti	onal						
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1
The following record repeat	s for each requested	feedlot.	_	=	-		
Feedlot ID —ID for feedlot selected to be included in version 3 output; Must match a valid feedlot id in the FEEDLOT DATA section.	"Feedlot_ID"				A10	2	1
Go to Layout Matrix	Go to Table of Contents						

OUTPUT OPTIONS DATA – FIELD POND

See Attachment A for a detailed explanation

Opti	onal						
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1
The following record repeats	for each requested fie	eld pond	•				
Field Pond ID —ID for feedlot selected to be included in version 3 output; Must match a valid feedlot id in the FEEDLOT DATA section.	"Field_Pond_ID"				A10	2	1
<u>Go to Layout Matrix</u>	Go to Table of Contents						

OUTPUT OPTIONS DATA – GULLY, CLASSIC

See Attachment A for a detailed explanation

Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.		
Field Header— Required unique field header for each field listed below.					Α	1	1		
The following record repeats for each requested classic gully.									
Classic Gully ID —ID for gully selected to be included in version 3 output; Must match a valid gully id in the CLASSIC GULLY DATA section.	"Classic_Gully_ID"				A10	2	1		
<u>Go to Layout Matrix</u>	Go to Table of Contents								

OUTPUT OPTIONS DATA – GULLY, EPHEMERAL

See Attachment A for a detailed explanation

Opti	onal						
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.					Α	1	1
The following record repeats for	each requested epher	neral gu	illy.				
Ephemeral Gully ID —ID for gully selected to be included in version 3 output; Must match a valid gully id in the EPHEMERAL GULLY DATA section.	"Ephemeral_Gully_ID"				A10	2	1
<u>Go to Layout Matrix</u>	Go to Table of Contents						

OUTPUT OPTIONS DATA – IMPOUNDMENT

See Attachment A for a detailed explanation

Opti	onal								
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.		
Field Header— Required unique field header for each field listed below.					Α	1	1		
The following record repeats fo	The following record repeats for each requested impoundment.								
Impoundment ID —ID for impoundment selected to be included in version 3 output; Must match a valid impoundment id in the IMPOUNDMENT DATA section.	"Impoundment_ID"				A10	2	1		
Go to Layout Matrix	Go to Table of Contents								

OUTPUT OPTIONS DATA – POINT SOURCE

See Attachment A for a detailed explanation

Opti	onal							
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.					Α	1	1	
The following record repeats for	or each requested poi	nt sourc	e.		_	-		
Point Source ID —ID for point source selected to be included in version 3 output; Must match a valid point source id in the POINT SOURCE DATA section.	"Point_Source_ID"				A10	2	1	
<u>Go to Layout Matrix</u>	Go to Table of Contents							

OUTPUT OPTIONS DATA – REACH

See Attachment A for a detailed explanation

Opti	onal								
Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.		
Field Header— Required unique field header for each field listed below.					Α	1	1		
The following record repea	ts for each requested	reach.							
Reach ID —ID for reach selected to be included in version 3 output; Must match a valid reach id in the REACH DATA section.	"Reach_ID"				A10	2	1		
<u>Go to Layout Matrix</u>	Go to Table of Contents								

OUTPUT OPTIONS DATA – WETLAND

See Attachment A for a detailed explanation

Optional

Description	Field Header for Record #1	Code Index	Col.	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.					Α	1	1	
The following record repeat	s for each requested v	vetland.		-				
Wetland ID—ID for wetland selected to be included in version 3 output; Must match a valid wetland id in the WETLAND DATA section.	"Wetland_ID"				A10	2	1	
<u>Go to Layout Matrix</u>	Go to Table of Contents							

PESTICIDE APPLICATION DATA

Optional unless referenced in Management Schedule Data or Simulation Period Data

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.				
Field Header— Required unique field header for each field listed below.				А	1	1-8				
The following record repeats for the number of pesticide application sets.										
Pesticide Application ID —Alphanumeric string identifying the pesticide application.	"Application_ID			A10	2	1				
Pesticide ID —Alphanumeric string identifying the pesticide. Must be the same as a pesticide reference ID (in Pesticide Reference Data).	"Reference_ID"			A40	2	2				
Pesticide Rate —Application rate for pesticide active ingredient.	"Application_R ate"	{lb / acre} [kg / hectare]	{0.0 to 100.0} [0.0 to 112.0]	F10	2	3				
Pesticide Depth —Soil depth to which is pesticide is applied. Zero depth implies the pesticide was not incorporated in the soil.	"Depth"	{in} [mm]	{0.0 to 60.0} [0.0 to 1500.0]	F10	2	4				
Pesticide Mixing code - Indicator of whether pesticide is mixed uniformly between the soil surface and the depth of incorporation: Acceptable values are: Y- yes, N—no (Blank indicates yes)	"Mixing_Code"		Blank, Y, or N	A10	2	5				
Pesticide Foliage Fraction —Decimal fraction of pesticide applied to the foliage	"Foliage_Fracti on"		0.0 to 1.0	F10	2	6				
Pesticide Soil Fraction—Decimal fraction of pesticide applied to the ground	"Soil_Fraction"		0.0 to 1.0	F10	2	7				
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	8				
Go to Layout Matrix		Go to	Table of Conten	ts						

PESTICIDE INITIAL CONDITIONS DATA

Optional

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.				
Field Header— Required unique field header for each field listed below.				А	1	1-5				
The following record repeats for the number of initial pesticide records. Enter only for pesticides that require other than the global initial condition defaults.										
Initial Pesticide ID —Alphanumeric string identifying the pesticide initially in the soil profile.	"Initial_Pesticid e_ID"			A40	10	1				
Crop Initial Pesticide Amount —Initial pesticide amount to be used for each cropland cell. Two soil layers are used first is top 8 in (200 mm), second is remaining soil profile. (Blank defaults to 0.0)	"Crop_Initial_A mount_1", Crop_Initial_A mount_2	mass-pest / mass-soil (nd)	Blank, or 0.0 to 1.0	2F10	10	2-3				
Non-crop Initial Pesticide Amount —Initial pesticide amount to be used for each non-cropland cell. Two soil layers are used first is top 8 in (200 mm), second is remaining soil profile. (Blank defaults to 0.0)	"Non- Crop_Initial_A mount_1", Non- Crop_Initial_A mount_2	mass-pest / mass-soil (nd)	Blank, or 0.0 to 1.0	2F10	10	4-5				
Go to Layout Matrix	Go to Table of Contents									

PESTICIDE REFERENCE DATA

Optional unless referenced in Pesticide Application Data or Simulation Period Data

Description	Field Header for Record #1	Units	Domain		Record No.	Field No.						
Field Header— Required unique field header for each field listed below.				А	1	1-9						
The following record repeats for t	The following record repeats for the number of pesticide references.											
Pesticide Reference ID —Alphanumeric string identifying the pesticide.	"Pesticide_Refer ence_ID"			A40	2	1						
Pesticide Solubility —Solubility of the pesticide in water; weight of the pesticide divided by the weight of the total solution (water).	"Solubility"	wt / wt (nd)	0 .0 to 100000000	F10	2	2						
Pesticide Partition —Soil pesticide partitioning coefficient "Koc" normalized for organic carbon.	"Partition"	ml/g	0.0 to 100000000.	F10	2	3						
Pesticide Soil Half-life —Time it takes half of the pesticide to degrade into or on the soil. This combines all the degradation methods e.g. chemical, biological, photo.	"Soil_Half-life"	days	0.1 to 50000.0	F10	2	4						
Pesticide Foliage Half-life —Time it takes half of the pesticide to degrade on the foliage. This is a combination of all degradation methods e.g. chemical, biological, photo.	"Foliage_Half- life"	days	0.1 to 50000.0	F10	2	5						
Pesticide Washoff —Fraction of pesticide that washes foliage once a threshold value of 0.1 inch (2.5 mm) of rainfall or spray irrigation is exceeded.	"Washoff"		0.0 to 1.0	F10	2	6						
Metabolite ID —Common scientific name of the resulting chemical after breakdown of the parent compound. Must be the same as a pesticide reference ID (in Pesticide Reference Data). Leave blank if no metabolite is to be considered.	"Metabolite_ID "			A40	2	7						
Metabolite Transformation —Fraction of pesticide that is transformed to the metabolite in one day. Leave blank if no Metabolite ID is provided. Blank (if a Metabolite ID is provided) defaults to 1.0	"Metabolite_Tr ansformation"		Blank or 0.0 to 1.0	F10	2	8						
Pesticide Reach Half-life —Time it takes half of the pesticide to degrade while in a reach (channel). This combines all the degradation methods e.g. chemical, biological, photo. Blank defaults to no pesticide degradation in the reach.	"Reach_Half- life"	days	Blank or 0.1 to 50000.0	F10	2	9						
<u>Go to Layout Matrix</u>		<u>Go to</u>	o Table of Conten	ts								

PL CALIBRATION DATA

Optional

				-	-					
	Field	Units	Domain	For	Record	Field				
Description	Header for	{English}	{English} [SI]	mat	No.	No.				
	Record #1	[51]								
Field Header— Required unique field header for each field listed				Α	1	1-28				
below.		· · · · Ph · · · · ·								
The following record repeats for watersned-scale calibration factors.										
Fields 1-/ are from of Organic contact from of generation factor, if	ganic carbon	sources.	Diants 0.0 an	E10	2	1				
present will be used as the default for organic carbon within field	es"	(na)	00000001 to	F10	Z	1				
where blank fields exist in this record from sheet & rill, feedlot,			9999999999.							
point source, gully, pond, and irrigation. If blank, coded defaults										
will be used. A "0.0" (at least one zero & a decimal) means that										
none of this pollutant originates from this source. A complete field										
("9999999999."), which is AnnAGNPS' real-number input for										
infinity, means that all physically-possible pollutant originates										
instantaneously from this source for each runoff event. Any real-										
number smaller than this "infinity" is taken to be exactly the value										
of this real-number.	"OC Shoot and	(D11-00	E10	2	2				
Organic carbon from sneet & rill – I his calibration factor, if	_Rill"	(nd)	Blank, 0.0 , or 0.000001 to	FIU	2	2				
If blank, the organic carbon calibration factor from sheet & rill			99999999999999.							
sources will be used. A "0.0" (at least one zero & a decimal) means										
that none of this pollutant originates from this source. A complete										
field for the real-number input consisting of nine 9's & a decimal										
(999999999),), which is AnnAGNPS real-number input for infinity means that all physically-possible pollutant originates										
instantaneously from this source for each runoff event. Any real-										
number smaller than this "infinity" is taken to be exactly the value										
of this real-number.										
Organic carbon from feedlot —This calibration factor, if present,	"OC_Feedlot"	(nd)	Blank, 0.0, or	F10	2	3				
watershed-scale global calibration factor will be used A "0 0" (at			.00000000110							
least one zero & a decimal) means that none of this pollutant			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,							
originates from this source. A complete field for the real-number										
input consisting of nine 9's & a decimal ("999999999."), which is										
AnnAGNPS' real-number input for infinity, means that all										
source for each runoff event. Any real-number smaller than this										
"infinity" is taken to be exactly the value of this real-number.										
Organic carbon from point source—This calibration factor, if	"OC_Point_Sou	(nd)	Blank, 0.0, or	F10	2	4				
present, will be used for all point source sources of organic carbon.	rce		.000000001 to							
If blank, the watershed-scale global calibration factor will be used. $\Lambda "0.0"$ (at least one zero & a decimal) means that none of this			99999999999.							
pollutant originates from this source. A complete field for the real-										
number input consisting of nine 9's & a decimal ("999999999."),										
which is AnnAGNPS' real-number input for infinity, means that all										
physically-possible pollutant originates instantaneously from this										
"infinity" is taken to be exactly the value of this real-number										
Organic carbon from gully—This calibration factor, if present,	"OC_Gully"	(nd)	Blank, 0.0, or	F10	2	5				
will be used for all gully sources of organic carbon yield to their		· · ·	.000000001 to							
mouths. If blank, the watershed-scale global calibration factor will			9999999999.							
be used. A "0.0" (at least one zero & a decimal) means that none of										
real-number input consisting of nine 9's & a decimal										
("999999999."), which is AnnAGNPS' real-number input for										
infinity, means that all physically-possible pollutant originates										
instantaneously from this source for each runoff event. Any real-										
number smaller than this "infinity" is taken to be exactly the value				1						

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Organic carbon from pond —This calibration factor, if present, will be used for all pond sources of organic carbon yield to their outlets. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"OC_Pond"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	6
Organic carbon from irrigation—This calibration factor, if present, will be used for all irrigation sources of organic carbon. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real- number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"OC_Irrigation "	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	7
Nitrogen from all sources —This calibration factor, if present, will be used as the default for nitrogen within field where blank fields exist in this record from sheet & rill, feedlot, point source, gully, pond, and irrigation. If blank, coded defaults will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_All_Sources	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	8
Nitrogen from sheet & rill —This calibration factor, if present, will be used for all sheet & rill sources of nitrogen. If blank, the nitrogen calibration factor from all sources will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Sheet_and_ Rill"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	9
Nitrogen from feedlot—This calibration factor, if present, will be used for all feedlot sources of nitrogen. If blank, the watershed- scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Feedlot"	(nd)	Blank, 0.0, or .000000001 to 999999999.	F10	2	10

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Nitrogen from point source —This calibration factor, if present, will be used for all point source sources of nitrogen. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number	"N_Point_Sourc e"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	11
Nitrogen from gully —This calibration factor, if present, will be used for all gully sources of nitrogen yield to their mouths. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Gully"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	12
Nitrogen from pond —This calibration factor, if present, will be used for all pond sources of nitrogen yield to their outlets. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Pond"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	13
Nitrogen from irrigation —This calibration factor, if present, will be used for all irrigation sources of nitrogen. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"N_Irrigation"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	14
Fields 15-21 are from	phosphorus sourc	ces.				
Phosphorus from all sources —1 his calibration factor, if present, will be used as the default for phosphorus within field where blank fields exist in this record from sheet & rill, feedlot, point source, gully, pond, and irrigation. If blank, coded defaults will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"r_AII_Sources	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	15

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Phosphorus from sheet & rill —This calibration factor, if present, will be used for all sheet & rill sources of phosphorus. If blank, the phosphorus calibration factor from all sources will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Sheet_and_ Rill"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	16
Phosphorus from feedlot —This calibration factor, if present, will be used for all feedlot sources of phosphorus. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Feedlot"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	17
Phosphorus from point source —This calibration factor, if present, will be used for all point source sources of phosphorus. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Point_Sourc e"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	18
Phosphorus from gully —This calibration factor, if present, will be used for all gully sources of phosphorus yield to their mouths. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real- number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Gully"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	19
Phosphorus from pond —This calibration factor, if present, will be used for all pond sources of phosphorus yield to their outlets. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Pond"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	20
Phosphorus from irrigation —This calibration factor, if present, will be used for all irrigation sources of phosphorus. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"P_Irrigation"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	21

			-			
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Sediment from all sources—This calibration factor, if present, will be used as the default for erosion/sediment yield within field where blank fields exist in this record from sheet & rill, feedlot, point source, gully, pond, and irrigation. If blank, coded defaults will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real- number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_All_ Sources"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	22
Sediment from sheet & rill—This calibration factor, if present, will be used for all sheet & rill sources of erosion. If blank, the watershed-scale sediment from all sources will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Shee t_and_Rill"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	23
Sediment from feedlot —This calibration factor, if present, will be used for all feedlot sources of erosion. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Feed lot"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	24
Sediment from point source—This calibration factor, if present, will be used for all point source sources of erosion. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Poin t_Source"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	25
Sediment from gully —This calibration factor, if present, will be used for all gully sources of sediment yield to their mouths. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real- number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Gull y"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	26

				-		
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Sediment from pond—This calibration factor, if present, will be used for all pond sources of sediment yield to their outlets. If blank, the watershed-scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("9999999999."), which is AnnAGNPS' real-number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Pon d"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	27
Sediment from irrigation—This calibration factor, if present, will be used for all irrigation sources of erosion. If blank, the watershed- scale global calibration factor will be used. A "0.0" (at least one zero & a decimal) means that none of this pollutant originates from this source. A complete field for the real-number input consisting of nine 9's & a decimal ("999999999."), which is AnnAGNPS' real- number input for infinity, means that all physically-possible pollutant originates instantaneously from this source for each runoff event. Any real-number smaller than this "infinity" is taken to be exactly the value of this real-number.	"Sediment_Irrig ation"	(nd)	Blank, 0.0, or .000000001 to 9999999999.	F10	2	28
<u>Go to Layout Matrix</u>		<u>Go to</u>	Table of Conten	ts		

POINT SOURCE DATA

Optional

00.							
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.				А	1	1-11	
The following record repeats for the number of point sources. Multiple point sources for a cell should be consecutive records here.							
Point Source ID —Alphanumeric string identifying the Point Source.	"Point_Source_I D"			A10	2	1	
Point Cell ID —Alphanumeric string identifying cell that contains the point source. Must be the same as a cell ID in the CELL DATA section.	"Cell_ID"			A10	2	2	
Point Flow —Constant runoff flow rate from point source.	"Point_Flow"	cfs $[m^3 / sec]$	{0.0033 to 10000.0} [0.00001 to 300.0]	F10	2	3	
Point Nitrogen —Concentration of elemental nitrogen in solution in the discharge	"Point_N"	ppm	0.0 to 1000.0	F10	2	4	
Point Phosphorus —Concentration of elemental phosphorus in solution in the discharge	"Point_P"	ppm	0.0 to 1000.0	F10	2	5	
Point Organic Carbon —Concentration of organic Carbon in solution in the discharge	"Point_OC"	ppm	0.0 to 100000.0	F10	2	6	
Organic Carbon Calibration Factor —used to calibrate point source organic carbon for this point source only. Defaults to the watershed-scale organic carbon from point sources calibration factor in the PL CALIBRATION DATA section.	"OC_Calib_Fctr	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	7	
Nitrogen Calibration Factor—used to calibrate point source nitrogen for this point source only. Defaults to the watershed-scale nitrogen from point sources calibration factor in the PL CALIBRATION DATA section.	"N_Calib_Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	8	
Phosphorus Calibration Factor —used to calibrate point source phosphorus for this point source only. Defaults to the watershed- scale phosphorus from point sources calibration factor in the PL CALIBRATION DATA section.	"P_Calib_Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	9	

AnnAGNPS Input File Specifications

Revision: December 4, 2019	File Name: Input_Specifications_v5.51_2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Erosion Calibration Factor—used to calibrate point source erosion for this point source only. Defaults to the watershed-scale sediment from point sources calibration factor in the PL CALIBRATION DATA section.	"Erosion_Calib_ Fctr"	(nd)	Blank, 0.0 or .000000001 to 9999999999.	F10	2	10
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	11
Go to Layout Matrix	Go to Table of Contents					

⁸RCN CALIBRATION DATA:

Optional unless referenced in Cell Data

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.				А	1	1-9	
The following record repeats	for each RCN	calibration	ID				
RCN Calibration ID —ID for an RCN calibration subwatershed. The 1 st record must have an ID of "WATERSHED" which is to apply to all cells not otherwise associated with an RCN calibration area.	"RCN_Calib_ID "			A10	2	1	
Target Average Annual Direct Runoff Load —the desired average annual direct (surface plus quick return flow) runoff at the subwatershed outlet associated with the RCN calibration ID. Blank defaults to no calibration iterations.	"Target_AA_Di rect_Runoff_Lo ad"	{in} [mm]	Blank, or {0.0 to 472.44} [0.0 to 12000.]	F10	2	2	
Previous Estimate for the Average Annual Direct Runoff Load—the average annual direct (surface plus quick return flow) runoff load at the subwatershed outlet associated with the RCN retention factor from a previous execution. Blank defaults to no previous estimate. Currently not used - Reserved.	"Prev_Estimate _for_Water_Loa d"	{in} [mm]	Blank, or {0.0 to 472.44} [0.0 to 12000.]	F10	2	3	
RCN Retention factor —RCN retention variable calibration factor applied to all cells above the subwatershed's outlet not otherwise specifically calibrated. The value for this field is used as the starting retention calibration factor if the target water load is given. Blank defaults to 1.	"RCN_Retentio n_Fctr"	(nd)	Blank, or 0 to ∞	F10	2	4	
Reach ID Site—Reach ID of outlet for the RCN Calibration ID subwatershed. Blank defaults to watershed outlet for one RCN Calibration ID only.	"Reach_ID_Site "			A10	2	5	
Reach Ratio —ratio of the reach's local cells' drainage area contribution at the RCN calibration site to the reach's total local cells' drainage area at the downstream end of the reach. (Blank defaults to 1.)	"Reach_Ratio"	(nd)	Blank, 0. to 1.	F10	2	6	
Available Soil Moisture, AMC-II —available soil moisture for AMC-II used to determine the daily RCN. Blank defaults to 0.50.	"Avbl_Soil_Moi st_AMC_II"	(nd)	Blank, 0. to 1.	F10	2	7	
Target Average Annual Total Streamflow —the desired total streamflow (direct runoff plus baseflow). Blank defaults to no baseflow. Currently not used - Reserved.	"Target_AA_To tal_Streamflow"	{in} [mm]	Blank, or {0.0 to 472.44} [0.0 to 12000.]	F10	2	8	
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	9	
Go to Layout Matrix	Go to Table of Contents						

⁸ This section must be placed immediately following the GLOBAL IDS, FACTORS, AND FLAGS DATA section.

REACH DATA

Required

Kee	uneu					
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.	International In			A	1	1-28
The following record repea	ts for the numl	oer of reach	es.	-	<u>-</u>	
Reach ID —Alphanumeric string identifying the channel reach.	"Reach_ID"			A10	2	1
Receiving Reach ID —Alphanumeric string identifying the receiving reach. Must be the same as a reach ID in the REACH DATA section or "Outlet". Use "Outlet" for the flow from the downstream most reach in the watershed.	"Receiving_Rea ch"			A10	2	2
Reach Vegetation code—Acceptable values are: 0 = Reach is not vegetated. 1 = Reach is vegetated 2 = Reach is submerged due to an impoundment. May be blank if Reach Manning's N is entered for reach. Blank defaults to 1.	"Vegetation_Co de"		Blank, 0,1 or 2	I10	2	3
Reach Elevation —Elevation of the downstream end of the reach	"Elevation"	{ft} [m]	{-3280.8. to 32808.3} [-1000.0 to 10000.0]	F10	2	4
Reach Slope —Average channel slope for the reach.	"Slope"	len-vert / len-horz (nd)	0.00001 to 10.0	F10	2	5
Reach Manning's n—Representative roughness coefficient for reach. Blank defaults to: 0.04—(for Reach Vegetation code of 1) 0.02—(for Reach Vegetation code of 0 or 2) If Reach Vegetation code is blank and Reach Manning's n is blank then this value defaults to 0.04.	"Mannings_n"		Blank, or 0.005 to 1.0	F10	2	6
Reach Infiltration Rate —Infiltration rate of the reach bottom. (Blank defaults to 0.0) Currently not used - Reserved.	"Infiltration_Ra te"	{in/hr} [mm/hr]	Blank, or {0.0 to 100.0 } [0.0 to 2540.0]	F10	2	7
Hydraulic Geometry ID —Alphanumeric string identifying the hydraulic geometry data. Must be the same as a Hydraulic Geometry ID in the HYDRAULIC GEOMETRY DATA section. Leave blank if the reach length, width, depth and valley width are entered unless required for cells contributingl directly to the reach and the cell's time of concentration (T _c) is not entered. Blank defaults to the default Hydraulic Geometry ID in Simulation Period Data section.	"Hydraulic_Geo m_ID"			A10	2	8
Reach Length —Length of the channel reach. Blank indicates AnnAGNPS computes from geometric relationship based on reach geometry ID specified above.	"Length"	{ft} [m]	Blank or {0.0 to 999999999.9} [0.0 to 30480000.0]	F10	2	9
Reach Top Width —Top width of the channel at bank full flow. Blank indicates AnnAGNPS computes from geometric relationship based on reach geometry ID specified above.	"Top_Width"	{ft} [m]	Blank or {0.0 to 984.} [0.0 to 300.0]	F10	2	10
Reach Flow Depth —Flow depth of the channel at bank full flow. Blank indicates AnnAGNPS computes from geometric relationship based on reach geometry ID specified above.	"Flow_Depth"	{ft} [m]	Blank or {0.0 to 984.} [0.0 to 300.0]	F10	2	11
Valley Width—Width of the floodplain. The width of the floodplain entered includes the bankfull top width of the channel, which will be subtracted out in AnnAGNPS. Blank indicates AnnAGNPS computes from geometric relationship based on reach geometry ID specified above.	"Valley_Width"	{ft} [m]	Blank or {0.0 to 98425.} [0.0 to 30000.0]	F10	2	12
Valley n—Floodplain Manning's "n" roughness coefficient. Blank defaults to 0 150	"Valley_Mannin gs_n"		Blank or 0.005 to 1.0	F10	2	13

	T• 1 1	T T •/		-		
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Start Diversion —Reach flow rate above which water is diverted from the reach to a sink/diversion. Water discharged to a sink/diversion is lost from the watershed. Zero or blank signifies no flow is diverted to the sink. Currently not used.	"Start_Diversio n"	{cfs} [m ³ /sec]	Blank or {0.0 to 35287552.} [0.0 to 1000000.0]	F10	2	14
Stop Diversion —Maximum capacity of the sink/diversion. Flows exceeding the maximum sink capacity continue down the reach. Zero or blank indicates no sink. Currently not used.	"Stop_Diversion "	{cfs} [m ³ /sec]	Blank or {0.0 to 35287552.} [0.0 to 1000000.0]	F10	2	15
Travel Time —travel time through reach. Default is to use the reach hydraulic parameters and assume uniform flow at bank full discharge.	"Travel_Time"	hr	Blank or 0.0 to 50 hr	F10	2	16
Clay Scour code—Code indicating if clay sized particles are to be allowed to scour the reach channel. Acceptable codes are : Y = Yes $N = No$ (Blank defaults to N) Currently not used.	"Clay_Scour_C ode"		Blank, Y, or N	A2	2	17
Silt Scour code—Code indicating if silt sized particles are to be allowed to scour the reach channel. Acceptable codes are : Y = Yes $N = No$ (Blank defaults to N)	"Silt_Scour_Co de"		Blank, Y, or N	A2	2	18
Sand Scour codeCode indicating if sand sized particles are to beallowed to scour the reach channel. Acceptable codes are : $Y = Yes$ $N = No$ (Blank defaults to N)	"Sand_Scour_C ode"		Blank, Y, or N	A2	2	19
Small Aggregate Scour codeCode indicating if small aggregatesized particles are to be allowed to scour the reach channel.Acceptable codes are : $Y = Yes$ $N = No$ (Blank defaults toN)	"Small_Agg_Sco ur_Code"		Blank, Y, or N	A2	2	20
Large Aggregate Scour code —Code indicating if large aggregate sized particles are to be allowed to scour the reach channel. Acceptable codes are: $Y = Yes$ $N = No$ (Blank defaults to N)	"Large_Agg_Sc our_Code"		Blank, Y, or N	A2	2	21
Valley Clay Scour code—Code indicating if clay sized particles are to be allowed to scour the reach valley (excluding channel). Acceptable codes are: Y = Yes N = No (Blank defaults to N) Currently not used.	"Valley_Clay_S cour_Code"		Blank, Y, or N	A2	2	22
Valley Silt Scour code —Code indicating if silt sized particles are to be allowed to scour the reach valley (excluding channel). Acceptable codes are : $Y = Yes$ N = No (Blank defaults to N)	"Valley_Silt_Sco ur_Code"		Blank, Y, or N	A2	2	23
Valley Sand Scour codeCode indicating if sand sized particlesare to be allowed to scour the reach valley (excluding channel).Acceptable codes are : $Y = Yes$ N = No(Blank defaults to N)	"Valley_Sand_S cour_Code"		Blank, Y, or N	A2	2	24
Valley Small Aggregate Scour codeAggregate sized particles are to be allowed to scour the reach valley(excluding channel). Acceptable codes are : $Y = Yes$ $N = No$ (Blank defaults to N)	"Valley_Small_ Agg_Scour_Cod e"		Blank, Y, or N	A2	2	25
Valley Large Aggregate Scour codeCode indicating if largeaggregate sized particles are to be allowed to scour the reach valley(excluding channel). Acceptable codes are : $Y = Yes$ $N = No$ (Blank defaults to N)	"Valley_Large_ Agg_Scour_Cod e"		Blank, Y, or N	A2	2	26
Delivery Ratio —Delivery ratio from all sources of sediment load at the reach's upstream end to its downstream end. Recommended procedure is the sediment transport algorithm which is the default procedure when this field is left blank.	"Delivery_Ratio "	(nd)	Blank, or 0. to 1.	F10	2	27
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	28
Go to Layout Matrix		Go t	o Table of Conten	ts		

REACH NUTRIENT HALF-LIFE

Optional

Description Field Header for each field listed	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
below.						
The following reco	rd does not re	peat.		-	-	
Reach Nitrogen Half-life —Time it takes half of the Nitrogen to degrade while in a reach (channel). This combines all the degradation methods e.g. chemical, biological, photo. Blank defaults to 730.0 days.	"N_Half-life"	days	Blank or 0.0 to 50000.0	F10	2	1
Reach Phosphorus Half-life —Time it takes half of the Phosphorus to degrade while in a reach (channel). This combines all the degradation methods e.g. chemical, biological, photo. Blank defaults to 730.0 days.	"P_Half-life"	days	Blank or 0.0 to 50000.0	F10	2	2
Reach Organic Carbon Half-life —Time it takes half of the organic Carbon to degrade while in a reach (channel). This combines all the degradation methods e.g. chemical, biological, photo. Blank defaults to 730.0 days.	"OC_Half-life"	days	Blank or 0.0 to 50000.0	F10	2	3
<u>Go to Layout Matrix</u>		<u>Go to</u>	Table of Conten	<u>ts</u>	_	

RIPARIAN BUFFER DATA

Optional

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.				А	1	1-20	
The following record repeats for the number of buffer records.							
Buffer ID— Alphanumeric string identifying the buffer record.	"Buffer_ID"			A40	2	1	
Location ID —Alphanumeric string identifying the cell or reach that contains the buffer. Must match a cell ID in the Cell Data section for cell-located buffers or a Reach ID in the Reach Data section for reach-located buffers.	"Location_ID"			A10	2	2	
Vegetative Type —Alphanumeric string identifying the vegetative type. The acceptable values are "forest" and "grass" (Blank defaults to "grass")	"Vegetative_Ty pe"		Blank, or "forest" or "grass"	F10	2	3	
Slope — Slope of the buffer. (For cell located buffers, blank defaults to slope of the cell. For reach located buffers, blank defaults to the reach slope.)	"Buffer_Slope"	[nd]	Blank, or 0.00001 to 3.0	F10	2	4	
Maximum Trapping Efficiency "TE-m"—Maximum buffer trapping efficiency. If the user enters a value for this parameter then values for buffer width and slope are ignored. (Blank indicates that AnnAGNPS will calculate value based on vegetative type and slope.)	"Max_Trap_Eff iciency"	[nd]	Blank, or [0.0 to 1.0]	F10	2	5	
Effective Buffer Width —Effective width of the buffer associated with the length of the flow path through the buffer. (Blank defaults to 10.0 m).	"Eff_Wdth_Thr u_Buffer"	{ft} [m]	Blank, or {0.0 to 5000.} [0.0 to 5000.]	F10	2	6	
Effective Concentrated Flow Width—Effective width for the concentrated flow path through the buffer. (For cell located buffers, blank defaults to the hydraulic geometry width of concentrated flow in the cell. For reach located buffers, blank defaults to the reach width.)	"Eff_Wdth_Alo ng_Buffer"	{ft} [m]	Blank, or {0.0 to 5000.} [0.0 to 5000.]	F10	2	7	
Buffer Location Code — If buffer is cell-located, code must be a "T" or blank. If buffer is reach-located, code must be an "F". Blank defaults to true "T"	"Buffer_Locatio n_Code"		Blank, "T" or "F"	L1	2	8	

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Drainage Area to Upstream Portion of Buffer —The total drainage area contributing to the flow entering the upstream portion of the buffer. For cell-located buffers, blank defaults to the cell's entire drainage area. For reach-located buffers, blank defaults to the drainage area at the downstream end of the reach.	"Drainage_Area _to_Buffer"	{acres} [hectares]	Blank, or {0.000025 to 9884.} [0.00001 to 4000.0]	F10	2	9
Actual Trapping Efficiency "TE-a"—Actual trapping efficiency for each particle size. If the user enters a value for this parameter then all other parameters are ignored. (Blank indicates that AnnAGNPS will calculate values.)	"Actual_Trap_E fficiency_Clay", "Actual_Trap_E fficiency_Silt", "Actual_Trap_E fficiency_Sand", "Actual_Trap_E fficiency_Sm_A gg", "Actual_Trap_E fficiency_Lg_Ag g"	[nd]	Blank, or [0.0 to 1.0]	5F10	2	10-14
Fraction Trapped "TE-ps" — Fraction trapped for each AnnAGNPS particle size. (Blank indicates that AnnAGNPS will calculate values.)	"Fraction_Trap ped_Clay", "Fraction_Trap ped_Silt", "Fraction_Trap ped_Sand", "Fraction_Trap ped_Sm_Agg", "Fraction_Trap ped_Lg_Agg"	[nd]	Blank, or 0.00001 to 3.0	5F10	2	15-19
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	20
Go to Layout Matrix		Go to	Table of Conten	ts		

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

RUNOFF CURVE NUMBER DATA (BUILT-IN)

There are two internally defined curve number sets.

RCN ID	Curve "A"	Curve "B"	Curve "C"	Curve "D"			
"Default Crop CN"	72. 81. 88.						
"Default Non-Crop CN"	68. 79. 86. 89.						
No layout matrix available	Go to Table of Contents						

RUNOFF CURVE NUMBER DATA

Required

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				A	1	1-5
The following record repeats for th	e number of r	unoff curve	numbers.			-
Curve Number ID —Alphanumeric string ID for the specific cover (cover type, treatment and hydrologic condition) for Runoff Curve Number data.	"Curve_Numbe r_ID"			A40	2	1
Curve Number "A" —Runoff Curve Number for Soil Hydrologic Group "A"	"CN_A"		30.0 to 100.0	F10	2	2
Curve Number "B"—Runoff Curve Number for Soil Hydrologic Group "B"	"CN_B"		30.0 to 100.0	F10	2	3
Curve Number "C"—Runoff Curve Number for Soil Hydrologic Group "C"	"CN_C"		30.0 to 100.0	F10	2	4
Curve Number "D"—Runoff Curve Number for Soil Hydrologic Group "D"	"CN_D"		30.0 to 100.0	F10	2	5
<u>Go to Layout Matrix</u>		<u>Go to</u>	Table of Conten	<u>ts</u>		

RUSLE2 DATA

Optional unless referenced in other data sections

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-3
The following record repeats for the number of RUSLE2 records.						
RUSLE2 ID — Alphanumeric string identifying the RUSLE2 record.	"RUSLE2_ID"			A40	2	1
RUSLE2 Filename — Filename of the csv-formatted RUSLE2 erosion file to use as input	"RUSLE2_Filen ame"		-	A250	2	2
RUSLE2 Erosion Flag — True if RUSLE2 erosion values are to be used. If this flag is set to false then RUSLE1 erosion will calculated. Default = true.	"RUSLE2_Erosi on_Flag"		Blank, T or F	L1	2	3
Go to Layout Matrix	Go to Table of Contents					

SIMULATION PERIOD DATA Required

	unea					
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-17
The following reco	ord does not re	peat.				
Simulation Begin Date—Month, day, and year the watershed simulation begins.	"Simulation_Be gin_Month", "Simulation_Be gin_Day", "Simulation_Be gin_Vear"	mm dd yyyy	1 to 12 1 to 31 1 to 9999	12, 12, 14	2	1-3
Simulation End Date—Month, day, and year the watershed simulation ends. Leave blank for a single event simulation .	"Simulation_En d_Month", "Simulation_En d_Day", "Simulation_En d_Year"	mm dd yyyy	Blank, or 1 to 12 1 to 31 1 to 9999	12, 12, 14	2	4-6
Rainfall factor—Average annual RUSLE rainfall factor.	"Rainfall_Fctr"	{100 ft- ton-in / (acre-hr year)} [megajoule -mm / hectare-hr year]	{0.0 to 2000.0} [0.0 to 34100.0]	F10	2	7
10-yr EI —RUSLE energy intensity for 10 year frequency rainfall.	"10-Year_El"	{100 ft- ton-in / (acre-hr)} [megajoule -mm / hectare-hr]	{0.000006 to 2000.0} [0.0001 to 34100.0]	F10	2	8
EI Number —Energy Intensity (EI) distribution number identifying the EI distribution curve. For values > 400, the EI curve accounts for frozen soil effects such as in the Pacific Northwest Palouse region. EI distributions from RUSLE automatically entered for EI codes 1- 149. Optionally, the user may enter the EI distribution values through the EI PERCENTAGE DATA section. To account for frozen soil effects, the user must enter a code value > 400, and enter the EI distribution values in the EI PERCENTAGE DATA section.	"EI_Number"		1 to 500	I10	2	9
Irrigation Climate code —Code indicating the climate category to use for irrigation. Acceptable values are: $1 = arid$; $2 = humid$ (Blank is 2)	"Irrigation_Cli mate_Code"		Blank, 1 or 2	I10	2	10
Soil Moisture Steps —Number of soil moisture computation time steps within a day. (Blank defaults to 8 time steps.)	"Soil_Moisture_ Steps"		Blank, 1 to 24	I10	2	11

	Field	Units				
Description	Header for	{English}	Domain {English} [SI]	For mat	Record No.	Field No.
Annual K-factor code—Code indicating whether average annual	Kecord #1 "Annual_K_Fct	[51]	Blank, Y or N	A10	2	12
USLE soil losses are based on nomographs and volcanic soil	r_Code"		,	_		
equations or to use K factors provided with soil data. Acceptable						
values are: $\mathbf{X} = \mathbf{X}_{-1}$ $\mathbf{N} = \mathbf{N}_{-1}$ (Distribute for the factor \mathbf{X}_{-1})						
Y = Y es $N = No$ (Blank defaults to Y es).	"Variable K Fc		Plank V or N	A 10	2	12
average annual USLE soil loss throughout the year. Acceptable	tr_Code"		Dialik, I of N	AIU	Z	15
values are:						
Y = Yes $N = No$ (Blank defaults to Yes).						
Number of Initialization Years—Number of climate data years to	"Number_Init_		Blank, 0 to 100	I10	2	14
run for initializing variables prior to watershed simulation. The	Years"					
climate data used for initialization will be based on the record in the						
climate data file that corresponds to the years of initialization in the						
data in the climate file for initialization, then the option selected in						
the Initialization Method Code parameter will be used to complete						
the initialization record that is missing in the climate file. (Blank						
defaults to 2)						
Initialization Method Code—Code for initialization method; This	"Init_Method_C		Blank, 0 to 1	I10	2	15
is only used if the "Number of Initialization Years" parameter is	ode"					
populated in the SIMULATION PERIOD DATA section (where						
two initialization years are defined by default) and there are not this						
climate data input file						
ennate data input inc.						
0 = typical weather; precipitation monthly normals are calculated						
based on the entire climate data input file. Then, a determination is						
made as to the month and year of the actual climate data that had the						
smallest deviation from the normal. The actual climate data from						
that month and year is used for the corresponding month of						
initialization and repeated for each year of initialization.						
1 = re-use input weather: AnnAGNPS calculates a pseudo-year						
based on the simulation beginning year less the current year of						
initizalization. The actual climate data for the pseudo-year is used in						
its entirety for the current year of initialization. The pseudo-year						
decrements for each subsequent year of initialization. If initialization						
years requested exceeds the years of actual climate data supplied,						
then the fast year of actual chillate data is repeated as needed.						
(Blank defaults to 0)						
Winter Bouts—not currently used – Reserved.	"Winter_Bouts"			10	2	16
Input Units Code— Code identifying whether input is in English or	"Input_Units_C			I1	2	17
SI units. $0 = \text{English}, 1 = \text{SI}$ (Blank defaults to the input units code	oue					
in the AnnAGNPS ID data section)						
Co to Layout Matrix		Cot	Cable of Conton	to		

SOIL DATA Required

Field DescriptionField Header for Record #1Units {English} [SI]Domain For Record #1For Record #1Record Fi								
Field Header— Required unique field header for each field listed below.				А	1	1-12		
The following record repeats for the number of soils. The accompanying required soil layer data is specified in the SOIL LAYER DATA section.								
Soil ID —Soil Survey area ID combined with soil survey mapping unit symbol. (e.g., 013Mbac)	"Soil_ID"			A10	2	1		

The finance input_operations_ford f_2019 initial debit	Revision: December 4, 2019	File Name:	Input_Specifications_	v5.51_2019.xx.xx.docx
--	----------------------------	------------	-----------------------	-----------------------

	Field	Units				
Description	Header for	{English}	Domain {English} [SI]	For mat	Record No.	Field No.
	Record #1	[51]				
Hydrologic Soil Group —Soil Hydrologic group (see TR-55). Acceptable values are : A, B, C, D	"Hydrologic_Soi l_Group"		A, B, C, or D	A10	2	2
K-factor—USLE & RUSLE K factor for whole soil: Note this has already been adjusted to add the rock fragments > 2 mm.	"K_Factor"	{(ton ac hr)/ (100 ac ft tonf in)} [(met. ton hec hr) / (hec Mj mm)]	{0.0 to 1.0} [0.0 to .1317]	F10	2	3
Albedo—Solar radiation reflection from the bare soil surface.	"Albedo"	Radiation reflected / radiation incoming	Blank or 0.0 to 1.0	F10	2	4
Time to consolidation —Time for 95% of effect of disturbance to have disappeared due to consolidation.	"Time_to_Cons olidation"	years	0.0 to 100.0	F10	2	5
Impervious Depth —Depth to impervious layer in soil column. Blank defaults to a depth of 1000 [m]. Note: RUSLE assumes that residue incorporation cannot occur within a soil depth of 2 inches or less.	"Impervious_De pth"	{in} [mm]	Blank, or {2.01 to 393701.0} [50.9 to 10000000.0]	F10	2	6
Specific Gravity —Average specific gravity for the total mass in the soil column. (Blank defaults to 2.65)	"Specific_Gravit y"	mass-soil / mass-H ₂ O (nd)	Blank or 0.0 to 4.0	F10	2	7
Initial Soil Conditions ID —Alphanumeric string identifying the initial conditions record to be used for this soil. This is an optional parameter. This ID, if present, must have a matching ID and record set in the "Soil Initial Conditions Data" section.	"Initial_Soil_Co nditions_ID"			A10	2	8
If this parameter is left blank, a check is made to see if a global "Default Initial Soil Conditions ID" has been specified in the "Global IDs, Factors, and Flags Data" section. If so, the associated global default "Soil Initial Conditions Data" ID and record set will be used.						
Therefore, if this ID parameter is populated –or– is blank and a global "Default Initial Soil Conditions ID" has been specified in the "Global IDs, Factors, and Flags Data" section, then the initial conditions parameters specified in the associated "Soil Initial Conditions Data" record are used for those associated parameters in this Soil Data section that are blank. Otherwise, internally-defined values will be used as defaults.						
Soil Name—Full name for the soil	"Soil_Name"			A40	2	9
Soil Texture —Unabbreviated soil texture (e.g., clay loam) Number Soil Layers —Number of soil layers for this soil. At least one soil layer record is required for each soil. The soil layer information is specified in the SOIL LAYER DATA section.	"Soil_Texture" "Number_of_So il_Layers"		1 to 2147483647	A40 I10	2 2	10 11
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	12
Go to Layout Matrix		Go to	Table of Conten	ts		

SOIL INITIAL CONDITIONS DATA Optional

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-20

	Field	Units				
Description	Header for	{English}	Domain	For	Record	Field
	Record #1	[SI]	{English} [51]	mat	INO.	NO.
The following record repeats for the number of soil initial cond	itions records.	This data so	ection is optional a	and is	needed or	nly if
the global soil initial conditions defaults are not sufficient for in initialization. if initialization years are defined, or the first of	day of simulati	ion if there a	ditions are effective are no initialization	ve on n vear	the first u rs defined	ay of
Initial Soil Conditions ID —Alphanumeric string identifying the	"Initial_Soil_Co			A10	2	1
initial conditions record.	nditions_ID"					
Landuse Type ID—Alphanumeric string describing the landuse	"Landuse_Type			A10	2	2
type. Acceptable values are: "Cropland" or "Non-Crop" (Blank	_10					
Inorganic N—Initial soil inorganic nitrogen to be used for each cell.	"Inorganic N 1	nnm	Blank or	2F10	2	3-4
Two soil layers are used, first is the top 200 mm (\sim 8 in), second is	" for layer 1 and	ppm	0.0 to 100000.0	21 10	2	J- 1
the remaining soil profile.	"Inorganic_N_2 " for layer 2					
Blank = -999. This means that a determination of the default value						
will be made after the "Organic Matter Ratio" parameter is read in						
the Soil Data section. The default value will then be calculated as						
(Organic Matter Katto us determined in the Sou Data section 2700).						
Inorganic P —Initial soil inorganic phosphorus to be used for each	"Inorganic_P_1	ppm	Blank, or	2F10	2	5-6
cell. Two soil layers are used, first is the top 200 mm (~8 in),	" for layer 1 and "Inorganic P 2		0.0 to 100000.0			
second is the remaining soil profile.	" for layer 2					
Blank = -999. This means that a determination of the default value						
the Soil Data section. The default value will then be calculated as						
(Organic Matter Ratio as determined in the Soil Data section *						
7500).						
Soil Moisture— Initial soil moisture, in non-dimensional units [nd]	"Soil_Moisture_	ht-H ₂ O /	Blank, or	2F10	2	7-8
of available soil moisture, to be used for each cell. The non-	1" for layer 1 and	ht-soil	0.0 to 1.0, or >1.			
dimensional units is a ratio of the initial soil moisture divided by the	"Soil_Moisture_	layer				
available soil moisture. Available soil moisture is the soil moisture at field concerts less the soil moisture at the wilting point. Two soil	2" for layer 2	(na)				
lavers are used: the first laver is the top 200 [mm] (8 [in]), the						
second layer is called the bottom layer and is the remaining soil						
profile. Blank defaults to 0.5 [nd] which is the average soil moisture						
between the field capacity, which is 1.[nd], and the wilting point,						
which is 0.[nd]. Any value greater than "1." will default to the total						
Saturation which is the entitle porosity. (Diank defaults to 0.5)	"Organic Matte		Dlamk on	2510		0.10
cell Two soil layers are used, first is the top 200 mm (~8 in).	r_1" for layer 1	mass—org matter /	0.0 to 1.0	2110	۷	9-10
second is the remaining soil profile. (Blank defaults to 0.05 for each	and "Organic Matte	mass-soil	0.0 10 1.0			
layer)	r_2" for layer 2	(nd)				
Organic N—Initial soil organic nitrogen to be used for each cell.	"Organic_N_1"	ppm	Blank, or	2F10	2	11-12
Two soil layers are used, first is the top 200 mm (~8 in), second is	for layer 1 and "Organic N 2"		0.0 to 100000.0			
the remaining soil profile.	for layer 2					
Blank = -999. This means that a determination of the default value will be made after the "Organic Matter Ratio" parameter is read in						
the Soil Data section. The default value will then be calculated as				l l		
(Organic Matter Ratio as determined in the Soil Data section *						
90000).						l
Organic P —Initial soil organic phosphorus to be used for each cell.	"Organic_P_1"	ppm	Blank, or	2F10	2	13-14
Two soil layers are used first is top 8 in (200 mm), second is	"Organic_P_2"f		0.0 to 100000.0			
remaining soil profile. $P_{1} = 0.00$ This means that a determination of the default value.	or layer 2			l l		
will be made after the "Organic Matter Ratio" parameter is read in						
the Soil Data section. The default value will then be calculated as						
(Organic Matter Ratio as determined in the Soil Data section *						
15000).						
Surface Residue—Initial surface residue to be used for each cell.	"Surface_Resid	$\{lb / acre\}$	{0.0 to	F10	2	15
(Blank defaults to 0.0)	ue	[kg /	100000.0}			
Manning's n Initial condition overland flow Manning's n to be	"Mannings n"	hectare	0.005 to 1000.01	F10	2	16
used for each cell if not defined as part of an initial operations data	Mannings_n		0.003 10 1.0	1.10	2	10
for the first field management. (Blank defaults to 0.035)						

AnnAGNPS Input File Specifications

Revision: December 4, 2019	File Name: Inpr	ut_Specifications_v5.5	1_2019.xx.xx.docx
----------------------------	-----------------	------------------------	-------------------

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Snow Depth —Initial condition depth of snow on ground to be used for each cell. (Blank defaults to 0.0)	"Snow_Depth"	{in} [mm]	Blank or {0.0 to 1000.0} [0.0 to 25400.0]	F10	2	17
Snow Density —Initial condition snow density if any snow is to be on ground for each cell. Must have value if positive snow depth is indicated above. Leave blank if no snow depth is indicated above. (Blank defaults to 0.0)	"Snow_Density"	$\{lb / ft^3\}$ [Mg / m ³]	Blank, or {0.0006 to 62.4} [0.00001 to 1.0]	F10	2	18
Surface Constant —Initial condition surface condition constant to be used for each cell if not defined as part of an initial operations data for the first field management. (Blank defaults to 0.15 for "Cropland"; 0.30 for "Non-crop")	"Surface_Const ant"		Blank, or 0.0 to 1.0	F10	2	19
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	20
<u>Go to Layout Matrix</u>		<u>Go to</u>	o Table of Conten	<u>ts</u>		

SOIL LAYER DATA Required

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-24
The following record repeats for the number of soil la All layer data must be sequential and contiguous	yers for each s for each speci	soil specified fied soil in tl	l in the SOIL DAT he SOIL DATA so	FA sec ection.	tion.	
Soil ID —Soil ID of the soil for which this record applies. Must match a Soil ID specified in the SOIL DATA section.	"Soil_ID"			A10	2	1
Layer Number—Sequential and contiguous layer number for the applicable soil.	"Layer_Number "		1 to 2147483647	I10	2	2
Layer Depth—Depth from the soil surface to the bottom of the soil layer	"Layer_Depth"	{in} [mm]	{1.0 to 120.0} [1.0 to 3000.0]	F10	2	3
Bulk Density —Dry bulk density of soil layer (assumed to be the consolidated stage for cropland top layer).	"Bulk_Density"	$ \{ lb / ft^3 \} \\ [Mg / m^3] $	{6.25 to 168.0} [0.1 to 2.7]	F10	2	4
Clay Ratio —Clay mass ratio to sum total mass of mineral soil (sand, silt, clay) excluding rock for the soil layer.	"Clay_Ratio"	mass-clay/ mass- mineral soil (nd)	0.0 to 1.0	F10	2	5
Silt Ratio—Silt mass ratio to sum total mass of mineral soil (sand, silt, clay) excluding rock for the soil layer.	"Silt_Ratio"	mass-silt/ mass- mineral soil (nd)	0.0 to 1.0	F10	2	6
Sand Ratio—Sand mass ratio to sum total mass of mineral soil (sand, silt, clay) excluding rock for the soil layer.	"Sand_Ratio"	mass-sand / mass- mineral soil (nd)	0.0 to 1.0	F10	2	7
Rock Ratio —Ratio of rock fragment mass (fraction greater than 2 mm) to sum total mass of mineral soil (sand, silt, clay, & rock)in the soil layer. Blank defaults to 0.0	"Rock_Ratio"	mass— rock / mass— mineral soil (nd)	Blank or 0.0 to 1.0	F10	2	8

Revision: December 4, 2019	File Name: Input_Specifications_v5.51_2019.xx.xx.docx
----------------------------	---

	17.14	TI				
Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Very Fine Sand Ratio —Ratio of very fine sand (0.05 mm to 0.1mm) to the sum of total mass of mineral soil (sand, silt, clay) excluding rock in the soil layer. Blank defaults to 0.0	"Very_Fine_San d_Ratio"	mass-very fine sand / mass- mineral soil	Blank or 0.0 to 1.0	F10	2	9
CaCO3—Calcium carbonate content of soil layer.	"CaCO3_Conte nt"	wt CaCO3 / wt < 2mm soil (nd)	Blank or 0.0 to 1.0	F10	2	10
Saturated Conductivity —-Saturated hydraulic conductivity of the soil layer.	"Saturated_Con ductivity"	{in / hr} [mm / hr]	{0.0 to 10000.0} [0.0 to 254000.0]	F10	2	11
Field Capacity —Fraction of water volume at field capacity (300 kPa) to the soil volume in the soil layer. Based on whole soil (includes rock fragments).	"Field_Capacity "	ht-H ₂ O / ht-soil layer (nd)	wilting point fraction to 1.0	F10	2	12
Wilting Point —Fraction of water volume at wilting point (1500 kPa) to the soil volume in the soil layer. Based on whole soil (includes rock fragments).	"Wilting_Point"	ht-H ₂ O / ht-soil layer (nd)	0.0 to 1.0	F10	2	13
Volcanic code —Code indicting whether soil layer is from a volcanic parent material. $Y = volcanic$ $N = not volcanic (Blank defaults to No)$	"Volcanic_Code "		Blank, Y, or N	A10	2	14
Base Saturation —Base saturation of the soil layer for volcanic soils. This parameter is only used if the Volcanic code is set to Yes and the Annual K-factor code is set to Yes in the Simulation Period Data Section. The use of this parameter should only be considered for those soils that are similar to soils found in Hawaii. (See Equation 3-2 in the RUSLE Handbook, Renard et al., 1997)	"Base_Saturatio n"	%	Blank, 0.0 to 100.0	F10	2	15
Unstable Aggregate Ratio —Ratio of unstable aggregates (< .25 mm) to sum total mass of soil (sand, silt, clay, rock, & organic matter) in the soil layer. Leave blank if soil layer is not from a volcanic parent material	"Unstable_Aggr egate_Ratio"	mass- unstable agg / mass- soil (nd)	Blank or 0.0 to 1.0	F10	2	16
pH —Representation of the Hydrogen ion concentration (pH) for the soil layer. pH which is the logarithm of the reciprocal of the H concentration (g atoms /l) is a measure of acidity / alkalinity. Blank defaults to 6.5	"рН"		Blank, 1.0 to 14.0	F10	2	17
Organic Matter Ratio—Ratio of organic matter to the sum of the total mass of soil (sand, silt, clay, rock, & organic matter) in the soil layer. Blank defaults to 0.05 if "Initial Soil Conditions ID" is not specified for this soil and "Default Initial Soil Conditions ID" is not specified in the "Global IDs, Factors, and Flags Data" section. Otherwise, blank will default to the "Organic Matter" in the "Soil Initial Conditions Data" record referenced by this soil or by the "Global IDs, Factors, and Flags Data" section.	"Organic_Matte r_Ratio"	mass-org matter / mass-soil (nd)	Blank, 0.0 to 1.0	F10	2	18
Organic N Ratio —Ratio of initial amount of organic nitrogen in the soil layer at the start of the simulation. Blanks defaults to (Organic Matter Ratio * 90000) if "Initial Soil Conditions ID" is not specified for this soil and "Default Initial Soil Conditions ID" is not specified in the "Global IDs, Factors, and Flags Data" section. Otherwise, blank will default to the "Organic N" in the "Soil Initial Conditions Data" record referenced by this soil or by the "Default Initial Soil Conditions ID" in the "Global IDs, Factors, and Flags Data" section.	"Organic_N_Ra tio"	ppm	Blank, 0.0 to 100000.0	F10	2	19

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Inorganic N Ratio —Ratio of initial amount of inorganic nitrogen in the soil layer at the start of the simulation. Blanks defaults to (Organic Matter Ratio * 2700) if "Initial Soil Conditions ID" is not specified for this soil and "Default Initial Soil Conditions ID" is not specified in the "Global IDs, Factors, and Flags Data" section. Otherwise, blank will default to the "Inorganic N" in the "Soil Initial Conditions Data" record referenced by this soil or by the "Default Initial Soil Conditions ID" in the "Global IDs, Factors, and Flags Data" section.	"Inorganic_N_R atio"	ppm	Blank, 0.0 to 100000.0	F10	2	20
Organic P Ratio —Ratio of initial amount of organic phosphorus in the soil layer at the start of the simulation. Blanks defaults to (Organic Matter Ratio * 15000) if "Initial Soil Conditions ID" is not specified for this soil and "Default Initial Soil Conditions ID" is not specified in the "Global IDs, Factors, and Flags Data" section. Otherwise, blank will default to the "Inorganic N" in the "Soil Initial Conditions Data" record as referenced by this soil or by the "Default Initial Soil Conditions ID" in the "Global IDs, Factors, and Flags Data" section.	"Organic_P_Rat io"	ppm	Blank, 0.0 to 10000.0	F10	2	21
Inorganic P Ratio —Ratio of initial amount of inorganic phosphorus in the soil layer at the start of the simulation. Blanks defaults to (Organic Matter Ratio * 7500) if "Initial Soil Conditions ID" is not specified for this soil and "Default Initial Soil Conditions ID" is not specified in the "Global IDs, Factors, and Flags Data" section. Otherwise, blank will default to the "Inorganic P" in the "Soil Initial Conditions Data" record referenced by this soil or by the "Default Initial Soil Conditions ID" in the "Global IDs, Factors, and Flags Data" section.	"Inorganic_P_R atio"	ppm	Blank, 0.0 to 10000.0	F10	2	22
Soil Structure code—Code indicating average aggregate soil structure size for soil layer. Acceptable values are: 1 = very fine granular (< 1mm) 2 = fine granular (1 to 2mm) 3 = medium or coarse granular (2 to 5mm) 4 = blocky, platey or massive (> 5mm) Blank defaults to 3.	"Soil_Structure _Code"		Blank, 1 to 4	I10	2	23
Input Units Code— Code identifying whether input is in English or SI units. 0 = English ,1 = SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"	Cott	Table of Contan	Il	2	24

STRIP CROP DATA

Optional unless referenced in Management Schedule Data

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.	
Field Header— Required unique field header for each field listed below.				А	1	1-3	
The following record repeats for	or the number	of strip croj	o sets.				
Strip Crop ID —Alphanumeric string identifying the Strip Crop Data.	"Strip_Crop_ID "			A10	2	1	
P Factor —The RUSLE conservation support practice associated with the entire impact of strip crop support practices applied on the entire field on the average annual erosion rate associated with the Strip Crop ID. Blank defaults to 1.0.	"P_Factor"	[nd]	Blank, 0. to 1.	F10	2	2	
Sediment Delivery Ratio —The fraction of the gross soil loss from all of the strip crops in the field to the amount of sediment leaving the strips at the bottom of the slope in the field. Blank defaults to 1.0.	"Delivery_Ratio "	[nd]	Blank, 0. to 1.	F10	2	3	
Go to Layout Matrix	Go to Table of Contents						

TILE DRAIN DATA

Optional unless referenced in Management Field Data

Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.					
Field Header— Required unique field header for each field listed below.				А	1	1-4					
The following line repeats for the number of tile drain sets (specified above).											
Tile Drain ID —Alphanumeric string identifying the Tile Drainage Scheme.	"Tile_Drain_ID "			A10	2	1					
Drain Rate —Daily reduction in height of water table. If entered by the user, this Value is used for all tile drainage calculations. Default value is 0.5 inches or 12.7 mm	"Drain_Rate"	{in / day} [mm / day]	0.0 to 999999.9	I10	2	2					
Invert Depth —Depth of installed tile drain invert below the surface. Default value is 800 [mm] or 31.5 [in]	"Invert_Depth"	{in} [mm]	0.0 to 9999999.9	F10	2	3					
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	4					
<u>Go to Layout Matrix</u>	Go to Table of Contents										

WATERSHED DATA Required

Keq	uncu					
Description	Field Header for Record #1	Units	Domain	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-5
The following reco	ord does not re	peat.	-	-	_	-
Watershed Name—Name of the watershed (Optional).	"Wshd_Name"		Blank or name up to 80 characters	A80	2	1
Watershed Description—Description of the watershed (Optional).	"Wshd_Descript ion"		Blank or description up to 80 characters	A80	3	2
Watershed Location—Location of the watershed (Optional).	"Wshd_Locatio n"		Blank or location up to 60 characters	A60	4	3
Latitude—Latitude for centroid of watershed. Latitude is expressed as "+" for North and "-" for South. (This is only used for reporting purposes; not used in any calculations therefore blank is acceptable and a "-999" will be reported in applicable verification output files)	"Latitude"	decimal °	Blank or -90. To 90.	F10	4	4
Longitude —Longitude for centroid of watershed. Longitude is expressed as "+" for East and "-" for West. (This is only used for reporting purposes; not used in any calculations therefore blank is acceptable and a "-999" will be reported in applicable verification output files)	"Longitude"	decimal °	Blank or -180. To 180.	F10	4	5
Go to Lavout Matrix		Go t	o Table of Conten	ts		

WETLAND DATA

Optional

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.			
Field Header— Required unique field header for each field listed				Α	1	1-17			
below.	<u> </u>	1'	<u> </u> '	<u> </u>	<u> </u>	1			
The following record repeats for the number of wetland records.									

AnnAGNPS Input File Specifications

		-	
Revision: December 4, 2019	File Name:	Input_Specifications_	_v5.51_2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Wetland ID—Alphanumeric string identifying the wetland record.	"Wetland_ID"			A10	2	1
Reach ID —Alphanumeric string identifying the reach that contains the wetland. Must be the same as a reach ID in the REACH DATA section. Note: AnnAGNPS currently only allows one wetland per reach.	"Reach_ID"			A10	2	2
Wetland Area—The area of the wetland. (Blank defaults to 2% of the drainage area at the upstream end of the reach. The maximum value allowed is less than or equal to the drainage area at the upstream end of the reach.)	"Wetland_Area "	{acre} [hectare]	Blank, or >0.0 to 9999999999.	F10	2	3
Initial Water Depth —Initial water depth in the wetland. (Blank defaults to 0.0)	"Initial_Water_ Depth"	{in} [mm]	Blank, or {0.0 to 1968.5} [0.0 to 50000.0]	F10	2	4
Minimum Water Depth—Minimum water depth in the wetland. (Blank defaults to 0.0)	"Min_Water_De pth"	{in} [mm]	Blank, or {0.0 to 196.8} [0.0 to 5000.0]	F10	2	5
Maximum Water Depth—Maximum water depth in the wetland. (Blank defaults to 1000.0; this value cannot be less than the Minimum Water Depth.)	"Max_Water_D epth"	{in} [mm]	Blank, or {>0.0 to 1968.5} [>0.0 to 50000.0]	F10	2	6
Water Temperature—Wetland water temperature. (Blank defaults to AnnAGNPS average air temperature.)	"Water_Temper ature"	{°F} [°C]	Blank, or {32.0 to 104.0} [0.0 to 40.0]	F10	2	7
Potential Daily Infiltration —Constant daily infiltration rate applied after water ponds in the wetland. (Blank defaults to; 1. First cell encountered in the sorted list of cell ids that has the same reach id as that of the wetland; or 2. Calculated watershed average daily infiltration value.)	"Potential_Daily _Infiltration"	{in/day} [mm/day]	Blank, or {0.0 to 100.0} [0.0 to 2540.0]	F10	2	8
Weir Coefficient—Weir coefficient. (Blank defaults to 2.0)	"Weir_Coef"	(nd)	Blank, or >0.0 to 10.0	F10	2	9
Weir Width—Width of weir opening. (Blank defaults to 1.0)	"Weir_Width"	{ft} [m]	Blank, or {0.03 to 328083} [0.01 to 100000.0]	F10	2	10
Weir Height—Height of the weir. (Blank defaults to 1.0)	"Weir_Height"	{ft} [m]	Blank, or {0.00328 to 32.8} [0.001 to 10.0]	F10	2	11
Soluble N Concentration —Initial concentration of soluble nitrogen. (Blank defaults to 0.0)	"Soluble_N_Con c"	{ppm} [mg/L]	Blank, or 0.0 to 100.0	F10	2	12
Nitrate-N Loss Rate—Nitrate-N loss rate. (Blank defaults to AnnAGNPS internally calculated value.)	"Nitrate- N_Loss_Rate"	${lb/ft^2/day} } [g/m^2/day]$	Blank, or >0.0 to 100.0	F10	2	13
Nitrate-N Loss Rate Coefficient —The area-based first order loss rate coefficient for nitrate-N at 20 °C (Blank defaults to 0.15)	"Nitrate- N_Loss_Rate_C oef"	{ft/day} [m/day]	Blank, or 0.05 to 0.5	F10	2	14
Temperature Coefficient —The temperature coefficient for nitrate- N loss. (Blank defaults to 1.09)	"Temperature_ Coef"	(nd)	Blank, or 0.1 to 2.5	F10	2	15
Weir Exponent—Weir coefficient. (Blank defaults to 1.5)	"Weir_Exp"	(nd)	Blank, or 0.0 to 100.0	F10	2	16
Input Units Code — Code identifying whether input is in English or SI units. $0 = \text{English}$, $1 = \text{SI}$ (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"			I1	2	17
Go to Layout Matrix		Go to	Table of Content	ts		

Input File Layout Matrix

ANNAGNPS ID

Record	Field 1	Field 2	Field 3	Field 4	Field 5				
1	Version	Input_Units	Output_Units	CCHE1D_Output_ Units	Screen_Output_U nits				
2	5.20								
Go to Data Section Description						<u>Go to</u>	Table of Conter	<u>its</u>	

AQUACULTURE POND DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Pond_ID	Cell_ID	Pond_Area	Pond_Depth	Seepage_Rat	e Sediment_Delivery _Ratio	Relative_Rotation_ Year	Mgmt_Schd_ID	OC_Calib_Fctr	N_Calib_Fctr
2										
Record	Field 11	Field 12	Field 13							
1	P_Calib_Fctr	Erosion_Calib_Fct r	Input_Units_Code							
2										
Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>		

AQUACULTURE POND MANAGEMENT SCHEDULE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Pond_Mgmt_Sc hd_ID	Month	Day	Year	Water_Operation_ Code	Aquaculture_ID	Crop_ID	Planting_Type_Co de	Gate_Status	Max_Pool_Depth
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Min_Pool_Dept h	Fill/Release_Vol	Fill/Drain_Time	Fill/Release_Rate	Fill/Drain_All_Co de	Total_Sed_Conc	Clay_Content	Silt_Content	Total_N	Dissolved_N
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
Record	Field 21 Total_P	Field 22 Dissolved_P	Field 23 Num_Pest_Apps	Field 24 Season_Adjust_Co nc	Field 25 Sed_Conc_Winter	Field 26 Total_N_Winter	Field 27 Dissolved_N_Wint er	Field 28 Total_P_Winter	Field 29 Dissolved_P_Wint er	Field 30 Sed_Conc_Spring
Record 1 2	Field 21 Total_P	Field 22 Dissolved_P	Field 23	Field 24 Season_Adjust_Co nc	Field 25 Sed_Conc_Winter	Field 26 Total_N_Winter	Field 27 Dissolved_N_Wint er	Field 28 Total_P_Winter	Field 29 Dissolved_P_Wint er	Field 30 Sed_Conc_Spring
Record 1 2 Record	Field 21 Total_P Field 31	Field 22 Dissolved_P Field 32	Field 23 Num_Pest_Apps Field 33	Field 24 Season_Adjust_Co nc Field 34	Field 25 Sed_Conc_Winter Field 35	Field 26 Total_N_Winter Field 36	Field 27 Dissolved_N_Wint er Field 37	Field 28 Total_P_Winter Field 38	Field 29 Dissolved_P_Wint er Field 39	Field 30 Sed_Conc_Spring Field 40
Record 1 2 Record 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Field 21 Total_P Field 31 Total_N_Spring	Field 22 Dissolved_P Field 32 Dissolved_N_Sprin g	Field 23 Num_Pest_Apps Field 33 Total_P_Spring	Field 24 Season_Adjust_Co nc Field 34 Dissolved_P_Sprin g	Field 25 Sed_Conc_Winter Field 35 Sed_Conc_Summe r	Field 26 Total_N_Winter Field 36 Total_N_Summer	Field 27 Dissolved_N_Wint er Field 37 Dissolved_N_Sum mer	Field 28 Total_P_Winter Field 38 Total_P_Summer	Field 29 Dissolved_P_Wint er Field 39 Dissolved_P_Sum mer	Field 30 Sed_Conc_Spring Field 40 Sed_Conc_Autum n

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

Record	Field 41	Field 42	Field 43	Field 44	Field 45					
1	Total_N_Autu mn	Dissolved_N_Autu mn	Total_P_Autumn	Dissolved_P_Autu mn	Input_Units_Code	e				
2										
		<u>Go to Data Se</u>	ection Description				<u>Go to</u>	Table of Content	<u>s</u>	

CELL DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Cell_ID	Soil_ID	Mgmt_Field_ID	Reach_ID	Reach_Location_C ode	Cell_Area	Time_of_Conc	Avg_Elevation	RCN_Calib_Fctr	Avg_Land_Slope
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Aspect	RUSLE_ls_Fctr	RCN_Rtn_Calib_ Fctr	Secondary_Climat e_File_ID	Sheet_Flow_Mann ings_n	Geology_ID	Conc_Flow_Slope	Conc_Flow_Lengt h	Hydraulic_Geom_ ID	Conc_Flow_Hydra ulic_Depth
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	Conc_Flow_Ma nnings_n	Sheet_Flow_Slope	Sheet_Flow_Lengt h	Shallow_Conc_Flo w_Slope	Shallow_Conc_Flo w_Length	Delivery_Ratio	Constant_USLE_ C_Fctr	Constant_USLE_P _Fctr	All_OC_Calib_Fct r	All_N_Calib_Fctr
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35					
1	All_P_Calib_Fc tr	Sheet_and_Rill_Er osion_Calib_Fctr	Gullies_Erosion_C alib_Fctr	RUSLE2_ID	Input_Units_Code					
2										
		Go to Data Se	ection Descriptio	<u>n</u>			<u>Go to</u>	Table of Conter	<u>its</u>	

CLASSIC GULLY DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Gully_ID	Cell_ID	Reach_ID	Soil_ID	Cell_Drainage_Ar ea	Reach_Drainage_ Area	Headcut_Depth	Erosion_Coef	Erosion_exp	Delivery_Ratio
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Mgmt_Field_ID	Cell_Drainage_Su barea	Load_Calib_Fctr	Rainfall/Runoff_In dicator	Units_Indicator	Gully_Location_C ode	OC_Calib_Fctr	N_Calib_Fctr	P_Calib_Fctr	Erosion_Calib_Fct r
2										
Record	Field 21									
1	Input_Units_Co de									
2										

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx Revision: December 4, 2019

Go to Data Section Description	Go to Table of Contents
--------------------------------	-------------------------

CONTOUR DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6				
1	Contour_ID	Ridge_Height_Cod e	Furrow_Slope	Disturbed_Cover	Consolated_Cover	Input_Units_Code				
2										
	Go to Data Section Description						<u>Go to</u>	Table of Conter	<u>nts</u>	

CROP DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Crop_ID	Yield_Units_Harve sted	Residue_Mass_Rat io	Surface_Decomp	Sub- surface_Decomp	USLE_C_Fctr	Moisture_Depletio n	Residue_Adjust_A mt	Crop_Residue_30 %	Crop_Residue_60 %
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Crop_Residue_ 90%	Annual_Crop_Cod e	Legume_Code	Senescence_Code	Yield_Unit_Name	Yield_Unit_Mass	Harvest_C- N_Ratio	Pre-Harvest_C- N_Ratio	Harvest_Water	N_Uptake
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	P_Uptake	Harvest_C- P_Ratio	Pre-Harvest_C- P_Ratio	Growth_Time_Ini	Growth_Time_Dev	Growth_Time_Ma t	Growth_Time_Sen	Growth_N_Uptake _Ini	Growth_N_Uptake _Dev	Growth_N_Uptake _Mat
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	Growth_N_Upt ake_Sen	Growth_P_Uptake _Ini	Growth_P_Uptake _Dev	Growth_P_Uptake _Mat	Growth_P_Uptake _Sen	Basal_Crop_Coef_ Ini	Basal_Crop_Coef_ Dev	Basal_Crop_Coef_ Mid	Basal_Crop_Coef_ End	Basal_Crop_Coef_ Climate_Adjust
2										
Record	Field 41									
1	Input_Units_Co de		-							
2										
		<u>Go to Data Se</u>	ection Description				<u>Go to</u>	Table of Content	<u>s</u>	

CROP GROWTH DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5					
1	Crop_Growth_ ID	Root_Mass	Canopy_Cover	Rain_Fall_Height	Input_Units_Co	de				
2										
		<u>Go to Data S</u>	ection Description				<u>Go to</u>	Table of Content	<u>s</u>	

EI PERCENTAGE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	"EI_Pct_01"	"EI_Pct_02"	"EI_Pct_03"	"EI_Pct_04"	"EI_Pct_05"	"EI_Pct_06"	"EI_Pct_07"	"EI_Pct_08"	"EI_Pct_09"	"EI_Pct_10"
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
Record 1	Field 11 "EI_Pct_11"	Field 12 "EI_Pct_12"	Field 13 "EI_Pct_13"	Field 14 "EI_Pct_14"	Field 15 "EI_Pct_15"	Field 16 "EI_Pct_16"	Field 17 "EI_Pct_17"	Field 18 "EI_Pct_18"	Field 19 "EI_Pct_19"	Field 20 "EI_Pct_20"

Record	Field 21	Field 22	Field 23	Field 24					
1	"EI_Pct_21"	"EI_Pct_22"	"EI_Pct_23"	"EI_Pct_24"					
2									
		Go to Data S	ection Description			<u>Go to</u>	Table of Content	<u>s</u>	

EPHEMERAL GULLY DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Gully_ID	Cell_ID	Reach_ID	Soil_ID	Drainage_Area_to _Mouth	Local_Drainage_A rea	Gully_Slope	Critical_Sheer_Str ess	Gully_Location_C ode	Mgmt_Field_ID
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Erosion_Depth	Cells_Drainage_Su bcell	Hydraulic_Geomet ry_ID	Width_Nachtergae le	Width_Hydraulic_ Geometry	Width_Non- submerging_Tailw ater	Width_Woodward s_Equilibrium	Width_Woodward s_Ultimate	Width_Wells_Eq.9	Delivery_Ratio
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
Record	Field 21 Mannings_n	Field 22 Replant_Period	Field 23 OC_Calib_Fctr	Field 24 N_Calib_Fctr	Field 25 P_Calib_Fctr	Field 26 Erosion_Calib_Fct r	Field 27 Headcut_Migratio n Barrier	Field 28 Headcut_Dtach/Er od_Coef_a	Field 29 Headcut_Dtach/Er od Exp Coef b	Field 30 Max_Trapping_Ef ficiency
Record 1 2	Field 21 Mannings_n	Field 22 Replant_Period	Field 23	Field 24	Field 25 P_Calib_Fctr	Field 26 Erosion_Calib_Fct r	Field 27 Headcut_Migratio n Barrier	Field 28 Headcut_Dtach/Er od Coef a	Field 29 Headcut_Dtach/Er od Exp Coef b	Field 30 Max_Trapping_Ef ficiency
Record 1 2 Record	Field 21 Mannings_n Field 31	Field 22 Replant_Period Field 32	Field 23 OC_Calib_Fctr Field 33	Field 24 N_Calib_Fctr Field 34	Field 25 P_Calib_Fctr Field 35	Field 26 Erosion_Calib_Fct r	Field 27 Headcut_Migratio n Barrier	Field 28 Headcut_Dtach/Er od Coef a	Field 29 Headcut_Dtach/Er od Exp Coef b	Field 30 Max_Trapping_Ef ficiency
Record 1 2 Record 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Field 21 Mannings_n Field 31 Width_Wells'_ Eq.8	Field 22 Replant_Period Field 32 Width_Reserved_i	Field 23 OC_Calib_Fctr Field 33 Width_Reserved_j	Field 24 N_Calib_Fctr Field 34 Width_Reserved_k	Field 25 P_Calib_Fctr Field 35 Input_Units_Code	Field 26 Erosion_Calib_Fct r	Field 27 Headcut_Migratio n Barrier	Field 28 Headcut_Dtach/Er od Coef a	Field 29 Headcut_Dtach/Er od Exp Coef b	Field 30 Max_Trapping_Ef ficiency
Record 1 2 Record 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Field 21 Mannings_n Field 31 Width_Wells'_ Eq.8	Field 22 Replant_Period Field 32 Width_Reserved_i	Field 23 OC_Calib_Fctr Field 33 Width_Reserved_j	Field 24 N_Calib_Fctr Field 34 Width_Reserved_k	Field 25 P_Calib_Fctr Field 35 Input_Units_Code	Field 26 Erosion_Calib_Fct r	Field 27 Headcut_Migratio n Barrier	Field 28 Headcut_Dtach/Er od Coef a	Field 29 Headcut_Dtach/Er od Exp Coef b	Field 30 Max_Trapping_Ef ficiency

FEEDLOT DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Feedlot_ID	Feedlot_Mgmt_ID	Open_Area	Paved_Ratio	Roof_Area	Upslope_Area	Initial_N	Initial_P	Initial_OC	Delta_N
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Delta_P	Delta_OC	Max_N	Max_P	Max_OC	Pack_N	Pack_P	Pack_OC	OC_Calib_Fctr	N_Calib_Fctr
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25					
1	P_Calib_Fctr	Erosion_Calib_Fct r	Cell_ID	Cell_Buffer_Lengt h	Input_Units_Code					
2										
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>s</u>	

FEEDLOT MANAGEMENT DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	Mgmt_ID	Month	Day	Year	Pack_Remove_Ra io	t Pack_Start_N	Pack_Start_P	Pack_Start_OC	Pack_Change_N	Pack_Change_P	
2											
Record	Field 11	Field 12									
1	Pack_Change_ OC	Input_Units_Code									
2											
		Go to Data Se	ection Description			Go to Table of Contents					

FERTILIZER APPLICATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6				
1	Application_ID	Name_ID	Application_Rate	Depth	Mixing_Code	Input_Units_Code				
2										
		<u>Go to Data S</u>	ection Description				<u>Go to</u>	Table of Content	<u>ts</u>	

FERTILIZER REFERENCE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Fiel	d 6	Field 7	Field 8	Field 9	Field 10	
1	Reference_ID	Nitrite	Nitrate	Inorganic_N	Organic_N	Amm	onia	Mineral_Ammonia	Elemental_P	Soluble_P	Inorganic_P	
2												
Record	Field 11	Field 12	Field 13									
1	Organic_P	Organic_Matter	Consistency_Code									
2												
	Go to Data Section Description						Go to Table of Contents					

FIELD POND DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10			
1	Pond_ID	Cell_ID	Pond_Area	Number_of_Rotati on_Years	Number_of_Gate_ Operations	Delivery_Ratio	Volume_of_Releas e_Water	Drain_Time	Release_Rate	Sediment_Conc			
2													
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20			
1	Clay_Content	Silt_Content	N_Conc	P_Conc	OC_Conc	Pesticide_Referenc e_ID	Pesticide_Conc	OC_Calib_Fctr	N_Calib_Fctr	P_Calib_Fctr			
2													
Record	Field 21	Field 22											
1	Erosion_Calib_ Fctr	Input_Units_Code											
2													
	Go to Data Section Description						Go to Table of Contents						

FIELD POND OPERATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5				
1	Pond_ID	Open/Close_Gate_ Action	Open/Close_Rotati on_Month	Open/Close_Rotati on_Day	Open/Close_Rotati on_Year				
2									
Go to Data Section Description						<u>Go to</u>	Table of Content	<u>ts</u>	

GEOLOGY DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	Geology_ID	Delay_Time	Water_Table	Aquifer_Sat_Hyd_ Conduct	Vadose_Sat_Hy Conduct	/d_ Porosity	Field_Capacity	Specific_Yield	Thickness	Soluble_N	
2											
Record	Field 11	Field 12									
1	Soluble_P	Input_Units_Code									
2											
Go to Data Section Description						Go to Table of Contents					

GLOBAL ERROR AND WARNING LIMITS DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5					
1	Keyword_ID	Warning_Min	Warning_Max	Error_Min	Error_Max	í.				
2										
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>s</u>	
GLOBAL IDS, FACTORS, AN	ND FLAGS DATA									
--------------------------------	---------------									
--------------------------------	---------------									

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	Hdct_Detachme nt Coef a	Hdct_Detachment Exp_Coef_b	Urban_Repair_Mo nth	Urban_Repair_Da v	Urban_Repair_Ye ar	Cropland_Repair_ Month	Cropland_Repair_ Dav	Cropland_Repair_ Year	Forest_Repair_Mo nth	Forest_Repair_Da v	
2				2							
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20	
1	Forest_Repair_ Year	Pasture_Repair_M onth	Pasture_Repair_D ay	Pasture_Repair_Y ear	Rangeland_Repair Month	Rangeland_Repair Day	Rangeland_Repair Year	Hdct_Erodibility_ Coef a	Hdct_Erodibility_ Exp_Coef_b	Width_Nachtergae le	
2											
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30	
1	Width_Hydraul ic_Geometry	Width_Non- submerging_Tailw ater	Width_Woodward s_Equilibrium	Width_Woodward s_Ultimate	Width_Wells_Eq.9	Erosion_Vrfy	Hydrograph_Vrfy	Nickpoint_Vrfy	Repair_Dates_Vrf y	Sed_Yield_to_Gull y_Mouth_Vrfy	
2											
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40	
1	Sed_Yield_to_R cvg_Reach_Vrf y	Min_Interception_ Evaporation	Max_Interception_ Evaporation	Detention_Coef_a	Detention_Coef_b	RCN_Convergence _Tolerance	RCN_Max_Iterati ons	Avbl_Soil_Moist_ Ratio_AMC_II	Max_Avbl_Sed_C onc_for_Sht_Flw	Max_Avbl_Sed_C onc_for_Conc_Flw	
2											
Record	Field 41	Field 42	Field 43	Field 44	Field 45	Field 46	Field 47	Field 48	Field 49	Field 50	
1	AA_Unit_Area Baseflow	RCN_Calib_Only	Calculate_Baseflo w	FAO_ET_Enhance ment	Basal_Crop_Coef_ Climate_Adjust	Wshd_Storm_Typ e_ID	Dflt_Geology_ID	Dflt_Hydraulic_Ge om ID	Dflt_Init_Soil_Con ditions_ID	Dflt_Crop_RCN_I D	
2											
Record	Field 51	Field 52	Field 53	Field 54	Field 55	Field 56	Field 57	Field 58	Field 59		
1	Dflt_Non- Crop_RCN_ID	Width_Wells'_Eq. 8	Width_Reserved_i	Width_Reserved_j	Width_Reserved_k	Critical_Shear_Str ess	RUSLE2_Flag	Dflt_RUSLE2_ID	Input_Units_Code		
2											
	Go to Data Section Description						Go to Table of Contents				

HYDRAULIC GEOMETRY DATA (USER DEFINED)

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	
1	Hydraulic_Geo m_ID	Channel_Length_ Coef	Channel_Length_ Exp	Channel_Width_C oef	Channel_Width_E xp	Channel_Depth_C oef	Channel_Depth_E xp	Valley_Width_Coe f	Valley_Width_Exp	
2										
Go to Data Section Description						Go to Table of Contents				

IMPOUNDMENT DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Impoundment_ ID	Infiltration	Seepage	Permanent_Pool_ Depth	Volume_Coef	Volume_Exp	Discharge_Coef	Discharge_Exp	Sed_Clean_Out_D epth	Sed_Clean_Out_Y ear
2										
Record	Field 11	Field 12								
1	Reach_ID	Input_Units_Code								
2										
		<u>Go to Data Se</u>	ection Description			Go to Table of Contents				

IRRIGATION APPLICATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Application_ID	Season_End_Mont h	Season_End_Day	Season_End_Year	Method_Code	Cycle_Duration	Amount_Lost	Application_Rate	Tailwater_Recover y	Depletion_Lower_ Limit
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18		
1	Application_A mount	Area_Fraction	Interval_Number	Interval_Days	Chemical_Multiple	Sediment_Rate	Depletion_Upper_ Limit	Input_Units_Code		
2										
		<u>Go to Data Se</u>	ection Description			Go to Table of Contents				

MANAGEMENT FIELD DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Field_ID	Landuse_Type_ID	Mgmt_Schd_ID	Greg_Yr_for_1 st _ Yr_of_Rotation	Percent_Rock_Cov er	Interrill_Erosion_ Code	Random_Roughne ss	Terrace_Horizonta l_Distance	Terrace_Grade	Tile_Drain_ID
2										
Record	Field 11									
1	Input_Units_Co de									
2										
		Go to Data Section Description					Go to Table of Contents			

MANAGEMENT OPERATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Mgmt_Operatio n_ID	Effect_Code_01	Effect_Code_02	Effect_Code_03	Effect_Code_04	Effect_Code_05	Residue_Cover_Re maining	Residue_Weight_R emaining	Area_Disturbed	Initial_Random_R oughness
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	
1	Final_Random_ Roughness	Operation_Tillage _Depth	Added_Surface_R esidue	Surface_Decomp	Subsurface_Decom p	Surface_Residue_3 0%	Surface_Residue_6 0%	Surface_Residue_9 0%	Input_Units_Code	
2										
	Go to Data Section Description					Go to Table of Contents				

MANAGEMENT SCHEDULE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Mgmt_Schd_ID	Event_Month	Event_Day	Event_Year	Contour_ID	New_Crop_ID	Strip_Crop_ID	New_Non- Crop_ID	Curve_Number_I D	Post_Event_Manni ngs_n
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Post_Event_Sur face_Constant	Operation_Residu e_Change	Fertilizer_Applicat ion_ID	Irrigation_Applica tion_ID	Mgmt_Operation_ ID	Tile_Drain_Contro lled_Status	Tile_Drain_Contro lled_Depth	Input_Units_Code	Pest_App_ID_1	Pest_App_ID_2
2										
Record	Field 21	Field 22	Field 23							
1	Pest_App_ID_3	Pest_App_ID_4	Pest_App_ID_5							
2										
	Go to Data Section Description					Go to Table of Contents				

NON-CROP DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Non-Crop_ID	Non- Crop_Description	Annual_Root_Mas s	Annual_Cover_Ra tio	Annual_Rain_Fall _Height	Surface_Cover_Re sidue	USLE_C-Fctr	Basal_Crop_Coef_ Mid	Growing_Season_ Start_Month	Growing_Season_ Start_Day
2										
Record	Field 11	Field 12	Field 13	Field 14						
1	Growing_Seaso n_End_Month	Growing_Season_ End_Day	Basal_Crop_Coef_ Climate_Adjust	Input_Units_Code						
2										
	Go to Data Section Description					Go to Table of Contents				

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Glbl_All_V3_cs v	Glbl_All_V3_dpp	Glbl_All_V3_npt	Glbl_All_V3_sim	Glbl_All_V3_txt	Log_to_File	Log_to_Screen	Warning_File	V1/2_Output_Files	Reserved
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Glbl_All_Cells	Glbl_All_Feedlots	Glbl_All_Fld_Pon ds	Glbl_All_Gullies	Glbl_All_Pt_Srcs	Glbl_All_Reaches	Glbl_All_Impound	Reserved	Glbl_All_AA_Nutr	Glbl_All_AA_Pest
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	Reserved	Reserved	Glbl_All_AA_Sed	Glbl_All_AA_Wtr	Glbl_All_EV_Nutr	Glbl_All_EV_Pest	Glbl_All_EV_Sed	Glbl_All_EV_Wtr	Reserved	Reserved
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	Glbl_All_V2/3_ Mass	Glbl_All_V2/3_Rat io	Glbl_All_V2/3_UA	Reserved	V2_Concepts	Reserved	V2_AA	V2_EV	V1_AA	V1_EV
2										
	Go to Data Section Description					Go to Table of Contents				

OUTPUT OPTIONS DATA – GLOBAL

OUTPUT OPTIONS DATA – AA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	AA_Feedlots_M ass	AA_Feedlots_Rati 0	AA_Feedlots_UA	AA_Fld_Ponds_M ass	AA_Fld_Ponds_Ra tio	AA_Fld_Ponds_U A	AA_Gullies_Erosio n	AA_Gullies_Sedim ent (Reserved)	AA_Gullies_Nutrie nts (Reserved)	AA_N_Ld_Mass	
2											
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20	
1	AA_N_Ld_Rati 0	AA_N_Ld_UA	AA_N_Yld_Mass	AA_N_Yld_Ratio	AA_N_YId_UA	AA_OC_Ld_Mass	AA_OC_Ld_Ratio	AA_OC_Ld_UA	AA_OC_Yld_Mass	AA_OC_Yld_Rati 0	
2											
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30	
1	AA_OC_Yld_U A	AA_Pest_Ld_Mass	AA_Pest_Ld_Rati 0	AA_Pest_Ld_UA	AA_Pest_Yld_Mas s	AA_Pest_Yld_Rati	AA_Pest_Yld_UA	AA_P_Ld_Mass	AA_P_Ld_Ratio	AA_P_Ld_UA	
2											
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40	
1	AA_P_Yld_Ma ss	AA_P_YId_Ratio	AA_P_Yld_UA	AA_Pt_Src_Mass	AA_Pt_Src_Ratio	AA_Pt_Src_UA	AA_Sed_Eros_Ma ss	AA_Sed_Eros_Rat io	AA_Sed_Eros_UA	AA_Sed_Ld_Mass	
2											
Record	Field 41	Field 42	Field 43	Field 44	Field 45	Field 46	Field 47	Field 48	Field 49	Field 50	
1	AA_Sed_Ld_R atio	AA_Sed_Ld_UA	AA_Sed_Yld_Mas s	AA_Sed_Yld_Rati 0	AA_Sed_Yld_UA	AA_Wtr_Ld_Mass	AA_Wtr_Ld_Rati 0	AA_Wtr_Ld_UA	AA_Wtr_Yld_Mas s	AA_Wtr_Yld_Rati 0	
2											
Record	Field 51										
1	AA_Wtr_Yld_ UA										
2											
	Go to Data Section Description						Go to Table of Contents				

OUTPUT	OPTIONS	DATA –	CSV
--------	----------------	--------	-----

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	All_Evt_Lds_C ell_to_DS_Rchs	All_AA	All_Events	All_N	All_OC	All_Pesticides	All_P	All_Sediment	All_Water	AA_N_Ld_Cel_to_ DS_Rchs
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	AA_N_Ld_in_ Rchs	AA_N_Yld_Cel_to Rcv_Rch	AA_OC_Ld_Cel_t o_DS_Rchs	AA_OC_Ld_in_Rc hs	AA_OC_Yld_Cel_ to_Rcv_Rch	AA_Pest_Ld_Cel_t o_DS_Rchs	AA_Pest_Ld_in_R chs	AA_Pest_Yld_Cel_ to_Rcv_Rch	AA_P_Ld_Cel_to_ DS_Rchs	AA_P_Ld_in_Rchs
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	AA_P_Yld_Cel _to_Rcv_Rch	AA_BB_Eros_in_ Rch	AA_BB_Ld_in_DS _Rchs	AA_Eros_in_Cels	AA_Gly_Yld_Cel_ to_Rcv_Rch	AA_LS_Eros_in_C els	AA_LS_Ld_Cels_t o_DS_Rchs	AA_LS_Ld_in_DS _Rchs	AA_LS_Yld_Cel_t o_Rcv_Rch	AA_Rill_Eros_in_ Cels
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	AA_SR_Yld_C el_to_Rcv_Rch	AA_Wtr_Ld_Cel_t o_DS_Rchs	AA_Wtr_Ld_in_D S_Rchs	AA_Wtr_Yld_Cel_ to_Rcv_Rch	N_Evt_Ld_Cel_to_ DS_Rchs	N_Evt_Ld_in_Rch s	N_Evt_Yld_Cel_to _Rcv_Rch	OC_Evt_Ld_Cel_t o_DS_Rchs	OC_Evt_Ld_in_Rc hs	OC_Evt_Yld_Cel_ to_Rcv_Rch
2										
Record	Field 41	Field 42	Field 43	Field 44	Field 45	Field 46	Field 47	Field 48	Field 49	Field 50
1	Pest_Evt_Ld_C el_to_DS_Rchs	Pest_Evt_Ld_in_R chs	Pest_Evt_Yld_Cel _to_Rcv_Rch	P_Evt_Ld_Cel_to_ DS_Rchs	P_Evt_Ld_in_Rch s	P_Evt_Yld_Cel_to _Rcv_Rch	Sed_Evt_BB_Eros _in_Rch	Sed_Evt_BB_Ld_i n_DS_Rchs	Sed_Evt_Gly_Eros _in_Cels	Sed_Evt_Gly_Yld_ Cel_to_Rcv_Rch
2										
Record	Field 51	Field 52	Field 53	Field 54	Field 55	Field 56	Field 57	Field 58	Field 59	Field 60
1	Sed_Evt_LS_Er	Sed_Evt_LS_Ld_C	Sed_Evt_LS_Ld_i	Sed_Evt_LS_Yld_ Cel_to_Rcy_Rch	Sed_Evt_SR_Eros	Sed_Evt_SR_Yld_ Cel_to_Rev_Rch	Wtr_Evt_Ld_Cel_ to DS Rchs	Wtr_Evt_Ld_in_D S_Rchs	Wtr_Evt_Pk_Disc	Wtr_Evt_Yld_Cel
2										
Record	Field 61									
1	With Ext Desel		•							
	ow									
2	OW									

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Acc_Setup	Cell_Initial	Cell_TOC	Crp_Grwth	Data_Prep_Pointe rs	Weather	Opr_Rotation	Pest_Metabolite	Process_Flag	Quadrature
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Hydraulic_Geo m	Rch_Routing	Rch_TOC	RUSLE_C_Fctr	RUSLE_C_Fctr_S C	Canopy_Cover	Crp_Residue	Dead_Roots	PreProc_C_Fctr	Dom_Contour
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	EI_Pcts	RUSLE_Grwth_D ays	RUSLE_Init_Loc_ Oprs	RUSLE_K_Fctr	RUSLE_LS_Fctr	RUSLE_Non- crp_C_Fctr	RUSLE_Num_SLy r_SRes	RUSLE_P_Fctr	RUSLE_P_Fctr_C ntrs	RUSLE_P_Fctr_St rp
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	RUSLE_P_Fctr _Strp_Rot	RUSLE_Prior_LU	RUSLE_Res_Coef	RUSLE_Seg_Res	RUSLE_Setup_Pr d_Seg	RUSLE_Soil_Mois ture	RUSLE_Surf_Cov er	RUSLE_Surf_Rou gh	RUSLE_Unique_R es	Sed_Part_Distrib
2										
Record	Field 41	Field 42	Field 43	Field 44	Field 45	Field 46				
1	Seg_EI_Prcp	Setup_Seg	Soil_Comp_Surf	Soil_Comp_Lyrs	Storm_Types	Climate_Daily_Wt hr		-		-
2										
		<u>Go to Data Se</u>	ection Description				<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – DPP

OUTPUT OPTIONS DATA – EV

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	EV_Feedlots_M ass	EV_Feedlots_Rati 0	EV_Feedlots_UA	EV_Fld_Ponds_M ass	EV_Fld_Ponds_Ra	EV_Fld_Ponds_U A	EV_Gullies_Mass	EV_Gullies_Ratio	EV_Gullies_UA	EV_N_Ld_Mass
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	EV_N_Ld_Rati 0	EV_N_Ld_UA	EV_N_Yld_Mass	EV_N_Yld_Ratio	EV_N_Yld_UA	EV_OC_Ld_Mass	EV_OC_Ld_Ratio	EV_OC_Ld_UA	EV_ OC_Yld_Mass	EV_ OC_Yld_Ratio
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	EV_ OC_YId_UA	EV_ Pest_Ld_Mass	EV_ Pest_Ld_Ratio	EV_Pest_Ld_UA	EV_ Pest_Yld_Mass	EV_ Pest_Yld_Ratio	EV_Pest_Yld_UA	EV_P_Ld_Mass	EV_P_Ld_Ratio	EV_P_Ld_UA
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	EV_P_Yld_Mas s	EV_P_Yld_Ratio	EV_P_YId_UA	EV_Pt_Src_Mass	EV_Pt_Src_Ratio	EV_Pt_Src_UA	EV_Sed_Eros_Ma ss	EV_Sed_Eros_Rat io	EV_Sed_Eros_UA	EV_Sed_Ld_Mass
2										
Record	Field 41	Field 42	Field 43	Field 44	Field 45	Field 46	Field 47	Field 48	Field 49	Field 50
1	EV_Sed_Ld_Ra tio	EV_Sed_Ld_UA	EV_Sed_Yld_Mass	EV_Sed_Yld_Rati 0	EV_Sed_Yld_UA	EV_Wtr_Ld_Mass	EV_Wtr_Ld_Ratio	EV_Wtr_Ld_UA	EV_Wtr_Yld_Mas s	EV_Wtr_Yld_Rati 0
2										
Record	Field 51	Field 52	Field 53	Field 54						
1	EV_Wtr_Yld_ UA	LS_Rnof_All_Srcs	LS_Yld_All_Srcs	EV_Gullies_Erosio n						
2										
		Go to Data Se	ection Description				<u>Go to</u>	Table of Content	<u>.s</u>	

				00110	1 01 110116					
Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	AnnAGNPS_ID	Cell	Climate_Station	Contour	Сгор	Feedlot	Fertilizer	Mgmt_Seq	Field_Pond	Glbl_Output_Opts
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Gully	Hydraulic_Geom	Impoundment	Irrigation	Landuse_Ref	Reserved	Output_Options	Pesticide	Point_Source	Reach
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	Runoff_Curve_ Num	Simulation_Period	Soil_Actual_Surfa ce	Strip_Crop	Tile_Drain	Mgmt_Field	Mgmt_Sched	Mgmt_Opr	Soil_Actual_Layer s	Aquaculture_Pond
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36				
1	Aquaculture_P ond_Mgmt_Sch d_A	Glbl_Err/Wrn	Soil_Init_Cond	Pest_Init_Cond	Wetland	Riparian_Buffers				
2										
		Go to Data Se	ection Description				<u>Go to</u>	Table of Conten	ts	

OUTPUT OPTIONS DATA – NPT

OUTPUT OPTIONS DATA – SIM

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Cell_Componen ts	Conversion_Units	Sht/Rill_Eros_Sed _Yld	Feedlots	Insitu_N_Inorg	Insitu_N_Org	Insitu_Residue	Insitu_OC	Insitu_P_Inorg	Insitu_P_Org
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Insitu_Soil_Moi st_Daily	Irrigation	Pesticide_App	Pesticide_Insitu	Gully	Reach_Acc_Mass	Reach_Acc_Ratio	LS_Yld_All_Srcs	Reach_Ld_Nutr	Reach_Ld_Pest
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28	Field 29	Field 30
1	Reach_Ld_Sed	Reach_Ld_Wtr	Impound_Routing _A	Reach_Routing_N utr	Reach_Routing_Pe st	Reach_Routing	Reach_Routing_W tr	Runoff_Curve_Nu m	Schd_Oprs	Soil_Part_Distrib
2										
Record	Field 31	Field 32	Field 33	Field 34	Field 35	Field 36	Field 37	Field 38	Field 39	Field 40
1	Pond_Release/ Yield	Winter_Thermal	Winter_Summary	USLE_Params	Baseflow	Insitu_Soil_Moist_ Wshd_Sum	Wetland_Effects	Pot_ET_Adjust	LS_Rnof_All_Srcs	Riparian_Buffers
2										
		Go to Data Se	ection Description			•	<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – TBL

Record	Field 1	Field 2	Field 3	Field 4	Field 5					
1	CCHE1D	CONCEPTS_XM L	Gaging_Station_H yd	REMM	Gaging_Station_l vt	E				
2										
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – MIN/MAX (LIMITS)

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Min_Evt_Date	Max_Evt_Date	Max_Number_Evt s	Min_Rnof_Evt	Min_Rnof_Cell	Min_Rnof_Outlet	Min_Subarea_ID	Max_Subarea_ID	Subarea_Units_Po sitn	Max_Vrfy_File_Ac cess
2										
Record	Field 11	Field 12								
1	Max_Vrfy_File _Bytes	Input_Units_Code								
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA - CELL

Record	Field 1									
1	Cell_ID									
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – FEEDLOT

Record	Field 1									
1	Feedlot_ID									
2										
	Go to Data Section Description						<u>Go ta</u>	<u>o Table of Content</u>	<u>s</u>	

OUTPUT OPTIONS DATA – FIELD POND

Record	Field 1									
1	Field_Pond_ID									
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – GULLY, CLASSIC

Record	Field 1									
1	Classic_Gully_I D									
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – GULLY, EPHEMERAL

Record	Field 1									
1	Ephemeral_Gul ly_ID									
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – IMPOUNDMENT

Record	Field 1									
1	Impoundment_ ID									
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – POINT SOURCE

Record	Field 1							
1	Point_Source_I D							-
2								
		<u>Go to Data S</u>	ection Description		<u>Go to</u>	Table of Content	<u>s</u>	

OUTPUT OPTIONS DATA – REACH

Record	Field 1											
1	Reach_ID											
2												
	Go to Data Section Description					<u>Go to</u>	Go to Table of Contents					

OUTPUT OPTIONS DATA – WETLAND

Record	Field 1									
1	Wetland_ID									
2										
	Go to Data Section Description						<u>Go te</u>	o Table of Conten	t <u>s</u>	

PESTICIDE APPLICATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8		
1	Application_ID	Reference_ID	Application_Rate	Depth	Mixing_Code	e Foliage_Fraction	Soil_Fraction	Input_Units_Code		
2										
		<u>Go to Data S</u>	ection Description				<u>Go ta</u>	Table of Content	<u>is</u>	

PESTICIDE INITIAL CONDITIONS DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5					
1	Initial_Pesticide _ID	Crop_Initial_Amo unt_1	Crop_Initial_Amo unt_2	Non- Crop_Initial_Amo unt_1	Non- Crop_Initial_An unt_2	10				
2										
		<u>Go to Data S</u>	ection Description				<u>Go to</u>	Table of Content	<u>ts</u>	

PESTICIDE REFERENCE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	
1	Pesticide_Refer ence_ID	Solubility	Partition	Soil_Half-life	Foliage_Half-life	Washoff	Metabolite_ID	Metabolite_Transf ormation	Reach_Half-life	
2										
		<u>Go to Data Se</u>	ection Description				<u>Go ta</u>	<u>o Table of Content</u>	<u>s</u>	

PL CALIBRATION DATA:

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	OC_All_Source	OC_Sheet_and_Ril l	OC_Feedlot	OC_Point_Source	OC_Gully	OC_Pond	OC_Irrigation	N_All_Sources	N_Sheet_and_Rill	N_Feedlot	
2											
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20	
1	N_Point_Sourc e	N_Gully	N_Pond	N_Irrigation	P_All_Sources	P_Sheet_and_Rill	P_Feedlot	P_Point_Source	P_Gully	P_Pond	
2											
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28			
1	P_Irrigation	Sediment_All_Sou rces	Sediment_Sheet_a nd_Rill	Sediment_Feedlot	Sediment_Point_S ource	Point_S e Sediment_Gully Sediment_Pond Sediment_Irrigatio n					
2											
		Go to Data Se	ection Description				<u>Go to</u>	Table of Content	<u>s</u>		

POINT SOURCE DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Point_Source_I D	Cell_ID	Point_Flow	Point_N	Point_P	Point_OC	OC_Calib_Fctr	N_Calib_Fctr	P_Calib_Fctr	Erosion_Calib_Fct r
2										
Record	Field 11									
1	Input_Units_Co de									
2										
		<u>Go to Data S</u>	ection Description				<u>Go te</u>	Table of Content	t <u>s</u>	

RCN CALIBRATION DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5		Field 6	Field 7	Field 8	Field 9	
1	RCN_Calib_ID	Target_AA_Direct _Runoff_Load	Prev_Estimate_for _Water_Load	RCN_Retention_F ctr	Reach_ID_Si	ite	Reach_Ratio	Avbl_Soil_Moist_ AMC_II	Target_AA_Total_ Streamflow	Input_Units_Code	
2											
		<u>Go to Data Se</u>	ection Description					<u>Go to</u>	Table of Content	<u>s</u>	

REACH DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Reach_ID	Receiving_Reach	Vegetation_Code	Elevation	Slope	Mannings_n	Infiltration_Rate	Hydraulic_Geom_ ID	Length	Top_Width
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Flow_Depth	Valley_Width	Valley_Mannings_ n	Start_Diversion	Stop_Diversion	Travel_Time	Clay_Scour_Code	Silt_Scour_Code	Sand_Scour_Code	Small_Agg_Scour_ Code
2										
Record	Field 21	Field 22	Field 23	Field 24	Field 25	Field 26	Field 27	Field 28		
1	Large_Agg_Sco ur_Code	Valley_Clay_Scour _Code	Valley_Silt_Scour_ Code	Valley_Sand_Scou r_Code	Valley_Small_Agg _Scour_Code	Valley_Large_Agg _Scour_Code	Delivery_Ratio	Input_Units_Code		
2										
	•	<u>Go to Data Se</u>	ection Description			·	<u>Go to</u>	Table of Content	<u>s</u>	

REACH NUTRIENT HALF-LIFE

Record	Field 1	Field 2	Field 3					
1	N_Half-life	P_Half-life	OC_Half-life					
2								
	•	<u>Go to Data Se</u>	ection Description		<u>Go to</u>	Table of Content	<u>s</u>	

RIPARIAN BUFFER DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Buffer_ID	Location_ID	Vegetative_Type	Buffer_Slope	Max_Trap_Efficie ncy	Eff_Buffer_Width	Eff_Conc_Flow_W idth	Drainage_Area_to _Buffer	Buffer_Location_ Code	Actual_Trap_Effic iency_Clay
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Actual_Trap_E fficiency_Silt	Actual_Trap_Effic iency_Sand	Actual_Trap_Effic iency_Sm_Agg	Actual_Trap_Effic iency_Lg_Agg	Fraction_Trapped _Clay	Fraction_Trapped _Silt	Fraction_Trapped _Sand	Fraction_Trapped _Sm_Agg	Fraction_Trapped _Lg_Agg	Input_Units_Code
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

RUNOFF CURVE NUMBER DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5				
1	Curve_Number _ID	CN_A	CN_B	CN_C	CN_D				
2									
Go to Data Section Description						<u>Go ta</u>	Table of Content	<u>is</u>	

RUSLE2 DATA

Record	Field 1	Field 2	Field 3						
1	RUSLE2_ID	RUSLE2_Databas e	"RUSLE2_Erosion _Flag"						
2									
Go to Data Section Description						<u>Go ta</u>	Table of Content	<u>s</u>	

SIMULATION PERIOD DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Simulation_Beg in_Month	Simulation_Begin_ Day	Simulation_Begin_ Year	Simulation_End_ Month	Simulation_End_D ay	Simulation_End_Y ear	Rainfall_Fctr	10-Year_EI	EI_Number	Irrigation_Climate _Code
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17			
1	Soil_Moisture_ Steps	Annual_K_Fctr_C ode	Variable_K_Fctr_ Code	Number_Init_Year s	Init_Method_Code	Winter_Bouts	Input_Units_Code			
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

SOIL DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Soil_ID	Hydrologic_Soil_G roup	K_Factor	Albedo	Time_to_Consolid ation	Impervious_Depth	Specific_Gravity	Initial_Soil_Condit ions_ID	Soil_Name	Soil_Texture
2										
Record	Field 11	Field 12								
1	Number_of_Soi l_Layers	Input_Units_Code								
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

SOIL INITIAL CONDITIONS DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Initial_Soil_Co nditions_ID	Landuse_Type_ID (Currently not used)	Inorganic_N_1	Inorganic_N_2	Inorganic_P_1	Inorganic_P_2	Soil_Moisture_1	Soil_Moisture_2	Organic_Matter_1	Organic_Matter_2
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Organic_N_1	Organic_N_2	Organic_P_1	Organic_P_2	Surface_Residue	Mannings_n	Snow_Depth	Snow_Density	Surface_Constant	Input_Units_Code
2										
		Go to Data Se	ection Description	•		•	<u>Go to</u>	Table of Content	<u>s</u>	

SOIL LAYER DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Soil_ID	Layer_Number	Layer_Depth	Bulk_Density	Clay_Ratio	Silt_Ratio	Sand_Ratio	Rock_Ratio	Very_Fine_Sand_ Ratio	CaCO3_Content
2										

AnnAGNPS Input File Specifications2019File Name: Input_Specifications_v5.51_2019.xx.xx.docx

Revision: December 4, 2019

Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20
1	Saturated_Con ductivity	Field_Capacity	Wilting_Point	Volcanic_Code	Base_Saturation	Unstable_Aggregat e_Ratio	рН	Organic_Matter_R atio	Organic_N_Ratio	Inorganic_N_Ratio
2										
Record	Field 21	Field 22	Field 23	Field 24						
1	Organic_P_Rat io	Inorganic_P_Ratio	Soil_Structure_Co de	Input_Units_Code						
2										
	Go to Data Section Description						<u>Go t</u>	Table of Content	<u>s</u>	

STRIP CROP DATA

Record	Field 1	Field 2	Field 3							
1	Strip_Crop_ID	P_Factor	Delivery_Ratio							
2										
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>ts</u>	

TILE DRAIN DATA

Record	Field 1	Field 2	Field 3	Field 4					
1	Tile_Drain_ID	Drain_Rate	Invert_Depth	Input_Units_Code					
2									
		<u>Go to Data Se</u>	ection Description			<u>Go to</u>	Table of Content	<u>s</u>	

WATERSHED DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5					
1	Wshd_Name	Wshd_Description	Wshd_Location	Latitude	Longitude					
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	t <u>s</u>	

WETLAND DATA

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Wetland_ID	Reach_ID	Wetland_Area	Initial_Water_Dep th	Min_Water_Depth	Max_Water_Dept h	Water_Temperatu re	Potential_Daily_In filtration	Weir_Coef	Weir_Width
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17			
1	Weir_Height	Soluble_N_Conc	Nitrate- N_Loss_Rate	Nitrate- N_Loss_Rate_Coef	Temperature_Coef	Weir_Exp	Input_Units_Code			
2										
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>s</u>	

Daily Climate Data

The daily climate data is in a separate file from other input for AnnAGNPS. The climate data can either be inputted as fixedformatted data as specified in the fixed-formatted version of the input specifications document or as csv-formatted data according to the tables below.

There are two possible climate related csv files; 1.) climate station csv file, 2.) daily climate data csv file. The filenames of the primary climate files are user-defined and placed in the csv master list file where AnnAGNPS will read the filenames and open the files for input. The required data section name for identification in the csv master list file for the climate station information is "Climate Data - Station". The required data section name for identification in the csv master list file for the daily climate data information is "Climate Data - Data - Daily".

Primary climate data is required for AnnAGNPS, therefore, if primary climate data is in csv-format then both station and daily csv input files are required in the AnnAGNPS csv master list. If climate data is not specified in the csv master list file then fixed-formatted climate data is assumed according to the fixed-formatted version of the input specifications document

Secondary climate files are optional. If secondary climate files are to be included, they must be in the same format as the primary climate file whether fixed-formatted or csv-formatted. Secondary filenames are not included in the csv master list file. Rather, they are internally defined within AnnAGNPS based on the primary climate filename. See the "Daily Climate input file" section of the document called "AnnAGNPS User's Guide for Input Files & Formats" for more information.

All data must be complete and have continuous daily record. Data can be from an actual weather station, a combination of weather stations, synthetically generated data, or a composite of weather stations & synthetically generated data. Note: Longitude, latitude, & wind direction are expressed in decimal degree units—<u>not</u> degrees & minutes for units.

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-24
The following reco	ord does not re	peat.		i	i	i
Climate Input Version ID — Unique alphanumeric string identifying the climate file's input version number	"Version"		5.51	A10	2	1
Climate Input Units code — Code identifying whether input is in English or SI units. $0 =$ English, $1 =$ SI (Blank defaults to the input units code in the AnnAGNPS ID data section)	"Input_Units_C ode"		Blank, 0 or 1	I1	2	2
Climate File Name — Alphanumeric string describing the climate station name and description.	"Climate_Statio n_Name"			A80	2	3
Climate Begin Date—Beginning month, day, & year for the weather data. Year can be specified as calendar years (e.g., 1960) or as simulation year which must be year 1. If blank, the date of the first daily record will be used.	"Beginning_Cli mate_Date"	mm/dd/yyyy	Blank, or mm—1 to 12 dd—1 to 31 yyyy—1 to 9999	A10	2	4
Climate End Date —Ending month, day, & year for the weather data. Year can be specified as calendar years (e.g., 1960) or as simulation year which must be the total number of years for the simulation. If blank, the date of the last daily record will be used.	"Ending_Climat e_Date"	mm/dd/yyyy	Blank, or mm—1 to 12 dd—1 to 31 yyyy—1 to 9999	A10	2	5
Station Latitude —Representative climate station latitude. Latitude is expressed as "+" for North and "-" for South.	"Latitude"	^O lat	-90. to 90.	F10	2	6
Station Longitude —Representative climate station longitude. Longitude is expressed as "+" for East and "-" for West.	"Longitude"	^o long	-180. to 180.	F10	2	7
Station Elevation—Representative climate station elevation.	"Elevation"	{ft} [m]	{-3280.8 to 32808.3} [-1000.0 to 10000.0]	F10	2	8
Adiabatic Air Temperature Lapse Rate—Air temperature change with respect to representative climate station elevation. Default is a decrease of 2°F for every 1000 ft increase in elevation (-0.002 °F/ft.).	"Temperature_ Lapse_Rate"	{°F / ft} [°C / m]	Blank or {-0.002 to 0.002} [-0.0032 to 0.0032]	F10	2	9

CLIMATE STATION INFORMATION

AnnAGNPS Input File Specifications

	Field	Unite				
Description	Header for	(English)	Domain	For	Record	Field
Description	Decord #1	{English; [SI]	{English} [SI]	mat	No.	No.
Descinitation Nitrogan Nitrogan concentration in precipitation	"Precipitation	$1 \sim N/1$	Dlank or	E10	r	10
Precipitation Introgen—Introgen concentration in precipitation.	N"	I g-IN / I Ma precip	0.0 ± 0.10	L10	2	10
Blank defaults to 0.0.		(nnm)	0.0 10 1.0			
Clabel Storm Type ID Storm type ID which will be used for all	"Clobal Storm	(ppm)	Diantz preset IDs	A 20	r	11
ctorms designated by this climate file excent where seasonal or daily	Type_ID"		for SCS synthetic	A20	2	11
IDs are given. Acceptable IDs are: (1) blank which defaults to the			storm types or			
Watershed Storm Type code designated in the watershed data input			user-defined IDs			
file's GLOBAL IDS. FACTORS. AND FLAGS DATA section: (2)			for user			
any of the preset storm type IDs: or (3) any new storm type IDs			requested input.			
given in the STORM TYPE DATA input file. Preset IDs are: (1)			Blank defaults to			
Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type II; (4)			the watershed			
Std. SCS Type III; (5) Std. Uniform; (6) Std. SCS NM60; (7) Std.			storm type as			
SCS NM65; (8) Std. SCS NM70; & (9) Std. SCS NM75. New storm			given by its code			
type IDs require user input for the rainfall distribution.			in the GLOBAL			
			IDS, FACTORS,			
			AND FLAGS			
			DATA section			
Elevation Difference (1)—1 st elevation with respect to	"1 st _Elevation_	{feet}	Blank, or	F10	2	12
representative climate station elevation for rainfall correction	Difference	[m]	{-1500.0 to			
relationship. Paired with 1 st spatial rainfall factor. Default is no			30000.0}			
change from representative station rainfall. Blank may be entered			[-500.0 to			
if no variation of precipitation with elevation is desired. If blank,			10000.0			
then Elevation Rain Factor (1), Elevation Difference (2), and						
Elevation Rain Factor (2) must also be blank.	((1 of 151)	5 1	51 1			10
Elevation Rain Factor (1)— 1^{st} average annual rainfall factor with	"1"_Elevation_ Rain Factor"	Depth-	Blank, or	F10	2	13
respect to representative climate station precipitation for rainfall	·····_·	annual	0.00001 to 10.0			
correction relationship. Paired with 1 st spatial rainfall elevation.		precip at				
Blank may be entered if no variation of precipitation with circulation		donth				
Difference (2) and Elevation Rain Factor (2) must also be blank		annual				
Difference (2) , and Elevation Rain ratio (2) must also be stank.		nrecin at				
		station				
		(dimension				
		less)				
Elevation Difference (2) —2 nd elevation with respect to	"2 nd _Elevation_	{feet}	Blank, or	F10	2	14
representative climate station elevation for rainfall correction	Difference"	[m]	{-1500.0 to			
relationship. Paired with 2 nd spatial rainfall factor. Blank may be			30000.0}			
entered if no variation of precipitation with elevation is desired. If			[-500.0 to			
blank, then Elevation Difference (1), Elevation Rain Factor (1), and			10000.0			
Elevation Rain Factor (2) must also be blank.						
Elevation Rain Factor (2)—2 nd average annual rainfall factor with	"2 nd _Elevation_	Depth-	Blank, or	F10	2	15
respect to representative climate station precipitation for rainfall	Kam_ractor	annual	0.00001 to 10.0			
correction relationship. Paired with 2 nd spatial rainfall elevation.		precip at				
Blank may be entered if no variation of precipitation with elevation		2 nd elev/				
is desired. It blank, then Elevation Difference (1), Elevation Kain $D_{1}^{(1)} = \frac{1}{2} \sum_{i=1}^{n} $		depth-				
Factor (1), and Elevation Difference (2) must also be blank.		annual				
		precip at				
		station				
2 V- 24 Hu Description Maximum 24 hour presinitation that is	"? Vr 24 hr P	Gin	(0.04 ± 12.0)	E10	r	16
2 11 24 mil riccipitation —Maximum 24 nour precipitation unat is	recipitation"	{III}}	$\{0.04 \ 10 \ 12.0\}$	F10	2	10

AnnAGNPS Input File Specifications Revision: December 4, 2019 File Name: Input_Specifications_v5.51_2019.xx.xx.docx

	T , 11	T T •/				
	Field	Units	Domain	For	Record	Field
Description	Header for	{English}	{English} [SI]	mat	No.	No.
	Record #1	[81]		F10		15
Rainfall Calibration or Areal Correction Coefficient—when used	"Calibration_or Areal Correcti		Blank or	F10	2	17
as an areal correction, then this field is an exponential curve	on_Coefficient"		it rainfall			
coefficient used to correct point rainfall to average rainfall over the			calibration, then			
climate station (units must be consistent within the power curve):			>0			
$correction = 1 - \left\{ coef \bullet \left\lfloor 1 - e^{-(\exp \bullet D_a)} \right\rfloor \right\}$			correction, then			
where: correction = areal rainfall correction factor [decimal]			0 < coef < 0.02			
coef = rainfall areal correction coefficient						
exp = rainfall areal correction exponent						
$D_a =$ drainage area of climate station [ac or na].						
A blank defaults to 1. If this field is blank, then the Areal Kalmal						
Desinfall Correction Coefficient field is blank. If the Areal						
rainfall collibration factor. Tin. TP40 fig 15 suggests that a close						
fit for a 24 hour storm's areal correction coefficient would be 0.002						
where D can be in either acres or bectares						
A real Dainfall Correction Exponent only used as an areal	"Calibration or		Blank or	E10	2	18
correction exponential curve exponent to correct point rainfall to	_Areal_Correcti		$\{0, 0, to 1\}$	110	2	10
average rainfall over the climate station :	on_Exponent"		[0.0 to 1.]			
			[0.0 to 1.]			
$correction = 1 - \left\{ coef \bullet \left\lfloor 1 - e^{-(\exp \bullet D_a)} \right\rfloor \right\}$						
where: correction = areal rainfall correction factor [decimal]						
coef = rainfall areal correction coefficient						
exp = rainfall areal correction exponent						
D_a = drainage area of climate station [ac or ha].						
A blank defaults to 0. If the Areal Rainfall Correction						
Coefficient field is blank, then this field must also be blank. Tip—						
TP40, fig.15 suggests that a corresponduing exponent to the TP40						
areal correction coefficient of 0.092 is 0.000023 when Da is acres						
and 0.000057 when D_a is in hectares. The difference between T(40						
and this exponential formula is less than ± 0.6 %.	(() K) T (-		
Minimum interception evaporation—Minimum evaporation	"Minimum_Inte	{ in }	Blank or	F10	2	19
associated with the interception for each precipitation event. Blank	ration"	[mm]	$\{0 \text{ to } 0.25\}$			
defaults to 1.000mm.	"M	()	[0 to 6.350]	F10	2	20
Maximum interception evaporation—Maximum evaporation	rception Evapo	{in}	Blank or	F10	2	20
associated with the interception for each precipitation event. Blank	ration"	[mm]	$\{0 \ 10 \ 0.25\}$			
defaults to 2.500mm.	"Winton Storm		[0 to 0.550]	1 20	2	21
for all storms designated by this alignets file during the winter	Type ID"		for SCS synthetic	A20	2	21
nor all storms designated by this climate the during the winter			storm types or			
Acceptable IDs are: (1) blank which defaults to the Watershed			user-defined IDs			
Storm Type code designated in the watershed data input file's			requiring			
SIMUL ATION PERIOD DATA section: (2) any of the preset storm			additional input			
type IDs: or (3) any new storm type IDs given in the storm type			Blank defaults to			
input file., Preset IDs are: (1) Std. SCS Type I: (2) Std. SCS Type			the global storm			
Ia: (3) Std. SCS Type II: (4) Std. SCS Type III: (5) Std. Uniform: (6)			type.			
Std. SCS NM60; (7) Std. SCS NM65: (8) Std. SCS NM70: & (9)			-7 F			
Std. SCS NM75. New storm type IDs require user input for the						
rainfall distribution.						

AnnAGNPS Input File Specifications Revision: December 4, 2019 File Name: Input_Specifications_v5.51_2019.xx.xx.docx

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Spring Season Storm Type ID —Storm type ID which will be used for all storms designated by this climate file during the spring months (21 Mar – 21 June) except where daily IDs are given. Acceptable IDs are: (1) blank which defaults to the Watershed Storm Type code designated in the watershed data input file's SIMULATION PERIOD DATA section; (2) any of the preset storm type IDs; or (3) any new storm type IDs given in the storm type input file Preset IDs are: (1) Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type II; (4) Std. SCS Type III; (5) Std. Uniform; (6) Std. SCS NM60; (7) Std. SCS NM65; (8) Std. SCS NM70; & (9) Std. SCS NM75. New storm type IDs require user input for the rainfall distribution.	"Spring_Storm_ Type_ID"		Blank, preset IDs for SCS synthetic storm types, or user-defined IDs requiring additional input. Blank defaults to the global storm type.	A20	2	22
Summer Season Storm Type ID—Storm type ID which will be used for all storms designated by this climate file during the summer months (22 Jun – 21 Sep) except where daily IDs are given. Acceptable IDs are: (1) blank which defaults to the Watershed Storm Type code designated in the watershed data input file's SIMULATION PERIOD DATA section; (2) any of the preset storm type IDs; or (3) any new storm type IDs given in the storm type input file Preset IDs are: (1) Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type II; (4) Std. SCS Type III; (5) Std. Uniform; (6) Std. SCS NM60; (7) Std. SCS NM65; (8) Std. SCS NM70; & (9) Std. SCS NM75. New storm type IDs require user input for the rainfall distribution.	"Summer_Stor m_Type_ID"		Blank, preset IDs for SCS synthetic storm types, or user-defined IDs requiring additional input. Blank defaults to the global storm type.	A20	2	23
Autumn Season Storm Type ID— Storm type ID which will be used for all storms designated by this climate file during the autumn months (22 Sep – 21 Dec) except where daily IDs are given. Acceptable IDs are: (1) blank which defaults to the Watershed Storm Type code designated in the watershed data input file's SIMULATION PERIOD DATA section; (2) any of the preset storm type IDs; or (3) any new storm type IDs given in the storm type input file Preset IDs are: (1) Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type II; (4) Std. SCS Type III; (5) Std. Uniform; (6) Std. SCS NM60; (7) Std. SCS NM65; (8) Std. SCS NM70; & (9) Std. SCS NM75 New storm type IDs require user input for the rainfall distribution.	"Autumn_Stor m_Type_ID"		Blank, preset IDs for SCS synthetic storm types, or user-defined IDs requiring additional input. Blank defaults to the global storm type.	A20	2	24
Go to Layout Matrix		Go to	o Table of Conten	ts		

DAILY CLIMATE INFORMATION

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.					
Field Header— Required unique field header for each field listed				Α	1	1-16					
below.											
The following line is repeated for each day of the climate record.											
Daily Weather Date—Month, day, & year according to the	"Month",	mm	mm—1 to 12	A2,	2	1-3					
Gregorian calendar for the weather data for the day specified. Years	"Day", "Vear"	dd	dd—1 to 31	A2,							
can be specified as calendar years (e.g., 1960) or as simulation years	i cai	уууу	yyyy—1 to 9999	A4							
which must begin with year 1.											
Daily Max Temperature—Maximum air temperature for the day	"Max_Air_Tem	${^{O}F}$	{-100.0 to 150.0}	F10	2	4					
specified.	perature"	[°C]	[-75.0 to 65.0]								
Daily Min Temperature—Minimum air temperature for the day	"Min_Air_Tem	${^{O}F}$	{-100.0 to 150.0}	F10	2	5					
specified.	perature"	[°C]	[-75.0 to 65.0]								
Daily Precipitation—Total precipitation for the day specified.	"Precip"	{in}	{0.0 to 30.0}	F10	2	6					
		[mm]	[0.0 to 750.]								
Daily Dew Point Temperature—24-hour average dew point	"Dew_Point"	{ ⁰ F}	{-100.0 to 150.0}	F10	2	7					
temperature for the day specified.		[°C]	[-75.0 to 65.0]								

	Field	IIta	-			
Description	Header for	{English}	Domain	For	Record	Field
	Record #1	[SI]	{English} [SI]	mat	No.	No.
Daily Sky Cover—24-hour average total opaque sky cover for the	"Sky_Cover"	%	Blank, 0.0 to	F10	2	8
day specified. If blank, then its corresponding daily solar radiation			100.0			
at ground level must be given.						
Daily Wind Speed—24-hour average wind speed for the day	"Wind_Speed"	{mph}	{0.0 to 200.0}	F10	2	9
specified.		[m / sec]	[0.0 to 90.0]			
Daily Wind Direction—24-hour average wind direction for the day	"Wind_Directio	Decimal ^o	0.0 to 360.0	F10	2	10
specified. Measured clockwise degrees from north.	n"					
Daily Solar Radiation at Ground Level—solar radiation at ground	"Solar_Radiatio	{BTU/hr/ft	{0.0 to 433.34}	F10	2	11
surface. Optional input for sky cover. If sky cover is blank, then	n"	^2}	[0.0 to 1367]			
this field will be used to calculate sky cover. Based on ASCE		[J/sec/m^2				
(1996).]				
Daily Storm Type ID—Storm ID for this day indicating one of the	"Storm_Type_I		Blank, preset IDs	A20	2	12
preset synthetic or user-requested storm type IDs to use with pre-	D"		for SCS synthetic			
calculated solutions using extended TR-55: Acceptable IDs are:			storm types, or			
blank which defaults to seasonal storm type. Preset IDs are: (1)			user-defined IDs			
Std. SCS Type I; (2) Std. SCS Type Ia; (3) Std. SCS Type I I; (4)			for user			
Std. SCS Type III; (5) Std. Uniform; (6 Std. SCS NM60; (7) Std.			requested input.			
SCS NM 65; (8) Std. SCS NM 70; & (9) Std. SCS NM75. Other			Blank defaults to			
IDs require user input for both the rainfall distribution & unit peak			seasonal storm			
discharge regression coefficients.			types.			
Potential ET—User supplied potential ET value for the current day.	"Potential_ET"	{in}	Blank, or	F10	2	13
If blank, the potential ET will be calculated internally.		[mm]	{0.0 to 30.0}			
			[0.0 to 750.]			
Actual ET—User supplied actual ET value for the current day. If	"Actual_ET"	{in}	Blank, or	F10	2	14
blank, the actual ET will be calculated internally.		[mm]	{0.0 to 30.0}			
			[0.0 to 750.]			
Actual EI—User supplied actual EI value for the current day. If	"Actual_EI"	{(100 ft-	Blank, or	F10	2	15
blank, the EI will be calculated internally.		ton-in) /	{0.0 to 1175.0}			
		(ac-hr)}	[0.0 to 20000.]			
		[MJ-				
		mm/ha-hr]				
Climate Input Units code — Code identifying whether input is in	"Input_Units_C		Blank, 0 or 1	I1	2	16
English or SI units. $0 =$ English $, 1 =$ SI (Blank defaults to the input	ode"					
units code in the AnnAGNPS ID data section)						
Go to Layout Matrix		<u>Go to</u>	Table of Conten	ts		

AnnAGNPS Input File Specifications Revision: December 4, 2019 File Name: Input_Specifications_v5.51_2019.xx.xx.docx

CLIMATE INPUT FILE LAYOUT MATRIX – STATION

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10	
1	Version	Input_Units_Code	Climate_Station_D escription	Beginning_Climate _Date	Ending_Climate_D ate	Latitude	Longitude	Elevation	Temperature_Lap se_Rate	Precipitation_N	
2	5.51										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20	
1	Global_Storm_ Type_ID	1st_Elevation_Diff erence	1st_Elevation_Rai n_Factor	2nd_Elevation_Dif ference	2nd_Elevation_Rai n_Factor	2_Yr_24_hr_Preci pitation	Calibration_or_Ar eal_Correction_Co efficient	Calibration_or_Ar eal_Correction_Ex ponent	Minimum_Interce ption_Evaporation	Maximum_Interce ption_Evaporation	
2											
Record	Field 21	Field 22	Field 23	Field 24							
1	Winter_Storm_ Type_ID	Spring_Storm_Ty pe_ID	Summer_Storm_T ype_ID	Autumn_Storm_T ype_ID							
2											
		<u>Go to Data Se</u>	ection Description			Go to Table of Contents					

CLIMATE INPUT FILE LAYOUT MATRIX – DAILY

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Month	Day	Year	Max_Air_Temper ature	Min_Air_Tempera ture	Precip	Dew_Point	Sky_Cover	Wind_Speed	Wind_Direction
2										
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 15				
1	Solar_Radiatio n	Storm_Type_ID	Potential_ET	Actual_ET	Actual_EI	Input_Units_Code				
2										
	Go to Data Section Description						<u>Go to</u>	Table of Content	<u>s</u>	

Storm Type Data

Storm type data is an optional input into AnnAGNPS. The storm type data can either be inputted as fixed-formatted data as specified in the fixed-formatted version of the input specifications document or as csv-formatted data according to the tables below.

There are two possible storm type related csv files; 1.) Rainfall distribution csv file (RFD), 2.) Unit peak discharge regression coefficients csv file (UPDRC). The filenames are user-defined and placed in the master list file where AnnAGNPS will read the filenames and open the files for input. Although storm type data is optional, if the UPDRC is present in the AnnAGNPS master list then the RFD file is required.

Data can be from an actual storm patterns, a combination of patterns, synthetically generated patterns, or a composite of actual patterns & synthetically generated data.

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.					
Field Header— Required unique field header for each field listed below.				А	1	1-243					
The following record repeats for the number of rainfall distribution records.											
Storm Type File Input Version ID — Unique alphanumeric string identifying the storm type file's input version number	"Version"		5.51	A10	2	1					
StormType ID — Unique storm type ID.	"Storm_Type_I D"			A20	2	2					
Accumulative Rainfall Amount—Monotonically increasing accumulative rainfall amount expressed as the ratio of rainfall at time t _i to the total 24-hr rainfall in 0.1 hour increments.	"0.0", "0.1", "0.2", "0.3", "24.0"	(mm-t _{i)} / (mm-t ₂₄)	0.0 - 1.0	F10	2	3-243					
Go to Layout Matrix	Go to Table of Contents										

RAINFALL DISTRIBUTION

UNIT PEAK DISCHARGE REGRESSION COEFFICIENTS

Description	Field Header for Record #1	Units {English} [SI]	Domain {English} [SI]	For mat	Record No.	Field No.
Field Header— Required unique field header for each field listed below.				А	1	1-121
The following record repeats for the n	umber of unit	peak discha	rge records.		-	
StormType ID — Unique storm type ID. Must match a "StormType ID" in the "Rainfall Distribution" table; otherwise, the record will be ignored.	"Storm_Type_I D"			A20	2	1
Regression Coefficients —the six (6) regression coefficients, a-f, for the rational polynominal: qup = $\{a+[c*tc]+[e*(tc**2)]\}/\{1+[b*tc]+[d*(tc**2)]+[f*(tc**3)]\}$ Where Ia/P ₂₄ which indicates the initial abstraction/24-hr rainfall ratio for the following fields from 0.00 to 0.95 in 0.05 increments.	" $a@0.00$ ", " $b@0.00$ ", " $c@0.00$ ", " $c@0.00$ ", " $f@0.00$ ", " $f@0.00$ ", " $f@0.05$ ", " $b@0.05$ ", " $t@0.05$ ", " $t@0.95$ ",	-	-∞ to +∞	E15	2	2-121
Go to Layout Matrix		<u>Go to</u>	o Table of Conten	<u>ts</u>		

STORM TYPE INPUT FILE LAYOUT MATRIX – RAINFALL DISTRIBUTION

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Version	Storm_Type_ID	0.0	0.1	0.2	0.3	0.4	0.5	0.6	0.7
2	5.51		amount at t _{0.0} =0.0	amount at t=0.1	amount at t=0.2	amount at t=0.3	amount at t=0.4	amount at t=0.5	amount at t=0.6	amount at t=0.7
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20-243
1	0.8	0.9	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7 – 24.0
2	amount at t=0.8	amount at t=0.9	amount at t=1.0	amount at t=1.1	amount at t=1.2	amount at t=1.3	amount at t=1.4	amount at t=1.5	amount at t=1.6	amount at t=1.7 – 24.0
Go to Data Section Description							<u>Go to</u>	Table of Content	<u>s</u>	

STORM TYPE INPUT FILE LAYOUT MATRIX – UNIT PEAK DISCHARGE REGRESSION COEFFICIENTS

Record	Field 1	Field 2	Field 3	Field 4	Field 5	Field 6	Field 7	Field 8	Field 9	Field 10
1	Storm_Type_I D	a@0.00	b@0.00	c@0.00	d@0.00	e@0.00	f@0.00	a@0.05	b@0.05	c@0.05
2		Coefficient "a" for Ia/P=0.00	Coefficient "b" for Ia/P=0.00	Coefficient "c" for Ia/P=0.00	Coefficient "d" for Ia/P=0.00	Coefficient "e" for Ia/P=0.00	Coefficient "f" for Ia/P=0.00	Coefficient "a" for Ia/P=0.05	Coefficient "b" for Ia/P=0.05	Coefficient "c" for Ia/P=0.05
Record	Field 11	Field 12	Field 13	Field 14	Field 15	Field 16	Field 17	Field 18	Field 19	Field 20-121
1	d@0.05	e@0.05	f@0.05	a@0.10	b@0.10	c@0.10	d@0.10	e@0.10	f@0.10	a@0.15 - f@0.95
2	Coefficient "d" for Ia/P=0.05	Coefficient "e" for Ia/P=0.05	Coefficient "f" for Ia/P=0.05	Coefficient "a" for Ia/P=0.10	Coefficient "b" for Ia/P=0.10	Coefficient "c" for Ia/P=0.10	Coefficient "d" for Ia/P=0.10	Coefficient "e" for Ia/P=0.10	Coefficient "f" for Ia/P=0.10	Coefficient "a" for Ia/P=0.15 - Coefficient "f" for Ia/P=0.95
Go to Data Section Description						Go to Table of Contents				

Appendix A: Output Files

Versions 1 & 2 Output-Related Options

Although the original four output-related data sections (five counting the "out-of-pocket" CONCEPTS output file request in the Watershed Data section) will remain within AnnAGNPS for an indefinite time; no further coding support will be provided for them. In fact, if any part of the output options feature is activated by input, AnnAGNPS will ignore entirely any of the versions 1 & 2 output-related data sections during execution. In short, if the "Output Options Data:" section(s) is activated, only the logic associated with the new output options will be used during execution.

The older versions of the output-related sections were very difficult to use and the various verification output data were all forced into the same file which made them almost unusable except when only an individual verification file was requested; and they did not always activate as indicated by the Editor.

Output File Structure

All current & future output will be under positive control of the user through Output Options Data section(s) and, therefore, will be accessible through the Editor or other software products.

Output files are typically categorized by type. The output filenames produced by AnnAGNPS have characters embedded that indicate the type of output contained in the file. The different types are "AA"- average annual, "CSV" - database, "DPP" - data preprocessing, "EV" - event, "NPT" - input, "SIM" - simulation, and "TBL" - tables. Output files are largely written in a fixed format. All extensions used are fixed format output files with the exception of ".csv" extensions. These are csv-formatted output files.

All output will be under user control according to typical AnnAGNPS global & local true (T), false (F), or blank fields.

DATABASE, TEXT, & VERIFICATION FILES

All output files are ASCII formatted. Some are meant to be used with database managers and use a standard comma separated variable format (*.csv). Others are heavily formatted with column headings, page & line counters and are meant to be viewed and interpreted using a text editor. Some are meant for the program coders and scientist to verify and analysis internal calculations for purposes of verification & validation (*.dpp, *.npt, & *.sim). Some of the output restrictions have certain rules and all have defaults.

The user will be able to restrict loadings—water, sediment, nutrients, & pesticides—in the verification and formatted text files. Further restrictions will be allowed to reduce the cell/reach(s) included in these output files. This will be done by allowing the user to explicitly indicate which: (1) reaches, in addition to the "outlet", will be included as a reference location for sediment tracking; (2) cells will be included, but the default will be all cells; (3) subareas will be included for those cell/reach IDs that are all numeric such as created by TopAGNPS'; (4) event outputs will be included by specifying a minimum/maximum dates that the event must fall within to be included in the event output; and (5) the minimum event runoff at the outlet before this particular event is included in any event output.

REACH RULES

The reach default is for only the outlet to be included in the output files. If additional reaches are indicated, they will be in included along with the outlet which will always be included.

CELL RULES

The cell default is for all cells to be included in the output files. To select specific cells, first set the global flag to false and then select the desired cells for output.

SUBAREA RULES

The subarea default is for all subareas to be included in the formatted & verification output files. The subarea restriction is by a minimum & maximum integer subarea ID. Any integer subarea IDs that are included within the minimum to maximum integer number, and any non-numeric IDs, will be included in the output files. A further detail is that for integer cell/reach IDs within any included subarea only, the subarea output can be restricted to include only source cells, left bank cells, right bank cells, or reaches. That is, the subarea output can be restricted to eliminate integer subareas outside of the min/max integer specification and, unless the default is left active, all cells & reaches within the specified integer subareas not specifically requested.

MINIMUM/MAXIMUM EVENT DATE RULES

The minimum/maximum event date default is for all events between the entire simulation period.

MINIMUM EVENT RUNOFF RULES

The minimum event runoff default is 1/4 inch at the outlet.

DATABASE (*.CSV) FILES

Standard-formatted database (*.csv) files will contain complete input and all output-generated event-related erosion, yield, & instream loading cell/reach data because the database manager(s) used will have their own macros to perform arithmetical operations, extract, & produce hard copies for reports and subsidiary files that will be used with other software such as ArcView.

TEXT (*.TXT) FILES

Text (*.txt) files are designed tables that have been hard-coded in Fortran with fixed formats that show column & row headings & dimension units associated with the output statistics. These files contain output of interest to the normal, non-model development user who is performing the analyses. They are designed to be easily read and viewed by text editors and/or MSWORD.

VERIFICATION (*.DPP, *.NPT, & *.SIM) FILES

Verification files (*.dpp, *.npt, & *.sim) are designed to be used by scientists and programmers to verify & validation the model. These files may also be used to calibrate the input.

Not all processes are available to verify the simulation output yet and the ease of reading any particular currently available verification file varies depending upon the effort of the original coder and any modifications subsequently made. The specific format within any one verification file may change as needs require.

Output File Names & Indices

The file names follow a formal, consistent convention to ensure a logical listing order in their directory. Every output file from AnnAGNPS begins with the "AnnAGNPS" and followed by their type of output grouping—database, formatted, input, preparation, & simulation. The extensions indicate their function. The "*.csv" files are designed to be used as standard database files that can be read by most database managers. All other extensions are for user-friendly, hard copy, output tables that are to be read by text. The verification files are designed to be used by the more experienced users (usually program developers—scientists and/or programmers) to analyze internal calculations. These files can also be used to calibrate input.